

THE STABLE HOMOTOPY CATEGORY HAS A UNIQUE MODEL AT THE PRIME 2

STEFAN SCHWEDE

1. INTRODUCTION

The stable homotopy category has been extensively studied by algebraic topologists for a long time. For many applications it is convenient or even necessary to work with point set level models of spectra as opposed to working up-to-homotopy, and the outcome of a calculation can depend on the choice of model. In recent years many new models for the stable homotopy category have been constructed. It is especially useful to have the structure of a *closed model category* in the sense of Quillen [Qui] and many examples of spectra categories fit into this context [BF, Rob87, EKMM, HSS, Lyd, MMSS]. Moreover all known examples capture the ‘same homotopy theory’ – in technical terms one speaks of *Quillen equivalent* model categories [Hov, Def. 1.3.12]. Hence not only the homotopy categories, but also higher order information such as Toda brackets, homotopy colimits and homotopy types of function spaces coincide. In two Quillen equivalent model categories the answer to every homotopy theoretic question comes out the same.

In a model category one can pass to the associated *homotopy category* by formally inverting the class of weak equivalences. However, passage to the homotopy category loses information and in general the ‘homotopy theory’ can not be recovered from the homotopy category, see 2.1 and 2.2 for two examples. In this paper we show that in contrast to the general case, the stable homotopy category completely determines the stable homotopy theory, at least 2-locally. We prove a uniqueness theorem which says that there is essentially only one model category structure underlying the stable homotopy category of 2-local spectra — the stable homotopy category has no ‘exotic’ models at the prime 2.

We call a pointed model category *stable* if it is cocomplete and the loop and suspension functors defined on its homotopy category are inverse equivalences. The homotopy category of a stable model category is naturally triangulated with suspension and cofibration sequences defining the shift operator and the distinguished triangles [Hov, Prop. 7.1.6].

Main Theorem: *Let \mathcal{C} be a stable model category. If the homotopy category of \mathcal{C} and the 2-local homotopy category of spectra are equivalent as triangulated categories, then there exists a Quillen equivalence between \mathcal{C} and the 2-local model category of spectra.*

We prove a stronger form of the main theorem as Theorem 3.5 below. The stronger version says that already the subcategory of *finite* 2-local spectra determines the model category structure up to Quillen equivalence of model categories. In particular there is only one way to ‘complete’ the homotopy category of finite 2-local spectra to a triangulated category with infinite coproducts — as long as some underlying model structure exists. This gives a partial answer to Margolis’ Uniqueness Conjecture [Mar, Ch. 2 §1] for the stable homotopy category, see Corollary 3.7.

The proof of the main theorem relies on the following characterization of self-equivalences of the 2-local stable homotopy category:

Theorem: *Let F be an exact endofunctor of the homotopy category of finite 2-local spectra. If F takes the 2-local sphere spectrum to itself (up to isomorphism), then F is a self-equivalence.*

This result is a combination of Propositions 3.1 and 3.2 below. To obtain the same conclusion at an odd prime, one has to assume in addition that the functor F does not annihilate the first p -torsion element in the stable homotopy groups of spheres, see Proposition 3.1 for the precise statement. The reason that the prime

2 behaves differently from the odd primes goes back to the ‘misbehavior’ of the mod-2 Moore spectrum that its identity map has order 4. In other parts of stable homotopy theory this is often a nuisance; for us it is the key to why we can prove the uniqueness theorem for 2-local spectra. Currently we have no replacement for this part of the argument at odd primes, see also Remark 5.1.

Our reference model is the category of spectra in the sense of Bousfield and Friedlander [BF, §2]. This is probably the simplest model category of spectra and its objects are sequences $\{X_n\}_{n \geq 0}$ of pointed simplicial sets together with maps $\Sigma X_n \rightarrow X_{n+1}$. Morphisms are given on every level and commute strictly with the structure maps. The weak equivalences are the *stable equivalences*, i.e., the morphisms which induce isomorphisms of stable homotopy groups. For the remaining details of the model structure see [BF, Thm. 2.3]. A variant of the stable model structure is the *p-local* model structure for a prime p ; here the weak equivalences are those morphisms which induce an isomorphism on stable homotopy groups tensored with $\mathbb{Z}_{(p)}$; see [SSa, 4.1] for details on the *p-local* model structure.

Acknowledgments: This paper owes a lot to several discussions with Mark Mahowald. He first explained to me why every element in the homotopy groups of spheres is a ‘higher order Toda bracket’ of Adams filtration one elements; this is now a main ingredient of the inductive argument in Lemma 4.1.

2. BACKGROUND AND RELATED RESULTS

2.1. A triangulated category with several models. In general, the triangulated homotopy category does not determine the Quillen equivalence type of a model category. As an example we consider the n -th Morava K -theory spectrum $K(n)$ for a fixed prime and some number $n > 0$. By a theorem of Robinson [Rob89] this spectrum admits the structure of an A_∞ -ring spectrum and so its module spectra form a stable model category. The ring of homotopy groups of $K(n)$ is the graded field $\mathbb{F}_p[v_n, v_n^{-1}]$ with v_n of dimension $2p^n - 2$. Hence the homotopy group functor establishes an equivalence between the homotopy category of $K(n)$ -module spectra and the category of graded $\mathbb{F}_p[v_n, v_n^{-1}]$ -modules.

Similarly the homology functor establishes an equivalence between the derived category of the graded field $\mathbb{F}_p[v_n, v_n^{-1}]$ and the category of graded $\mathbb{F}_p[v_n, v_n^{-1}]$ -modules. This derived category arises from a model category structure on differential graded $\mathbb{F}_p[v_n, v_n^{-1}]$ -modules with weak equivalences the quasi-isomorphisms. So the two stable model categories of $K(n)$ -module spectra and DG-modules over $\mathbb{F}_p[v_n, v_n^{-1}]$ have equivalent triangulated homotopy categories. On the other hand they are not Quillen equivalent: if they were, then the homotopy types of the function spaces would agree [DK, Prop. 5.4]. But for DG-modules all function spaces are products of Eilenberg-MacLane spaces, which is not the case for $K(n)$ -modules.

2.2. Franke’s algebraic model for the $E(n)$ -local stable homotopy category. In [Fra], Franke constructs an exotic model for the homotopy category of $E(n)$ -local spectra at a ‘large’ prime. Earlier Bousfield [Bou] had given an algebraic description of the isomorphism classes of K -local spectra at an odd prime. However Bousfield could not determine whether his algebraic model describes the morphisms between the spectra correctly. As one application of a general uniqueness theorem [Fra, Sec. 2.2 Thm. 5], Franke provides an algebraic derived category which is equivalent, as a triangulated category, to the homotopy category of K -local spectra, see [Fra, Sec. 3.1 Thm. 6]. Franke’s uniqueness theorem applies more generally to the homotopy categories of $E(n)$ -local spectra whenever $n^2 + n < 2p - 2$. Hence he obtains an equivalence of triangulated categories between the derived category of an abelian model category and the homotopy category of $E(n)$ -local spectra, see [Fra, Sec. 3.5 Thm. 10]. By the same reasoning as in 2.1, these two kinds of model categories are *not* Quillen equivalent, see also [Fra, Sec. 3.1 Rem. 1].

We currently do not know whether there exist exotic models for the stable homotopy category at an odd prime. Via Proposition 3.1 this problem reduces to a question about the α_1 -map. Franke’s exotic equivalences are relevant for these considerations since the α_1 -map survives K -localization.

2.3. Reduction to ‘exotic sphere spectra’. Suppose that \mathcal{C} is a stable model category and let $\Phi : \mathrm{Ho}(\mathrm{Spectra}) \xrightarrow{\cong} \mathrm{Ho}(\mathcal{C})$ be an equivalence of triangulated categories. Then the image of the sphere spectrum $P = \Phi(S^0)$ is a small weak generator (see [SSa, Def. 3.1]) of the homotopy category of \mathcal{C} . In [SSb], B. Shipley and the author associate to an object P of a stable model category a ring spectrum $\mathrm{End}_{\mathcal{C}}(P)$ called the *endomorphism ring spectrum*. The ring of homotopy groups of $\mathrm{End}_{\mathcal{C}}(P)$ is isomorphic

to the graded ring of self-maps of P in the homotopy category of \mathcal{C} . If P is a small weak generator then we also show that the model category \mathcal{C} is Quillen equivalent to the category of modules over the endomorphism ring spectrum $\text{End}_{\mathcal{C}}(P)$.

The original equivalence Φ establishes an isomorphism between the ring π_*^s of stable homotopy groups of spheres and the graded ring of self-maps of P in $\text{Ho}(\mathcal{C})$; this in turn is isomorphic to the homotopy groups of the ring spectrum $\text{End}_{\mathcal{C}}(P)$. Since Φ is an exact functor, the isomorphism between π_*^s and $\pi_* \text{End}_{\mathcal{C}}(P)$ also preserves Toda brackets. Hence the endomorphism ring spectrum $\text{End}_{\mathcal{C}}(P)$ looks very much like the sphere spectrum, and the original model category \mathcal{C} is Quillen equivalent to the category of spectra if and only if the unit map $S^0 \rightarrow \text{End}_{\mathcal{C}}(P)$ of the endomorphism ring spectrum is a stable equivalence. In other words: the question whether there are exotic models for the stable homotopy category can be reduced to the question about the existence of ‘exotic sphere spectra’, i.e., ring spectra which are not equivalent to the sphere spectrum, but whose derived category is equivalent to the stable homotopy category. While this reduction gives a better idea of what possible exotic models look like, we will not use the results of [SSb] here and rather prove our uniqueness theorem directly.

2.4. An integral uniqueness result assuming additional structure. The homotopy category of a stable model category admits additional structure which one can take into account when proving a uniqueness result. The homotopy category of every model category admits an action of the homotopy category of simplicial sets [Hov, Thm. 4.3.4]. If the model category is stable, then this action induces an action of the graded ring π_*^s of stable homotopy groups of spheres, see [SSa, 2.3].

In [SSa] B. Shipley and the author show that with this extra structure the stable homotopy category determines the model category structure up to Quillen equivalence. More precisely we show that if \mathcal{C} is a stable model category and if the homotopy category of \mathcal{C} admits a π_*^s -linear equivalence to the homotopy category of spectra, then \mathcal{C} is Quillen equivalent to the Bousfield-Friedlander stable model category of spectra. Hence the result of the present paper is a strengthening of the Uniqueness Theorem of [SSa], at least 2-locally. While [SSa] mainly depends on model category arguments, we have to use more information about the structure of the stable homotopy category here.

3. PROOF OF THE 2-LOCAL UNIQUENESS THEOREM

In this section we deduce our main theorem from other results which should be of independent interest. The first two results concern properties of the stable homotopy category. Proposition 3.1 is an elaboration on the idea that the stable homotopy groups of spheres are generated under ‘higher order Toda brackets’ by the elements of Adams filtration one (see [Coh] for a precise formulation of this fact). For a prime p the *mod- p Adams filtration* of a map of spectra is the largest number n such that the map can be factored as a composite of n maps all of which induce the trivial map on mod- p cohomology. When the prime is understood we simply speak of the filtration of a map. Adams showed [Ada] that the only positive dimensional elements in $\pi_* S_{(2)}^0$ which have filtration one are multiples of the Hopf maps η, ν and σ in dimensions 1, 3 and 7 respectively. For odd primes the only such elements are in the first non-trivial p -torsion homotopy group $\pi_{2p-3} S_{(p)}^0$, see [Liu, Thm. 1.2.1].

An *exact functor* between triangulated categories is an additive functor F which commutes with shift and preserves distinguished triangles. More precisely: F is endowed with a natural isomorphism $\iota_X : F(X[1]) \cong F(X)[1]$ such that for every distinguished triangle (homotopy cofibre sequence)

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} X[1]$$

the sequence

$$F(X) \xrightarrow{F(f)} F(Y) \xrightarrow{F(g)} F(Z) \xrightarrow{\iota_X \circ F(h)} F(X)[1]$$

is again a distinguished triangle. An *equivalence of triangulated categories* is an equivalence of categories which is exact and whose inverse functor is also exact. In what follows square brackets $[-, -]$ denote morphisms in the homotopy category of spectra, possibly graded when decorated with a subscript.

Proposition 3.1. *Let p be a prime number and let F be an exact endofunctor of the homotopy category of finite p -local spectra which takes the p -local sphere spectrum to itself (up to isomorphism). If every element of Adams filtration one in the graded endomorphism ring $[F(S_{(p)}^0), F(S_{(p)}^0)]_*$ is in the image of F , then F is a self-equivalence.*

The next result says that the prime 2 is special because the Hopf maps are always taken care of. We do not know whether the analogue of the following proposition is true for odd primes, see also Remark 5.1.

Proposition 3.2. *Let F be an exact endofunctor of the homotopy category of finite 2-local spectra which takes the 2-local sphere spectrum to itself (up to isomorphism). Then all maps of Adams filtration one in the graded endomorphism ring $[F(S_{(2)}^0), F(S_{(2)}^0)]_*$ are in the image of F .*

We prove Propositions 3.1 and 3.2 in Sections 4 and 5 respectively.

In order to state the next auxiliary result we recall the notion of a *compactly generated* triangulated category. An object A of a triangulated category \mathcal{T} is called *compact* (also called *small* or *finite*) if the representable functor $\mathcal{T}(A, -)$ preserves infinite coproducts. A full subcategory \mathcal{S} of a triangulated category \mathcal{T} is *closed under extensions* if whenever two of the objects X, Y and Z in a distinguished triangle

$$X \longrightarrow Y \longrightarrow Z \longrightarrow X[1]$$

belong to \mathcal{S} , the third object also belongs to \mathcal{S} (this implies that \mathcal{S} contains the zero objects and is closed under isomorphisms, finite sums and shift in either direction). A triangulated category \mathcal{T} is *compactly generated* if \mathcal{T} is the only subcategory which contains the compact objects and is closed under extensions and infinite coproducts. The compact objects of the stable homotopy category are precisely the finite spectra, and similarly for the p -local stable homotopy category. The stable homotopy category and its full subcategory of p -local spectra are compactly generated. However, there are triangulated categories in which the zero objects are the only small objects — see [HS, Cor. B.13] for examples which are Bousfield localizations of the stable homotopy category.

The following lemma is well-known and we give the easy proof at the end of Section 4.

Lemma 3.3. *Let F be an exact functor between compactly generated triangulated categories with infinite coproducts. If F preserves coproducts and restricts to an equivalence between the full subcategories of compact objects, then F is an equivalence.*

Finally, we quote a result from [SSa] which is entirely model category theoretic. It roughly says that the model category of spectra is the ‘free stable model category on one object’. Here spectra are understood in the sense of Bousfield and Friedlander, endowed with the stable model structure of [BF, 2.3]. In particular the weak equivalences are those maps which induce isomorphisms of stable homotopy groups. The p -local model structure is the localization of the stable model structure of spectra in which the weak equivalences are the maps inducing an isomorphism of stable homotopy groups tensored with $\mathbb{Z}_{(p)}$, see [SSa, 4.1] for details. A *left Quillen functor* is a functor between model categories which has a right adjoint and which preserves cofibrations and acyclic cofibrations.

Proposition 3.4. [SSa, 5.1] *Let \mathcal{C} be a stable model category and X a cofibrant and fibrant object of \mathcal{C} . Then there exists a left Quillen functor from the category of spectra to \mathcal{C} which takes the sphere spectrum to X . If the endomorphism ring of X in the homotopy category of \mathcal{C} is a $\mathbb{Z}_{(p)}$ -algebra, then the functor is also a left Quillen functor with respect to the p -local stable model structure for spectra.*

Now we can state and prove a uniqueness result which has the Main Theorem of the introduction as a special case. This version is stronger since the hypothesis only refer to the full subcategory of compact, or finite, objects in the triangulated homotopy category.

Theorem 3.5. *Let \mathcal{C} be a stable model category whose homotopy category is compactly generated. Suppose that the full subcategory of compact objects in the homotopy category of \mathcal{C} and the homotopy category of finite 2-local spectra are equivalent as triangulated categories. Then there exists a Quillen equivalence between \mathcal{C} and the 2-local model category of spectra, whose left adjoint ends in \mathcal{C} .*

Proof. Let Φ be an equivalence of triangulated categories from the homotopy category of finite 2-local spectra to the compact objects in the homotopy category of \mathcal{C} . We choose a cofibrant and fibrant object X of \mathcal{C} which is isomorphic to $\Phi(S_{(2)}^0)$ in the homotopy category of \mathcal{C} . Lemma 3.4 yields a left Quillen functor, with respect to the 2-local stable model structure, from the category of spectra to \mathcal{C} which takes the sphere spectrum to X . We denote the functor by $X \wedge -$. This left Quillen functor has an exact total left derived functor $X \wedge^L -$ on the level of homotopy categories (see [Qui, I.4 Prop. 2] or [Hov, Prop. 6.4.1]).

The derived functor $X \wedge^L -$ takes the localized sphere spectrum to the compact object X , hence it takes compact objects to compact objects; we denote by $(X \wedge^L -)|_{\text{small}}$ the restriction to finite 2-local spectra. The composite functor $F = \Phi^{-1} \circ (X \wedge^L -)|_{\text{small}}$ takes the 2-local sphere spectrum to itself, up to isomorphism, so by Propositions 3.1 and 3.2 it is a self-equivalence of the finite 2-local stable homotopy category. Since F and Φ^{-1} are equivalences of categories, so is $(X \wedge^L -)|_{\text{small}}$. By Lemma 3.3, the functor $X \wedge^L -$ is then an equivalence of categories, so the left Quillen functor $X \wedge -$ and its right adjoint are in fact a Quillen equivalence [Hov, Prop. 1.3.13]. \square

Warning: The equivalence Φ takes the 2-local sphere spectrum to the object X , and the same is true for the left derived functor $X \wedge^L -$. If there was a natural transformation between Φ and $(X \wedge^L -)|_{\text{small}}$ which induces an isomorphism at $S_{(2)}^0$, then the natural transformation would be a natural isomorphism, so $X \wedge^L -$ would also be an equivalence on compact objects. However, there is no reason why such a natural transformation should exist, and so there is no a priori reason why $X \wedge^L -$ should be an equivalence.

In particular we do *not* claim that the left Quillen equivalence $X \wedge -$ lifts the triangulated equivalence Φ . Hence we leave open the question whether there are exotic self-equivalences of the 2-local stable homotopy category, i.e., self-equivalences not induced from a Quillen equivalence (or what is the same: self-equivalences other than iterated (de-)suspensions).

Remark 3.6. (Margolis' Uniqueness Conjecture) In 'Spectra and the Steenrod algebra', H. R. Margolis introduces a set of axioms for a *stable homotopy category* [Mar, Ch. 2 §1]. The stable homotopy category of spectra satisfies the axioms, and Margolis conjectures [Mar, Ch. 2, §1] that this is the only model, i.e., that any category which satisfies the axioms is equivalent to the stable homotopy category.

As part of the structure Margolis requires a triangulation, infinite coproducts and that the whole category be generated by a single compact object. Furthermore, Margolis' Axiom 5 asks for an equivalence between the full subcategory of compact objects and the Spanier-Whitehead category of finite CW-complexes. So the Uniqueness Conjecture really concerns possible 'completions' of the category of finite spectra to a triangulated category with infinite coproducts. Margolis also assumes the existence of a compatible symmetric monoidal smash product, but the smash product does not enter into our present considerations.

Margolis shows [Mar, Ch. 5 Thm. 19] that modulo phantom maps each model of his axioms is equivalent to the standard model. Moreover, Christensen and Strickland show in [CS] that in any model the ideal of phantom maps is equivalent to the phantoms in the standard model.

One can consider a 2-primary analog of Margolis' Uniqueness Conjecture by modifying his Axiom 5 and instead requiring the full subcategory of compact objects in the stable homotopy category to be equivalent, as a triangulated category, to the homotopy category of finite 2-local spectra. Our main theorem proves the following 2-primary analog of Margolis' Uniqueness Conjecture for stable homotopy categories with a model:

Corollary 3.7. *Suppose that \mathcal{S} is a 2-primary stable homotopy category (in the sense of [Mar, Ch. 2 §1]) which is equivalent, as a triangulated category, to the homotopy category of a stable model category. Then \mathcal{S} is triangulated equivalent to the homotopy category of 2-local spectra.*

Note that we do not assume any internal smash product on the model category, and the corollary does not give that the equivalence between \mathcal{S} and the stable homotopy category of 2-local spectra preserves the smash products.

4. A CHARACTERIZATION OF SELF-EQUIVALENCES OF THE STABLE HOMOTOPY CATEGORY

In this section we prove Proposition 3.1. Throughout, p denotes any prime and F is an exact endofunctor of the homotopy category of finite p -local spectra. We assume further that F takes the p -local sphere spectrum to itself (up to isomorphism) and that all filtration one maps of positive dimension from $F(S_{(p)}^0)$ to itself are in the image of F . We want to show that F is then a self-equivalence.

Cohomology will always be spectrum cohomology with mod- p coefficients. If K is a finite spectrum we denote by $\beta(K)$ (resp. $\tau(K)$) the smallest (resp. largest) dimension in which the mod- p cohomology of K is non-trivial. As before square brackets $[-, -]$ denote morphisms in the homotopy category of spectra, possibly graded when decorated with a subscript.

Lemma 4.1. *Suppose that the map of graded rings $[S_{(p)}^0, S_{(p)}^0]_* \rightarrow [F(S_{(p)}^0), F(S_{(p)}^0)]_*$ induced by the functor F is an isomorphism below and including dimension n for some $n \geq 0$.*

- (1) *Let K and L be two finite p -local spectra. Then the map $F : [K, L] \rightarrow [F(K), F(L)]$ is an isomorphism if $\tau(K) - \beta(L) < n$ and an epimorphism if $\tau(K) - \beta(L) = n$.*
- (2) *Let K be a finite p -local spectrum satisfying $\tau(K) - \beta(K) \leq n + 1$. Then there exists a finite p -local spectrum K' with $\beta(K') \geq \beta(K)$ and $\tau(K') \leq \tau(K)$ and an isomorphism $K \cong F(K')$ in the homotopy category of spectra.*
- (3) *Every map from $F(S_{(p)}^{n+1})$ to $F(S_{(p)}^0)$ of Adams filtration at least two is in the image of F .*

Proof. (1) When K and L are localized sphere spectra, the claim hold by assumption. The general case is obtained by cell inductions for K and L .

We first prove the claim when L is a wedge of localized sphere spectra of a fixed dimension using induction on the total dimension of the mod- p cohomology of K . If H^*K is trivial, then K is contractible and the statement is true. Otherwise we can pinch off the top cells of K , i.e., we can choose a distinguished triangle

$$\bigvee_I S^{\tau(K)-1} \xrightarrow{\alpha} M \longrightarrow K \longrightarrow \bigvee_I S^{\tau(K)} \quad (*)$$

with M a finite p -local spectrum with $\tau(M) < \tau(K)$ and with strictly smaller cohomology. Hence by induction the claim is true for M and L . Taking homomorphism groups $[-, L]$ from the triangle $(*)$ gives a long exact sequence of abelian groups. The functor F preserves distinguished triangle, and so taking homomorphism from the image sequence into $F(L)$ yields a similar exact sequence and F gives a map between the sequences. Using that L is a wedge of spheres and that the claim holds for M with estimate increased by one, the five lemma proves the statement for K and this special L .

Now we do a similar induction on the dimension of H^*L . This time we collapse the bottom cells of L , i.e., we embed L in a triangle

$$\bigvee_J S^{\beta(L)} \longrightarrow L \longrightarrow L' \longrightarrow \bigvee_J S^{\beta(L)+1}$$

where the dimension of H^*L' is strictly smaller than that of L and $\beta(L') > \beta(L)$. By induction the claim holds for the spectra K and L' , and using the five lemma and the previous paragraph we deduce it for K and L .

(2) We argue by induction on the difference $\tau(K) - \beta(K)$. If the cohomology of K is concentrated in at most one dimension, then K is equivalent to a (possibly empty) wedge of localized spheres and the statement is true. Otherwise there exists a distinguished triangle $(*)$ as in part (1) with M a finite p -local spectrum which satisfies $\tau(M) < \tau(K)$ and $\beta(M) = \beta(K)$. By induction there exists a finite spectrum M' with $\beta(M') \geq \beta(M)$, $\tau(M') \leq \tau(M)$ and an isomorphism between $F(M')$ and M . By part (1) the composite

$$F(\bigvee_I S^{\tau(K)-1}) \xrightarrow{\cong} \bigvee_I S^{\tau(K)-1} \xrightarrow{\alpha} M \xrightarrow{\cong} F(M')$$

is of the form $F(\alpha')$ for some $\alpha' \in [\bigvee_I S^{\tau(K)-1}, M']$. We let K' be some mapping cone of the map α' , obtained by embedding α' in a distinguished triangle. Then we have the inequalities $\beta(K') \geq \beta(M') \geq$

$\beta(M) = \beta(K)$ and $\tau(K') \leq \max\{\tau(M'), \tau(K)\} = \tau(K)$. We end up with a diagram

$$\begin{array}{ccccccc}
 \bigvee_I S^{\tau(K)-1} & \xrightarrow{\alpha} & M & \xrightarrow{\quad} & K & \xrightarrow{\quad} & \bigvee_I S^{\tau(K)} \\
 \downarrow \cong & & \downarrow \cong & & \downarrow \text{dashed} & & \downarrow \cong \\
 F(\bigvee_I S^{\tau(K)-1}) & \xrightarrow{F(\alpha')} & F(M') & \xrightarrow{\quad} & F(K') & \xrightarrow{\quad} & F(\bigvee_I S^{\tau(K)})
 \end{array}$$

in which both rows are distinguished triangles and the left square commutes. Hence we can choose a map $K \rightarrow F(K')$ which makes the entire diagram commute, and this map is the isomorphism we are looking for.

(3) We first claim that every map $F(S_{(p)}^{n+1}) \rightarrow F(S_{(p)}^0)$ of filtration at least two can be factored through a spectrum of the form $F(K')$ where K' is a finite p -local spectrum with cohomology concentrated in dimensions 1 through n . Both maps in such a factorization are in the image of F by part (1), hence the original map is also in the image.

Except for the use of the functor F the claim is a well-known argument – it is e.g. used by Cohen [Coh, Thm. 4.2] who attributes it to Adams. We choose an isomorphism between $F(S_{(p)}^0)$ and $S_{(p)}^0$ and denote by $\overline{H\mathbb{Z}}_{(p)}$ and $\overline{H\mathbb{F}}_p$ the fibers of the composite Hurewicz maps

$$F(S_{(p)}^0) \xrightarrow{\cong} S_{(p)}^0 \xrightarrow{\text{Hurewicz}} H\mathbb{Z}_{(p)} \quad \text{and} \quad F(S_{(p)}^0) \xrightarrow{\cong} S_{(p)}^0 \xrightarrow{\text{Hurewicz}} H\mathbb{F}_p$$

to the p -local and mod- p Eilenberg-MacLane spectra respectively. A map to $F(S_{(p)}^0)$ has Adams filtration two if and only if it lifts to the fiber $\overline{H\mathbb{F}}_p$ and if some (hence any) such lift is trivial in mod- p cohomology.

Now let $\alpha : F(S_{(p)}^{n+1}) \rightarrow F(S_{(p)}^0)$ be a map of filtration at least two. Since $n+1 > 0$, α lifts to a map $F(S_{(p)}^{n+1}) \rightarrow \overline{H\mathbb{Z}}_{(p)}$. Furthermore the lift induces the trivial map in mod- p cohomology since the map $\overline{H\mathbb{Z}}_{(p)} \rightarrow \overline{H\mathbb{F}}_p$ is surjective in mod- p cohomology.

Since $F(S_{(p)}^{n+1})$ is an $(n+1)$ -dimensional p -local sphere, the lift of α factors through the $(n+1)$ -dimensional skeleton of the spectrum $\overline{H\mathbb{Z}}_{(p)}$. Since in addition α is trivial in mod- p cohomology and the cohomology of $\overline{H\mathbb{Z}}_{(p)}$ is acyclic with respect to the Bockstein operator, the lift can even be factored through the n -skeleton of $\overline{H\mathbb{Z}}_{(p)}$. This n -skeleton is a finite p -local spectrum with mod- p cohomology concentrated in dimensions 1 through n . By part (2) of this lemma there exists a finite p -local spectrum K' with cohomology in dimensions 1 through n and an isomorphism between the n -skeleton of $\overline{H\mathbb{Z}}_{(p)}$ and $F(K')$, which proves the claim. \square

Now we can give the

Proof of Proposition 3.1. Suppose F is an exact endofunctor of the homotopy category of finite p -local spectra which takes the p -local sphere spectrum to itself, up to isomorphism. Furthermore all filtration one maps of positive dimension from $F(S_{(p)}^0)$ to itself are in the image of F . We claim that the map of graded rings $F : [S_{(p)}^0, S_{(p)}^0]_* \rightarrow [F(S_{(p)}^0), F(S_{(p)}^0)]_*$ is an isomorphism. Since $F(S_{(p)}^0)$ is isomorphic to $S_{(p)}^0$ the map is necessarily an isomorphism in non-negative dimensions, and in positive dimensions both sides of the map are finite groups of the same order. Suppose the claim was false and let $m > 0$ be the smallest dimension in which F is not bijective, hence not surjective. By Lemma 4.1 (3), any element not in the image has filtration one, which contradicts the assumptions. Hence the above map is indeed bijective in all dimensions.

So the hypothesis of Lemma 4.1 is satisfied for arbitrarily large n ; conclusion (1) of that Lemma shows that F is full and faithful and conclusion (2) shows that F is surjective on isomorphism classes. Thus F is an equivalence of categories. \square

It remains to prove Lemma 3.3 which allows us to detect an equivalence of triangulated categories on compact objects.

Proof of Lemma 3.3. Let $F : \mathcal{S} \longrightarrow \mathcal{T}$ be an exact functor between compactly generated triangulated categories with infinite coproducts. Suppose that F preserves coproducts and restricts to an equivalence between the full subcategories of compact objects. We want to show that F itself is an equivalence.

We fix a compact object A of \mathcal{S} and consider the full subcategory of \mathcal{S} consisting of those Y for which the map

$$F : \mathcal{S}(A, Y) \longrightarrow \mathcal{T}(F(A), F(Y))$$

is bijective. By assumption this subcategory contains all compact objects. Since F is exact, the subcategory is closed under extensions. Since A and $F(A)$ are compact and F preserves coproducts, this subcategory is also closed under coproducts. Since \mathcal{S} is compactly generated, the map $F : \mathcal{S}(A, Y) \longrightarrow \mathcal{T}(F(A), F(Y))$ is thus bijective for all compact A and arbitrary Y .

Similarly for arbitrary but fixed Y the full subcategory of \mathcal{S} consisting of those X for which the map $F : \mathcal{S}(X, Y) \longrightarrow \mathcal{T}(F(X), F(Y))$ is bijective is closed under extensions, coproducts and contains the compact objects. Hence this subcategory coincides with \mathcal{S} which means that F is full and faithful.

Now we consider the full subcategory of \mathcal{T} of objects which are isomorphic to an object in the image of F . By assumption this category contains all compact objects, and it is closed under coproducts since these are preserved by F . We claim that this subcategory is also closed under extensions. Since \mathcal{T} is compactly generated this shows that F is essentially surjective and hence an equivalence.

To prove the last claim we consider a distinguished triangle

$$X \xrightarrow{f} Y \longrightarrow Z \longrightarrow X[1].$$

Since the subcategory under consideration is closed under isomorphism and shift in either direction we can assume that $X = F(X')$ and $Y = F(Y')$ are objects in the image of F . Since F is full there exists a map $f' : X' \longrightarrow Y'$ satisfying $F(f') = f$. We can then choose a mapping cone for the map f' and a compatible map from Z to $F(\text{Cone}(f'))$ which is necessarily an isomorphism. \square

5. TAKING CARE OF THE HOPF MAPS

In this final section we prove Proposition 3.2. Here F denotes an exact endofunctor of the homotopy category of 2-local spectra which takes the 2-local sphere spectrum to itself (up to isomorphism). We want to show that all maps of Adams filtration one in the graded endomorphism ring $[F(S_{(2)}^0), F(S_{(2)}^0)]_*$ are in the image of F . We introduce a slight abuse of notation: after choosing an isomorphism between $F(S_{(2)}^0)$ and the 2-local sphere spectrum we identify the ring $[F(S_{(2)}^0), F(S_{(2)}^0)]_*$ with the ring of 2-local stable homotopy groups of spheres (this identification does not depend on the choice of isomorphism). With this convention we have to show that the Hopf maps η, ν and σ are in the image of F in dimensions 1, 3 and 7 respectively.

We start by showing that the map $F(\eta)$ is non-trivial. Since F is exact, both rows in the diagram

$$\begin{array}{ccccccc} F(S_{(2)}^0) & \xrightarrow{\times 2} & F(S_{(2)}^0) & \longrightarrow & F(M(2)) & \longrightarrow & F(S_{(2)}^1) \\ \cong \downarrow & & \downarrow \cong & & \vdots & & \downarrow \cong \\ S_{(2)}^0 & \xrightarrow{\times 2} & S_{(2)}^0 & \longrightarrow & M(2) & \longrightarrow & S_{(2)}^1 \end{array}$$

are distinguished triangles (here $M(2)$ denotes the mod-2 Moore spectrum). Since the left square commutes we can choose a map between $F(M(2))$ and $M(2)$ making the entire diagram commute, and this map is necessarily an isomorphism.

The identity map of the Moore spectrum $M(2)$ has additive order 4. Since $F(M(2))$ is isomorphic to $M(2)$, its identity map also has order 4. Since F is additive, it does not annihilate the degree 2 map of the Moore spectrum. On the other hand this degree 2 map factors as the composite

$$M(2) \xrightarrow{\text{pinch}} S^1 \xrightarrow{\eta} S^0 \xrightarrow{\text{incl.}} M(2) .$$

Hence $F(\eta)$ has to be non-zero, which forces $F(\eta) = \eta$.

Because of the relation $4\nu = \eta^3$ (see e.g. [Tod, Theorem 14.1 (i)]) we know that $4 \cdot F(\nu) = F(4\nu) = F(\eta^3) = \eta^3 = 4\nu$. Since ν generates the cyclic group $\pi_3 S_{(2)}^0 \cong \mathbb{Z}/8$, we conclude that $F(\nu) = u \cdot \nu$ with u a 2-local unit. Hence ν is in the image of F .

For the last Hopf map σ we exploit the Toda bracket relation $4\sigma = \langle \nu, 8\iota, \nu \rangle$ (see e.g. [Tod, Lemmas 5.13 and 5.14]) which holds without indeterminacy since the fourth stable homotopy group of spheres is trivial. Since F preserves triangles it takes threefold Toda brackets to threefold Toda brackets and we obtain

$$4 \cdot F(\sigma) = F(4\sigma) = F(\langle \nu, 8\iota, \nu \rangle) \subseteq \langle F(\nu), F(8\iota), F(\nu) \rangle = u^2 \cdot \langle \nu, 8\iota, \nu \rangle = 4u^2 \cdot \sigma .$$

Since the latter Toda-bracket has no indeterminacy we conclude that $4 \cdot F(\sigma) = 4u^2 \cdot \sigma$. Again since σ generates the cyclic group $\pi_7 S_{(2)}^0 \cong \mathbb{Z}/16$, we conclude that $F(\sigma)$ coincides with σ up to a 2-local unit, so σ is in the image of F . This finishes the proof of Proposition 3.2.

Remark 5.1. The fact that multiplication by 2 is non-trivial on the mod-2 Moore spectrum is equivalent to the fact that $M(2)$ does not admit a product, i.e. there is no map $M(2) \wedge M(2) \rightarrow M(2)$ in the stable homotopy category which splits the two inclusions $j \wedge \text{id}, \text{id} \wedge j : M(2) \rightarrow M(2) \wedge M(2)$, where $j : S^0 \rightarrow M(2)$ is the inclusion of the bottom cell.

For an odd prime p the Moore spectrum $M(p)$ admits a unique and commutative product, which is also associative for $p \geq 5$. Hence $M(p)$ is a ring spectrum in the stable homotopy category. However, the product on $M(p)$ can not be made associative up to ‘coherent higher homotopy’, i.e., $M(p)$ does not admit the structure of an A_∞ -ring spectrum. In fact the element $\alpha_1 \in \pi_{2p-3} S_{(p)}^0$ is the obstruction to p -th order homotopy associativity. One could try to use this to detect α_1 and to prove the odd primary version of Proposition 3.2.

For $p = 3$ the associativity obstruction can be described more concretely, and we explain why we expect that the odd primary uniqueness problem cannot be decided by only looking at products on the Moore spectrum. Let $\mu : M(3) \wedge M(3) \rightarrow M(3)$ denote the unique and commutative multiplication on the mod-3 Moore spectrum. This product is not associative and in fact the associator $\mu \circ (\mu \wedge \text{id}) - \mu \circ (\text{id} \wedge \mu) : M(3) \wedge M(3) \wedge M(3) \rightarrow M(3)$ factors as the composite

$$M(3) \wedge M(3) \wedge M(3) \xrightarrow{\text{pinch}} S^3 \xrightarrow{\alpha_1} S^0 \xrightarrow{j} M(3) .$$

Suppose F is an exact endofunctor of the homotopy category of 3-local spectra which takes the 3-local sphere spectrum to itself. The same argument as for $p = 2$ yields an isomorphism $f : F(M(3)) \rightarrow M(3)$ under the isomorphism $F(S_{(3)}^0) \cong S_{(3)}^0$, which is unique since the first stable homotopy groups of spheres has no 3-torsion.

The multiplication μ splits the inclusion $\text{id} \wedge j : M(3) \rightarrow M(3) \wedge M(3)$, so the smash product of two copies of the mod-3 Moore spectrum is isomorphic to the wedge $M(3) \vee \Sigma M(3)$. Similarly for three smash factors there is an isomorphism

$$M(3) \wedge M(3) \wedge M(3) \xrightarrow{\cong} M(3) \vee \Sigma M(3) \vee \Sigma M(3) \vee \Sigma^2 M(3)$$

which we can choose in such a way that the projection onto the first summand is the map $\mu \circ (\mu \wedge \text{id}) : M(3) \wedge M(3) \wedge M(3) \rightarrow M(3)$. The exact functor F commutes with wedge and suspension, so together with (suspensions of) the isomorphism $f : F(M(3)) \cong M(3)$ we obtain an isomorphism

$$\overline{f} : F(M(3) \wedge M(3) \wedge M(3)) \longrightarrow M(3) \wedge M(3) \wedge M(3)$$

such that the diagram

$$\begin{array}{ccc}
 F(M(3) \wedge M(3) \wedge M(3)) & \xrightarrow{\bar{f}} & M(3) \wedge M(3) \wedge M(3) \\
 \downarrow F(\mu \circ (\mu \wedge \text{id})) & & \downarrow \mu \circ (\mu \wedge \text{id}) \\
 F(M(3)) & \xrightarrow{f} & M(3)
 \end{array}$$

commutes. However, it is not clear at this point if the choices can be made so that the isomorphisms \bar{f} and f also compatible with the map $\mu \circ (\text{id} \wedge \mu) : M(3) \wedge M(3) \wedge M(3) \longrightarrow M(3)$ which gives the other summand in the associator. Hence it is conceivable that the functor F annihilates the associator, and if that happens, then F does not detect α_1 .

REFERENCES

- [Ada] J. F. Adams, *On the non-existence of elements of Hopf invariant one*, Ann. of Math. II. **72** (1960), 20–104.
- [Bou] A. K. Bousfield, *On the homotopy theory of K-local spectra at an odd prime*, Am. J. Math. **107** (1985), 895–932.
- [BF] A. K. Bousfield and E. M. Friedlander, *Homotopy theory of Γ -spaces, spectra, and bisimplicial sets*, Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II (M. G. Barratt and M. E. Mahowald, eds.), Lecture Notes in Math., vol. 658, Springer, Berlin, 1978, pp. 80–130.
- [CS] J. D. Christensen and N. P. Strickland, *Phantom Maps and Homology Theories*, Topology **37** (1998), 339–364.
- [Coh] J. M. Cohen, *The decomposition of stable homotopy*, Ann. of Math. (2) **87** (1968), 305–320.
- [DK] W. G. Dwyer and D. M. Kan, *Function complexes in homotopical algebra*, Topology **19** (1980), 427–440.
- [EKMM] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May, *Rings, modules, and algebras in stable homotopy theory. With an appendix by M. Cole*, Mathematical Surveys and Monographs, vol. 47, American Mathematical Society, Providence, RI, 1997, xii+249 pp.
- [Fra] J. Franke, *Uniqueness theorems for certain triangulated categories possessing an Adams spectral sequence*, K-theory Preprint Archives #139 (1996), <http://www.math.uiuc.edu/K-theory/>.
- [Hov] M. Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, Providence, RI, 1999, xii+209 pp.
- [HSS] M. Hovey, B. Shipley, and J. Smith, *Symmetric spectra*, J. Amer. Math. Soc. **13** (2000), 149–208.
- [HS] M. Hovey and N. P. Strickland, *Morava K-theories and localisation*, Mem. Amer. Math. Soc. **139** (1999)
- [Liu] A. Liulevicius, *The factorization of cyclic reduced powers by secondary cohomology operations*, Mem. Amer. Math. Soc. **42** (1962)
- [Lyd] M. Lydakis, *Simplicial functors and stable homotopy theory*, Preprint, Universität Bielefeld, 1998. <http://hopf.math.purdue.edu/>.
- [MMSS] M. A. Mandell, J. P. May, S. Schwede and B. Shipley, *Model categories of diagram spectra*, Proc. London Math. Soc., to appear.
- [Mar] H. R. Margolis, *Spectra and the Steenrod algebra. Modules over the Steenrod algebra and the stable homotopy category*, North-Holland Mathematical Library **29**, North-Holland Publishing Co., Amsterdam-New York, 1983, xix+489 pp.
- [Qui] Daniel G. Quillen, *Homotopical algebra*, Lecture Notes in Mathematics, vol. 43, Springer-Verlag, 1967.
- [Rob87] A. Robinson, *Spectral sheaves: a model category for stable homotopy theory*, J. Pure Appl. Algebra **45** (1987), 171–200.
- [Rob89] A. Robinson, *Obstruction theory and the strict associativity of Morava K-theories*, Advances in homotopy theory (Cortona, 1988), Cambridge Univ. Press 1989, 143–152.
- [SSa] S. Schwede and B. Shipley, *A uniqueness theorem for stable homotopy theory*, Preprint, 2000. <http://www.mathematik.uni-bielefeld.de/~schwede/>
- [SSb] S. Schwede and B. Shipley, *Classification of stable model categories*, In Preparation.
- [Tod] H. Toda, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies **49**, Princeton University Press, 1962, v+193 pp.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, 33615 BIELEFELD, GERMANY
E-mail address: schwede@mathematik.uni-bielefeld.de