

Three problems for polynomials of small measure

ARTŪRAS DUBICKAS

Abstract. The problems of estimating the multiplicity of a polynomial at a certain complex point, of finding an upper bound for the height of a polynomial of small Mahler measure, as well the problem of finding how close an algebraic number can be to 1 are all related. We discuss these three problems, and show that the progress in each of the three sharpens either lower or upper bounds in the remaining two. In particular, we give an almost complete answer to the multiplicity at -1 problem for polynomials with coefficients equal to zero or one. An explicit example of a polynomial having small Mahler measure, but large norm, is also presented.

1. Introduction. Let $f(x) = \sum_{k=0}^D a_k x^k$ be a nonzero polynomial with complex coefficients. To measure the complexity (“height”) of the polynomial f one may use the notions of the length of f , $L(f) = \sum_{k=0}^D |a_k|$, the “classical” height of f , $H(f) = \max_{0 \leq k \leq D} |a_k|$, the maximum of the modulus of f in the unit circle, $\|f\| = \max_{|z| \leq 1} |f(z)|$, the Mahler measure of f ,

$$M(f) = \exp \left(\int_0^{2\pi} \log |f(e^{it})| dt \right),$$

or say the value of f at unity, $f(1)$.

Set $r_\varrho(f)$ for the order of vanishing of f at a complex number ϱ . Below, we shall use this notation for only two values of ϱ : $\varrho = 1$ and $\varrho = -1$.

A nonzero algebraic number $\alpha \in \overline{\mathbb{Q}}^*$ is described by its unique normalised minimal polynomial $P = P_\alpha$, whose coefficients are relatively prime integers and whose leading coefficient is positive. The minimal polynomial P is irreducible over the field of rational numbers \mathbb{Q} . Let $\mathcal{N} : \overline{\mathbb{Q}}^* \rightarrow \mathbb{Q}$ be the norm function. For every $\alpha \in \overline{\mathbb{Q}}^*$, we have

$$|\mathcal{N}(\alpha - 1)| \leq |P(1)| \leq \|P\| \leq L(P),$$

where $P = P_\alpha$.

We can now state three related problems for polynomials of small Mahler measure.

MULTIPLICITY PROBLEM. For a given set of complex numbers \mathcal{A} , find or estimate

$$r_\varrho(\mathcal{A}, D) = \max r_\varrho(f),$$

2000 *Mathematics Subject Classification*: 11R04, 11R09, 11B83.

where the maximum is taken over all nonzero polynomials f of degree D with coefficients in the set \mathcal{A} .

HEIGHT PROBLEM. Estimate the maximum of some “height” (say $|P(1)|$, $H(P)$, or $L(P)$) over all irreducible integer polynomials P of degree d and of Mahler measure at most M .

APPROXIMATION PROBLEM. Estimate the minimum of $|\alpha + 1|$ over all algebraic numbers of degree d and of Mahler measure at most M .

In the subsequent sections we discuss the problems and give some new results. Some of these (see sections 3 and 4) show that the problems are indeed related, and the progress in each of the three would sharpen either upper or lower bounds in the remaining two.

2. Multiplicity problem. The multiplicity problem for integer polynomials of bounded height, namely, with coefficients in the set

$$\mathcal{A}_H = \{a \in \mathbb{Z} : |a| \leq H\},$$

is an old one. Note that $r_1(\mathcal{A}_H, D) = r_{-1}(\mathcal{A}_H, D)$, since $\mathcal{A}_H = -\mathcal{A}_H$. Various bounds for $r_1(\mathcal{A}_H, D)$ were obtained by Bloch and Pólya [7], Schur [28], Mignotte [23], Bombieri and Vaaler [8], [9], Amoroso [2], Borwein, Erdélyi and Kós [10]. In particular, if $\varepsilon > 0$ for the $\{-1, 0, 1\}$ polynomials we have the bounds

$$(2 - \varepsilon) \sqrt{\log 2 \frac{D}{\log D}} < r_1(\mathcal{A}_1, D) < \frac{21}{13} \sqrt{D}$$

for all D sufficiently large. Here the lower bound follows from the work of Bombieri and Vaaler [8] (see also [18] for a different proof). The upper bound (see [20]) strengthens the constant $16/7$ obtained by Borwein, Erdélyi and Kós [10].

The problem of closing the remaining sublogarithmic gap is almost certainly a difficult one. In particular, the sharpening of the lower bound would have considerable applications in Diophantine approximation. Borwein and Mossinghoff [12] gave some computational results for the degrees of polynomials with coefficients in $\mathcal{A}_1 = \{-1, 0, 1\}$ and prescribed vanishing at 1.

In some cases the magnitude of $r_1(\mathcal{A}, D)$ is considerably smaller. Boyd [13] proved the inequalities

$$\log D \ll r_1(\mathcal{A}^*, D) < (1 + \varepsilon) \frac{(\log D)^2}{\log \log D},$$

where $\varepsilon > 0$, $\mathcal{A}^* = \{-1, 1\}$, and D is a sufficiently large integer. His upper bound is true for every bounded by an absolute constant set of odd integers \mathcal{A}^* .

If $\mathcal{B} = \{0, 1\}$, then $r_1(\mathcal{B}, D) = 0$. For $r_{-1}(\mathcal{B}, D)$ we can give an almost complete answer. Below [...] stands for the integral part.

THEOREM 1. *Let $D \geq 2$ be a positive integer. Then either*

$$r_{-1}(\mathcal{B}, D) = \left\lfloor \frac{\log D}{\log 2} \right\rfloor,$$

or

$$r_{-1}(\mathcal{B}, D) = \left\lfloor \frac{\log D}{\log 2} \right\rfloor - 1.$$

Furthermore, each possibility occur infinitely often.

PROOF. The upper bound

$$r_{-1}(\mathcal{B}, D) \leq \frac{\log(D+1)}{\log 2}$$

is given in [10]. We shall replace $D+1$ by D .

It follows easily that $r_{-1}(\mathcal{B}, 1) = r_{-1}(\mathcal{B}, 2) = 1$. Suppose that $2^r > D > 2$, where $r = r_{-1}(\mathcal{B}, D)$. Let g be a nonzero polynomial with $\{0, 1\}$ coefficients such that $r_{-1}(g) = r$. Clearly,

$$g(x) \neq 1 + x + x^2 + \dots + x^D,$$

since $r > 1$. At least one coefficient of g , therefore, is equal to 0. Hence

$$2^r \leq g(1) \leq D,$$

a contradiction. Since r is an integer, we deduce the inequality

$$r = r_{-1}(\mathcal{B}, D) \leq \left\lfloor \frac{\log D}{\log 2} \right\rfloor.$$

Set $b_k = (2^{k+1} + (-1)^k)/3$. Of course, $\{b_k\}_{k=1,2,\dots}$ is a sequence of positive odd integers. For any positive integer k one can easily verify the inequality

$$b_1 + b_2 + \dots + b_k < b_{k+1}$$

(in fact, the sum on the left hand side is equal to $b_{k+1} - 1$ or to $b_{k+1} - 2$ for k even or odd, respectively). Therefore,

$$B_r(x) = \prod_{k=1}^r (x^{b_k} + 1)$$

is a polynomial with coefficients in the set $\mathcal{B} = \{0, 1\}$. Set now

$$r = \left\lceil \frac{\log(3D/4)}{\log 2} \right\rceil,$$

and

$$B_r^*(x) = x^{D-b_1-b_2-\dots-b_r} \prod_{k=1}^r (x^{b_k} + 1).$$

Since

$$b_1 + \dots + b_r \leq b_{r+1} - 1 < \frac{4}{3}2^r \leq D,$$

$B_r^*(x)$ is a $\{0, 1\}$ polynomial of degree D .

Note that

$$\begin{aligned} r_{-1}(B_r^*) &= r = \left\lceil \frac{\log(3D/4)}{\log 2} \right\rceil \\ &= \left\lceil \frac{\log D}{\log 2} - 0,415\dots \right\rceil \geq \left\lceil \frac{\log D}{\log 2} \right\rceil - 1. \end{aligned}$$

This shows that there are only two possibilities for $r_{-1}(\mathcal{B}, D)$. Furthermore, if

$$\left\{ \frac{\log D}{\log 2} \right\} \geq 2 - \frac{\log 3}{\log 2} = 0.415\dots,$$

where $\{\dots\}$ stands for the fractional part, then we have the first possibility, namely,

$$r_{-1}(\mathcal{B}, D) = r = \left\lceil \frac{\log(3D/4)}{\log 2} \right\rceil = \left\lceil \frac{\log D}{\log 2} \right\rceil.$$

Clearly, there is an infinite number of such integers D .

Suppose now that D is a power of 2: $D = 2^m$, $D \geq 8$. We claim that

$$r_{-1}(\mathcal{B}, 2^m) = m - 1.$$

Indeed, assume that there is a $\{0, 1\}$ polynomial g of degree $D = 2^m$ for which

$$r_{-1}(g) = m.$$

Then, there is a coefficient of g equal to zero, for otherwise $g(-1) \neq 0$. Suppose that there is only one such a coefficient. Since $m \geq 3$, we see that $g(-1) = 0$, $g'(-1) = 0$ and $g''(-1) = 0$. From the first two of these equalities one can easily find that

$$g(x) = \frac{x^{D+1} - 1}{x - 1} - x^{D/2}.$$

But then

$$g''(-1) = \frac{D^2 + 2D}{4} \neq 0,$$

a contradiction. Thus, among the coefficients of g there are at least two zeros. Then, as above,

$$2^m = 2^{r_{-1}(g)} \leq D - 1 < 2^m,$$

a contradiction. Consequently, for every $m \geq 3$ we have the second possibility, namely, $r_{-1}(\mathcal{B}, 2^m) = m - 1$. \blacksquare

Note that Theorem 1 is true for every set $\mathcal{B} = \{0, b\}$, where b is a nonzero complex number, by considering the $\{0, 1\}$ polynomials g/b instead of the $\{0, b\}$ polynomials g .

3. Height problem. For the height and approximation problems we assume that $1 < M < \exp(d^{1-\delta})$, where δ is a small positive constant. Here we use the letter d instead of D for the degree to show that the involved polynomials are irreducible.

Mignotte [24] showed that the inequality

$$L(P) \leq (2d + 1)M \exp \left(\sqrt{2d(4 + \log d) \log((2d + 1)^{1/2} M)} \right)$$

holds for irreducible integer polynomials P of degree d and of measure at most M . For large M , say $M \geq e^d / \sqrt{d}$, the Liouville inequality $L(P) \leq 2^d M(P)$ gives a stronger bound. This is also the case with cyclotomic polynomials ($M = 1$). Amoroso [5] and the author [21] slightly improved the constants in Mignotte's inequality. For instance, in the range $1 < M < d^\theta$ the inequality

$$H(P) \leq \|P\| \leq L(P) < \exp \left((1 + \theta) \sqrt{d/2} \log d \right)$$

is true for all d sufficiently large [21].

For the height $P(1)$, Bugeaud [14] obtained the bound

$$|P(1)| \leq M^4 \exp \left(2\sqrt{d \log d \log M} \right).$$

The proof is based on the analytical inequality

$$\left| \prod_{k=1}^{K-1} (z^k - 1)^{K-k} \right| \leq \max\{1, |z|\}^{K(K^2-1)/6} K^{K/2},$$

where z is a complex number and $K \geq 2$ is an integer. The above inequality follows from Hadamard's inequality, by representing the polynomial in the left hand side as

$$z^{-K(K-1)(K-2)/6} \text{Det}(x^{ij})_{0 \leq i, j \leq K-1}.$$

In some sense this proof can be compared with Zagier's method [32] used for a lower bound of $(M(\alpha)M(1-\alpha))^{1/d}$.

We remark that there is more information in the above inequality than just an upper bound for $|P(1)|$. Taking a product over all d conjugates of $\alpha \in \overline{\mathbb{Q}}^*$, $\alpha \neq 1$, with the leading coefficient a one easily gets the inequality

$$\sum_{k=1}^{K-1} \left(1 - \frac{k}{K}\right) \log |\mathcal{N}(\alpha^k - 1)| \leq \frac{K^2 - 1}{6} \log \left(\frac{M(\alpha)}{a}\right) + \frac{d}{2} \log K.$$

This “one line proof” sharpens the essentially Blanksby and Montgomery inequality [6] (see Silverman [29], [30], [31]) obtained using the Fourier averaging techniques. Silverman [30] speculates that such an inequality can be used to approach the Lehmer conjecture.

The author used some more subtle determinants [16], [17] which gave the inequality

$$|P(1)| < \exp \left(\left(\frac{\pi}{4} + \varepsilon \right) \sqrt{d \log d \log M} \right)$$

for any $\varepsilon > 0$ and for all d sufficiently large. These proved to be useful in estimating the number of cyclotomic factors of a polynomial [19]. The latter problem was first investigated by Schinzel [27]. See also the paper of Pinner and Vaaler [26].

Amoroso [1] showed that these bounds are not far from being sharp. He proved that for $\varepsilon > 0$ the inequality

$$|P(1)| > \exp \left((1 - \varepsilon) \sqrt{d \log M(P)} \right)$$

has infinitely many solutions in irreducible integer polynomials P such that $d^2/4 \leq M(P) \leq d^2/2$. By simplifying his example, we show that it also has infinitely many solutions for smaller $M(P)$, say $d < M(P) < (1 + \varepsilon)d$. It would be of interest to reduce the upper bound for $M(P)$ to, for instance, $M(P) \leq \sqrt{d}$, if possible.

For every integer $q \geq 2$, set

$$d = \sum_{p \leq q} (p - 1),$$

where the sum is taken over primes. Let $p(d)$ be a prime number greater than d , and let

$$\Phi_d(x) = x^{1+d-p(d)} \left(x^{p(d)-d} \prod_{p \leq q} (1 + x + x^2 + \dots + x^{p-1}) \right)'.$$

Clearly, $\Phi_d(x)$ is an integer polynomial of degree d .

THEOREM 2. Φ_d is an irreducible integer polynomial such that $M(\Phi_d) = p(d)$. Furthermore, for any $\varepsilon > 0$ and d sufficiently large we have

$$\Phi_d(1) > \exp\left((1 - \varepsilon)\sqrt{d \log d}\right).$$

Note that if $p(d) < \exp\left(\varepsilon\sqrt{d \log d}\right)$, then for any root α of Φ_d

$$\mathcal{N}(\alpha - 1) = \frac{\Phi_d(1)}{p(d)} > \exp\left((1 - 2\varepsilon)\sqrt{d \log d}\right),$$

because the coefficients of Φ_d are all nonnegative. Also, if say $p(d) \leq d^{1+\varepsilon}$, then

$$\mathcal{N}(\alpha - 1) > \exp\left((1 - 2\varepsilon)\sqrt{d \log M(\alpha)}\right).$$

PROOF. The polynomial

$$x^{p(d)-d} \prod_{p \leq q} (1 + x + \dots + x^{p-1})$$

has no multiple roots on the unit circle $|z| = 1$. Hence, the zeros of Φ_d are all strictly inside the unit circle, since the smallest closed convex set containing all the zeros of a polynomial also contains all the zeros of the derivative of the polynomial. Suppose that

$$\Phi_d(x) = p(d)x^d + \dots + p(d) - d$$

is reducible. From the fact that $p(d)$ is a prime number, we deduce that there is an irreducible integer polynomial Φ^* with the leading coefficient $p(d)$ so that $\Phi^* \mid \Phi_d$. The remaining zeros of the ratio Φ_d/Φ^* must contain a full set of conjugates of an algebraic integer. This is, however, impossible, because they are all strictly inside the unit circle $|z| < 1$, and $\Phi_d(0) \neq 0$. Consequently, $\deg \Phi_d = d$ and $M(\Phi_d) = p(d)$.

For the last inequality, note that if $\varepsilon > 0$ and if q is sufficiently large then

$$d = \sum_{p \leq q} (p - 1) < \left(1 + \frac{\varepsilon}{2}\right) \frac{q^2}{2 \log q}.$$

Hence $q > (1 - \varepsilon/2)\sqrt{d \log d}$. Since the coefficients of Φ_d are all nonnegative, it follows that

$$\Phi_d(1) > \prod_{p \leq q} p > \exp\left(\left(1 - \frac{\varepsilon}{2}\right)q\right) > \exp\left((1 - \varepsilon)\sqrt{d \log d}\right)$$

as required. ■

Our next statement shows that the multiplicity problem and the height problem are indeed related. Let

$$r(L, D) = \max r_1(f),$$

where the maximum is taken over all integer polynomials of degree D and of length at most L .

PROPOSITION 1. *For a given positive integers L, d , and an integer irreducible polynomial P of degree d , $d \geq D$, we have*

$$r(L, D) \log |P(1)| \leq D \log M(P) + d \log L.$$

PROOF. There is no loss of generality to assume that $r = r(L, D) > 0$. Hence, none of the roots of P is a root of f , where $r_1(f) = r \geq 1$. Let us take a product over all roots of the polynomial P , $\alpha_1, \alpha_2, \dots, \alpha_d$, and multiply by a^r , where a is its leading coefficient. Using the inequality $|f(z)| \leq \max\{1, |z|\}^D L$, we obtain

$$|P(1)|^r = a^r \prod_{j=1}^d |\alpha_j - 1|^r \leq a^r |f(\alpha_1) \dots f(\alpha_d)| \leq M(P)^D L^d,$$

whence the result. ■

The best known lower bound for $r(L, D)$ can be used to replace the constant $\pi/4$ by $1/\sqrt{2}$ in the above upper bound for $|P(1)|$ provided that $M(P)$ is not too small.

COROLLARY. *Let $0 < \nu < 2/3$. If P is an irreducible integer polynomial of degree d , where $d > d(\nu)$, such that $M(P) \geq d^{1/\nu}$, then*

$$|P(1)| \leq \exp \left(\frac{1+\nu}{\sqrt{2}} \sqrt{d \log d \log M(P)} \right).$$

PROOF. Let us fix a small positive constant ε , and let D be sufficiently large. By considering the polynomials with coefficients in \mathcal{A}_H , where $H = \lceil D^{1/\nu} / (D+1) \rceil$, and applying Theorem 3 in [2], we get

$$r(L, D) \geq r_1(\mathcal{A}_H, D) > (1 - \varepsilon) \sqrt{8 \left(\frac{1}{\nu} - 1 \right) D}.$$

Here $L = D^{1/\nu}$. Taking

$$d = \left\lceil \frac{\nu D \log M(P)}{\log D} \right\rceil + 1 \geq \left\lceil \frac{D \log d}{\log D} \right\rceil + 1 > \frac{D \log d}{\log D}$$

we see that $d > D$. On applying the inequality of Proposition 1, we get

$$\log |P(1)| < (1 + 2\varepsilon) \sqrt{\frac{\nu D}{2(1 - \nu)}} \log M(P).$$

Since $\nu D < d \log d / \log M(P)$ and $(1 - \nu)^{-1/2} < 1 + \nu$, the statement follows. \blacksquare

The example given in Theorem 2 via Proposition 1 implies the essentially Schur's bound for $r(L, D)$:

$$r(L, D) \leq (2 + \varepsilon) \sqrt{D \log L}.$$

A better example would imply stronger bounds.

Suppose $H = [D^\xi]$ with a fixed constant $\xi > 0$. From [2], [8] or [23] it follows that

$$\sqrt{D} \ll r_1(\mathcal{A}_H, D).$$

Let $\varphi(d)$ be any function such that $\varphi(d) \rightarrow \infty$ as $d \rightarrow \infty$. Suppose that there exist a positive constant γ and a sequence of irreducible integer polynomials P of degree d (where the sequence of the degrees is sufficiently dense) so that $M(P) \geq d^{\varphi(d)}$ and

$$|P(1)| \geq \exp \left(\gamma \sqrt{d \log d \log M(P)} \right).$$

Then, taking say $d = \lceil D \log M(P) / \xi \log D \rceil$ in Proposition 1, we would get that

$$r_1(\mathcal{A}_H, D) \ll \sqrt{D}.$$

4. Approximation problem. How well a fixed algebraic number β can be approximated by algebraic numbers α of small Mahler measure? This question was first investigated by Mignotte [22]. The worst rate of approximation seems to occur in the case when β is a root of unity (see also [11]). On replacing α by its power, or by $-\alpha$, if possible, we see that it suffices to investigate the case $\beta = -1$. The lower bounds for $|\alpha + 1|$ were obtained by Mignotte [22], Mignotte and Waldschmidt [25], Bugeaud, Mignotte and Normandin [15], and by the author [16]. For a fixed $\varepsilon > 0$ and d sufficiently large the inequality

$$|\alpha + 1| > \exp \left(- \left(\frac{\pi}{4} + \varepsilon \right) \sqrt{d \log d \log M(\alpha)} \right)$$

is true say in the range $1 < M(\alpha) < \exp(d^{1-\delta})$ (see [16]).

The existence of algebraic numbers close to 1 or -1 was first shown by Amoroso [3], [4]. The author [17] showed that there are infinitely many positive integers d such that for every such a d there is an algebraic number α of degree d , satisfying the inequalities $0 < \log M(\alpha) \ll \log d$, and

$$|\alpha + 1| < \exp\left(-2.8\sqrt{\frac{d \log M(\alpha)}{\log d}}\right).$$

PROPOSITION 2. For $\alpha \in \overline{\mathbb{Q}}^*$, $\alpha \neq 1$, of degree d , we have

$$|\alpha + 1| > \frac{3}{4dL(\alpha)}.$$

PROOF. The proposition clearly holds for $d = 1$. If $d \geq 2$, then $L(\alpha) \geq 3$. Hence, there is no loss of generality to assume that $|\alpha + 1| \leq 1/4d$. On replacing α by $1/\alpha$, if necessary, we may assume that $|\alpha| \geq 1$. Therefore

$$\begin{aligned} 1 \leq |P(-1)| &= |P(-1) - P(\alpha)| \leq |1 + \alpha| \max_{1 \leq |z| \leq |\alpha|} |P'(z)| \leq \\ &\leq |1 + \alpha| dL(P) \left(1 + \frac{1}{4d}\right)^{d-1} < \frac{4}{3} |1 + \alpha| dL(\alpha), \end{aligned}$$

and the statement follows. ■

The following bounds show the relation between the multiplicity and approximation problems.

THEOREM 3. Let H and D be two positive integers. For any root α , $\alpha \neq -1$, of a polynomial of degree D with coefficients in \mathcal{A} , where $\mathcal{A} \subset \mathcal{A}_H$, we have

$$|\alpha + 1| > \frac{1}{2H(D+1)} \left(\frac{r_D + 1}{eD}\right)^{r_D+1}.$$

Here $r_D = r_{-1}(\mathcal{A}, D)$.

Furthermore, let N and $s > 1$ be two positive even and odd integers, respectively, such that $(N+1)(2s+1) \leq D+1$, and let \mathcal{A} be a set of complex numbers. Then there is a polynomial of degree at most D with coefficients in \mathcal{A} and a root α , $\alpha \neq -1$, so that

$$|\alpha + 1| < s^{2-r_N},$$

where $r_N = r_{-1}(\mathcal{A}, N) \geq 1$.

PROOF. The first part is a simplified version of the method used in [11]. Let $f(x)$ be a polynomial of degree D with coefficients in \mathcal{A} so that $r = r_D = r_{-1}(\mathcal{A}, D)$. The modulus of the left hand side in the identity

$$\left| \frac{f(x)}{(x+1)^r} \right|_{x=1} = -(1+\alpha) \frac{1}{r!} \left(\frac{f(x)}{x-\alpha} \right)^{(r)} \Big|_{x=-1},$$

where α , $\alpha \neq -1$, is a root of f , is bounded below by 1. There is no loss of generality in assuming that $1 \leq |\alpha| \leq 1 + 1/2(D+1)$. In order to establish the lower bound note that

$$\begin{aligned} \left| \frac{1}{r!} \left(\frac{f(x) - f(\alpha)}{x - \alpha} \right)^{(r)} \right|_{x=-1} &\leq \max_{1 \leq |z| \leq |\alpha|} \left| \frac{f^{(r+1)}(z)}{(r+1)!} \right| \\ &\leq \frac{L(f)D^{r+1}|\alpha|^{D-1}}{(r+1)!} < 2H(D+1) \left(\frac{De}{r+1} \right)^{r+1}. \end{aligned}$$

For the second part we use the following statement which is true for any set of complex numbers \mathcal{A} .

OBSERVATION. *For a given positive integers N , s , and a set of complex numbers \mathcal{A} , let $f(x)$ be a polynomial of degree N with coefficients in \mathcal{A} . Then the coefficients of the polynomial*

$$f(x) + x^{N+1}(x^{Ns+s} + 1)(x^{s-1} + x^{s-2} + \dots + 1)f(x^s)$$

are all in \mathcal{A} .

Let now $f(x)$ be a polynomial of degree N with coefficients in \mathcal{A} so that $r_{-1}(f) = r_N = r_{-1}(\mathcal{A}, N) = r$. Set

$$\begin{aligned} g(x) &= \frac{f(x)}{(x+1)^r}, \\ G(x) &= g(x) + x^{N+1}(x^{Ns+s} + 1) \frac{x^s - 1}{x - 1} \left(\frac{x^s + 1}{x + 1} \right)^r g(x^s) \end{aligned}$$

with s so that $2Ns + 2s + N \leq D$.

Suppose first that $g'(-1)g(-1) \leq 0$. Since N is even and $s > 1$ is odd, it follows that

$$\begin{aligned} G(-1) &= g(-1) \neq 0, \\ G'(-1) &= g'(-1) + (-1)^{N+1}(Ns + s)s^r g(-1) \\ &= g'(-1) - (N+1)s^{r+1}g(-1). \end{aligned}$$

Thus, the polynomial G has a root α , $\alpha \neq -1$, such that

$$\left| \frac{G'(-1)}{G(-1)} \right| \leq \frac{\deg G}{|\alpha + 1|} = \frac{2Ns + 2s + N - r}{|\alpha + 1|} < \frac{3(N+1)s}{|\alpha + 1|}.$$

Here the expression in the left hand side is at least $(N+1)s^{r+1}$, so

$$|\alpha + 1| < \frac{3}{s^r} \leq \frac{1}{s^{r-1}}.$$

Also, α is a root of the polynomial $(x+1)^r G(x)$ of degree $2Ns+2s+N \leq D$ and with coefficients in \mathcal{A} (see Observation).

In the case $g'(-1)g(-1) > 0$ we take a slightly different polynomial:

$$G(x) = x^{2Ns+2s}g(x) + (x^{Ns+s} + 1)\frac{x^s - 1}{x - 1}\left(\frac{x^s + 1}{x + 1}\right)^r g(x^s).$$

This time

$$\begin{aligned} G(-1) &= g(-1), \\ G'(-1) &= -(2Ns + 2s)g(-1) + g'(-1) + (Ns + s)s^r g(-1) \\ &= g'(-1) + (N+1)s(s^r - 2)g(-1), \end{aligned}$$

and for a root α , $\alpha \neq -1$, of G we have

$$|\alpha + 1| < \frac{3(N+1)s}{(N+1)s(s^r - 2) + g'(-1)/g(-1)} < \frac{3}{s^r - 2} \leq \frac{3}{s^{r-1}} \leq \frac{1}{s^{r-2}}.$$

Again, α is a root of the polynomial

$$(x+1)^r G(x) = x^{2Ns+2s}f(x) + (x^{Ns+s} + 1)(x^{s-1} + \dots + 1)f(x^s)$$

of degree $2Ns+2s+N \leq D$ and with coefficients in \mathcal{A} . This completes the proof. \blacksquare

Let ε be a small positive constant. Taking $s = 3$ and applying Theorem 3 to the $\{-1, 0, 1\}$ polynomials, we obtain that

$$\exp\left(-\left(\frac{\log 3}{\sqrt{7}} + \varepsilon\right)r_D\right) < \min |\alpha + 1| < \exp\left(-\left(\frac{1}{2} - \varepsilon\right)r_D \log D\right),$$

the minimum being taken over all roots α , $\alpha \neq -1$, of the $\{-1, 0, 1\}$ polynomials of a sufficiently large degree D . Recall that here r_D is in the range $\sqrt{D/\log D} \ll r_D < \frac{21}{13}\sqrt{D}$.

Let N and s be the even and odd numbers, respectively, among $[\sqrt{D/2}] - 1$, $[\sqrt{D/2}] - 2$. Using the bounds

$$\frac{\log D}{\log 2} - 2 < r_D = r_{-1}(B, D) \leq \frac{\log D}{\log 2}$$

(see Theorem 1), from Theorem 3 we now deduce that

$$\exp\left(-(1+\varepsilon)\frac{(\log D)^2}{\log 2}\right) < \min |\alpha + 1| < \exp\left(-(1-\varepsilon)\frac{(\log D)^2}{4\log 2}\right).$$

Here the minimum is taken over all roots α , $\alpha \neq -1$, of the $\{0, 1\}$ polynomials of a sufficiently large degree D . Borwein and Pinner [11] obtained the constant $4 \log 3 = 4.394 \dots$ instead of $4 \log 2 = 2.772 \dots$.

Finally, taking N and s as above,

$$\sqrt{D/2} - 3 < N, s \leq \sqrt{D/2} - 1,$$

and using Boyd's [13] result (see also [11])

$$\log D \ll r_D = r_{-1}(\mathcal{A}^*, D) \ll \frac{(\log D)^2}{\log \log D},$$

we have

$$\exp\left(-(1 + \varepsilon)r_D \log D\right) < \min |\alpha + 1| < \exp\left(-\left(\frac{1}{8} - \varepsilon\right)r_D \log D\right).$$

Here the minimum is taken over all roots α , $\alpha \neq -1$, of the $\{-1, 1\}$ polynomials of a sufficiently large degree D .

This research was partially supported by a grant from Lithuanian Foundation of Studies and Science.

References

- [1] F. Amoroso, *Sur des polynômes de petites mesures de Mahler*, C. R. Acad. Sci. Paris **321** (1995), 11–14.
- [2] F. Amoroso, *Polynomials with prescribed vanishing at roots of unity*, Bolletino Un. Mat. Ital. B (7) **9** (1995), 1021–1042.
- [3] F. Amoroso, *Algebraic numbers close to 1 and variants of Mahler's measure*, J. Number Theory **60** (1996), 80–96.
- [4] F. Amoroso, *Algebraic numbers close to 1: results and methods*, in: Number Theory (V.K. Murthy, M. Waldschmidt eds.), Contemporary Mathematics **210**, 1997, 305–316.
- [5] F. Amoroso, *Upper bounds for the resultant and Diophantine applications*, in: Number Theory: Diophantine, Computational and Algebraic Aspects (K. Györy, A. Pethö, V.T. Sós eds.), Walter de Gruyter, Berlin, 1998, 23–36.
- [6] P.E. Blanksby and H.L. Montgomery, *Algebraic integers near the unit circle*, Acta Arithm. **18** (1971), 355–369.
- [7] A. Bloch and G. Pólya, *On the roots of certain algebraic equations*, Proc. London Math. Soc. **33** (1932), 102–114.
- [8] E. Bombieri and J.D. Vaaler, *On Siegel's lemma*, Invent. Math. **73** (1983), 11–32.
- [9] E. Bombieri and J.D. Vaaler, *Polynomials with low height and prescribed vanishing*, in: Analytic Number Theory and Diophantine Problems (A.C. Adolphson, J.B. Conrey, A. Ghosh and R.I. Yager

- eds.), Progress in Mathematics Series **70**, Birkhäuser, Basel, 1987, 53–73.
- [10] P. Borwein, T. Erdélyi and G. Kós, *Littlewood-type problems on $[0, 1]$* , Bull. London Math. Soc. **79** (1999), 22–46.
 - [11] P. Borwein and C. Pinner, *Polynomials with $\{0, +1, -1\}$ coefficients and a root close to a given point*, Canad. J. Math. **49** (1997), 887–915.
 - [12] P. Borwein and M.G. Mossinghoff, *Polynomials with height 1 and prescribed vanishing at 1*, Experiment. Math. (to appear).
 - [13] D.W. Boyd, *On a problem of Byrnes concerning polynomials with restricted coefficients*, Math. Comp. **66** (1997), 1697–1703.
 - [14] Y. Bugeaud, *Algebraic numbers close to 1 in non-archimedean metrics*, The Ramanujan J. **2** (1998), 449–457.
 - [15] Y. Bugeaud, M. Mignotte and F. Normandin, *Nombres algébriques de petite mesure et formes linéaires en un logarithme*, C. R. Acad. Sci. Paris **321** (1995), 517–522.
 - [16] A. Dubickas, *On algebraic numbers of small measure*, Liet. Matem. Rink. **35** (1995), 421–431.
 - [17] A. Dubickas, *On algebraic numbers close to 1*, Bull. Australian Math. Soc. **58** (1998), 423–434.
 - [18] A. Dubickas, *Polynomials with high multiplicity at unity and Tarry’s problem*, Math. Notes **65** (1999), 684–688.
 - [19] A. Dubickas, *On a polynomial with large number of irreducible factors*, in: Number Theory in Progress, vol. **1**: Diophantine Problems and Polynomials (K. Györy, H. Iwaniec, J. Urbanowicz eds.), Walter de Gruyter, Berlin, 1999, 103–110.
 - [20] A. Dubickas, *On the order of vanishing at 1 of a polynomial*, Liet. Matem. Rink. **39** (1999), 461–468.
 - [21] A. Dubickas, *On certain geometric mean of the values of a polynomial*, (to appear).
 - [22] M. Mignotte, *Approximation des nombres algébriques par des nombres algébriques de grand degré*, Ann. Fac. Sci. Toulouse **1** (1979), 165–170.
 - [23] M. Mignotte, *Estimations élémentaires effectives sur les nombres algébriques*, Journées de Théorie Analytique et Élémentaire des Nombres, Caen (1980), 24–34.
 - [24] M. Mignotte, *An inequality about irreducible factors of integer polynomials*, J. Number Theory **30** (1988), 156–166.
 - [25] M. Mignotte and M. Waldschmidt, *On algebraic numbers of small height: linear forms in one logarithm*, J. Number Theory **47** (1994), 43–62.
 - [26] C. Pinner and J.D. Vaaler, *The number of irreducible factors of a polynomial. I*, Trans. Amer. Math. Soc. **339** (1993), 809–834.

- [27] A. Schinzel, *On the number of irreducible factors of a polynomial*, Colloq. Math. Soc. János Bolyai **13** (1976), 305–314.
- [28] I. Schur, *Untersuchungen über algebraische Gleichungen. I, Bemerkungen zu einem Satz von E. Schmidt*, Sitzungsber. Preuss. Acad. Wiss., Phys.-Math. Kl. **7/10** (1933), 403–428.
- [29] J.H. Silverman, *Lehmer’s conjecture and primes of small norm*, Brown University, preprint (1994).
- [30] J.H. Silverman, *Exceptional units and numbers of small Mahler measure*, Experiment. Math. **4** (1995), 69–83.
- [31] J.H. Silverman, *Small Salem numbers, exceptional units, and Lehmer’s conjecture*, Rocky Mountain J. Math. **26** (1996), 1099–1114.
- [32] D. Zagier, *Algebraic numbers close to both 0 and 1*, Math. Comp. **61** (1993), 485–491.

Present address:

Fakultät für Mathematik
 Universität Bielefeld
 Postfach 100 131
 D-33501 Bielefeld
 Germany

Permanent address:

Department of Mathematics and Informatics
 Vilnius University
 Naugarduko 24
 2600 Vilnius, Lithuania
 E-mail: arturas.dubickas@maf.vu.lt