

INVARIANTS, TORSION INDICES AND ORIENTED COHOMOLOGY OF COMPLETE FLAGS

B. CALMÈS, V. PETROV, AND K. ZAINOULLINE

CONTENTS

1. Introduction	1
Part 1. Invariants, torsion indices and formal group laws.	4
2. Formal group rings	4
3. Differential operators Δ_α and C_α	7
4. Endomorphisms of a formal group ring	10
5. Torsion indices and augmented operators	12
6. Invariants and the characteristic map	15
7. The product structure of the cohomology	17
 Part 2. Bott-Samelson resolutions for oriented theories.	 19
8. Algebraic cobordism and oriented cohomology theories	19
9. A sequence of split \mathbb{P}^1 -bundles	21
10. An elementary step of the Bott-Samelson resolution	23
11. Bott-Samelson varieties	25
12. Bott-Samelson resolutions	26
 Part 3. Algebraic and geometric comparison. Applications.	 27
13. Comparison results	27
14. Formulas for push-forwards	30
15. Algorithm for multiplying in $h^*(G/B)$	32
16. Landweber-Novikov operations	32
17. Examples of computations	33
References	35

1. INTRODUCTION

Let H be an algebraic cohomology theory endowed with Chern classes c_i such that for any two line bundles \mathcal{L}_1 and \mathcal{L}_2 over a variety X we have

$$(1) \quad c_1(\mathcal{L}_1 \otimes \mathcal{L}_2) = c_1(\mathcal{L}_1) + c_2(\mathcal{L}_2).$$

The basic example of such a theory is the Chow group CH of algebraic cycles modulo rational equivalence.

Let G be a split semi-simple linear algebraic group over a field k and let T be a split maximal torus inside G contained in a Borel subgroup B . Consider the variety G/B of Borel subgroups of G with respect to T . In two classical papers [7] and [8]

The first author was supported by EPSRC grant EP/E01786X/1. The second author is supported by RFBR 08-01-00756, 09-01-00878, 09-01-90304, 09-01-91333. The third author is supported by NSERC Discovery Grant.

Demazure studied the cohomology ring $H(G/B; \mathbb{Z})$ and provided an algorithm to compute $H(G/B; \mathbb{Z})$ in terms of generators and relations.

The main object of his consideration was the so called characteristic map

$$(2) \quad \mathbf{c}: S^*(M) \rightarrow H(G/B; \mathbb{Z}),$$

where $S^*(M)$ is the symmetric algebra of the character group M of T . In [7] Demazure interpreted this map from the point of view of invariant theory of the Weyl group W of G by identifying its kernel with the ideal generated by non-constant invariants $S^*(M)^W$. The cohomology ring $H(G/B; \mathbb{Z})$ was then replaced by a certain algebra constructed in terms of operators and defined in purely combinatorial terms.

In the present paper, we generalize most of the results of [7] to the case of an arbitrary oriented cohomology theory \mathfrak{h} , i.e. when (1) is replaced by

$$c_1(\mathcal{L}_1 \otimes \mathcal{L}_2) = F(c_1(\mathcal{L}_1), c_1(\mathcal{L}_2))$$

where F is the formal group law associated to \mathfrak{h} . Such theories were extensively studied by Levine-Morel [13], Panin-Smirnov [16], Merkurjev [14] and others. Apart from the Chow ring, other examples include algebraic K -theory, étale cohomology $H_{\text{ét}}^*(-, \mu_m)$, $(m, \text{char}(k)) = 1$, Morava K -theories, connective K -theory, elliptic cohomology and the universal such theory: algebraic cobordism Ω .

To generalize the characteristic map (2), we first introduce a substitute for the symmetric algebra $S^*(M) = \text{CH}^*(BT)$. This new combinatorial object, which we call a *formal group ring*, is denoted by $R[[M]]_F$, where $R = \mathfrak{h}(pt)$ is the coefficient ring, and can be viewed as a substitute of the cohomology ring of the classifying space $\mathfrak{h}(BT)$ of T . As in [7], we introduce a subalgebra $\mathcal{D}(M)_F$ of the R -linear endomorphisms of $R[[M]]_F$ generated by specific differential operators and by taking its R -dual we obtain $\mathcal{H}(M)_F$, a combinatorial substitute for the cohomology ring $\mathfrak{h}(G/B; \mathbb{Z})$. The characteristic map (2) then turns into the map

$$\mathbf{c}: R[[M]]_F \rightarrow \mathcal{H}(M)_F.$$

The Weyl group W acts naturally on $R[[M]]_F$ and the main result of our paper (theorems 6.4 and 6.9) says that:

Theorem. *If the torsion index of G is invertible in R and R has no 2-torsion, then the characteristic map is surjective and its kernel is generated by W -invariant elements in the augmentation ideal.*

Demazure's methods to prove this theorem in the special case of the additive formal group law do not work in general, for the following reason: the main objects used in his proofs are operators Δ_w for every $w \in W$. They are defined first for simple reflections, and afterwards for any w by decomposing it into simple reflections and composing the corresponding operators. It is then proved that the resulting composition is independent of the decomposition of w . For more general formal group laws, similar operators can still be defined for simple reflections (see Definition 3.5), but independence of the decomposition does not hold, as it was observed in [5]. Geometrically, it can be translated into the fact that the cobordism class of a desingularized Schubert variety depends on the desingularization, and not only on the Schubert variety itself (see Lemma 13.9). We overcome this problem by working with suitable filtrations such that the associated graded structures are covered by the additive case of Demazure (see in particular Proposition 4.4). We therefore encourage the reader unfamiliar with [7] to start by having a quick look at it before reading our sections 3 to 7.

As an immediate application of the developed techniques, we provide an efficient algorithm for computing the cohomology ring $\mathcal{H}(M)_F = \mathfrak{h}(G/B; \mathbb{Z})$. To do this

we generalize the Bott-Samelson approach introduced in [1] and [8]. For oriented topological theories, some algorithms were considered by Bressler-Evens in [4, 5] and for algebraic theories in characteristic 0 by Hornbostel-Kiritchenko in [12]. See remark 15.2 for a comparison.

Note that the theorem also provides another approach to computing the cohomology ring $\mathfrak{h}(G/B; \mathbb{Z})$ by looking at the subring of invariants $R[[M]]_F^W$. Observe that in the classical case when $\mathfrak{h} = \text{CH}$ (or K_0) and G is simply-connected it is known that $R[[M]]_F^W$ is a power series ring in basic polynomial invariants (resp. fundamental representations). In general, the structure of $R[[M]]_F^W$ remains unknown.

Finally, for the reader primarily interested in topology, let us mention that while all our proofs are algebraic and written in the language of algebraic geometry, the results apply as they are to topological cobordism or other complex oriented theories. Indeed, there is a canonical ring morphism

$$\Omega^*(G/B) \rightarrow \text{MU}^*(G/B(\mathbb{C}))$$

(see [13, Ex. 1.2.10]) that is an isomorphism because both are free modules over the Lazard ring with bases corresponding to each other (given by desingularized Schubert cells).

The paper consists of three parts.

In the first part, we generalize the results of [7] by introducing and studying the generalized characteristic map $\mathfrak{c}: R[[M]]_F \rightarrow \mathcal{H}(M)_F$. In section 2, we introduce the formal group ring $R[[M]]_F$ and prove its main properties. In section 3, we define the main operators Δ and C on $R[[M]]_F$. In sections 4 and 5 we study the subalgebra $\mathcal{D}(M)_F$ of operators generated by Δ (resp. C) and multiplications. In section 6 we define $\mathcal{H}(M)_F$ and prove the main theorem. In section 7 we introduce a product on $\mathcal{H}(M)_F$ compatible with the characteristic map \mathfrak{c} .

In the second part, we generalize some of the results of [8] to arbitrary oriented cohomology theories. In section 8, we discuss properties of oriented theories. In sections 9 to 12, we carry out the Bott-Samelson desingularization approach.

In the last part, we apply the results of the first and the second parts to obtain information about the ring structure in $\mathfrak{h}(G/B; \mathbb{Z})$. In section 13, we prove that our algebraic replacement $\mathcal{H}(M)_F$ is isomorphic (as a ring) to the oriented cohomology $\mathfrak{h}^*(G/B; \mathbb{Z})$ and that the characteristic maps $\mathfrak{c}_{G/B}$ and \mathfrak{c} correspond to each other via this isomorphism. In section 14, we give an algebraic description of the push-forward to the point and we prove various formulas. In sections 15 and 16, we explain an algorithm to compute the ring structure of $\mathfrak{h}^*(G/B; \mathbb{Z})$ and the Landweber-Novikov operations on algebraic cobordism. Finally, in section 17, we give multiplication tables for $\Omega^*(G/B; \mathbb{Z})$ for groups G of rank 2.

Notation. Let k denote a base field of arbitrary characteristic. A variety over k means a reduced irreducible scheme of finite type over k . By X and Y we always mean smooth varieties over k . The base point $\text{Spec } k$ is denoted by pt .

A ring always means a commutative ring with a unit and R always denotes a ring. A ring R' is called an R -algebra if it comes equipped with an injective ring homomorphism $R \hookrightarrow R'$. The letter M always denotes an abelian group.

All formal group laws are assumed to be one-dimensional and commutative. Let F denote a formal group law and let \mathbb{L} denote the Lazard ring, i.e. the coefficient ring of the universal formal group law U .

Acknowledgments. Our sincere gratitude goes to Michel Demazure for his answers to our questions concerning the paper [7]. We would like to thank the authors of the Macaulay 2 software [10], and, in particular Dan Grayson for his quick and accurate responses on the Macaulay 2 mailing list. We would also like to

thank Vladimir Chernousov, Fabien Morel and Burt Totaro for their encouraging attention to our work.

Part 1. Invariants, torsion indices and formal group laws.

2. FORMAL GROUP RINGS

Let R be a ring, let M be an abelian group and let F be a formal group law over R . In the present section we introduce and study the *formal group ring* $R[[M]]_F$. For this, we use several auxiliary facts concerning topological rings and their completions which can be found in [3, III, §2]. The main result here is the decomposition theorem 2.10. At the end we provide some examples of computations of $R[[M]]_F$.

2.1. Definition. Let R be a ring and let S be a set. Let $R[x_S] := R[x_s, s \in S]$ denote the polynomial ring over R with variables indexed by S . Let $\epsilon : R[x_S] \rightarrow R$ be the augmentation morphism which maps any x_s to 0. Consider the $\ker(\epsilon)$ -adic topology on $R[x_S]$ given by ideals $\ker(\epsilon)^i$, $i \geq 0$, which form a fundamental system of open neighborhoods of 0. Note that a polynomial is in $\ker(\epsilon)^i$ if and only if its valuation is at least i , hence, we have $\bigcap_i \ker(\epsilon)^i = \{0\}$ and the $\ker(\epsilon)$ -adic topology is Hausdorff.

We define $R[[x_S]]$ to be the $\ker(\epsilon)$ -adic completion of the polynomial ring $R[x_S]$.

2.2. Remark. By definition, an element in $R[[x_S]]$ can be written uniquely as a formal sum

$$\sum_{s_1, \dots, s_m \in S} a_{s_1, \dots, s_m} \cdot x_{s_1} \dots x_{s_m}, \quad a_{s_1, \dots, s_m} \in R,$$

where for any positive m there is only a finite number of non-zero coefficients a_{s_1, \dots, s_m} . In particular, when S is a finite set of order n , the ring $R[[x_S]]$ is the usual ring of power series in n variables.

2.3. Let F be a formal group law over a ring R (see [13, p.4]). Given an integer $m \geq 0$ we use the notation

$$x +_F y = F(x, y), \quad m \cdot_F x = \underbrace{x +_F \dots +_F x}_{m \text{ times}} \quad \text{and} \quad (-m) \cdot_F x = -_F(m \cdot_F x)$$

By associativity of F , for any $m_1, m_2 \in \mathbb{Z}$ we have

$$(m_1 + m_2) \cdot_F x = (m_1 \cdot_F x) +_F (m_2 \cdot_F x).$$

Now, here comes the definition playing a central role in the sequel.

2.4. Definition. Let M be an abelian group and let F be a formal group law over a ring R . Consider the ring $R[[x_M]]$ and let \mathcal{J}_F be the closure of the ideal generated by the elements x_0 and $x_{\lambda+\mu} - (x_\lambda +_F x_\mu)$ for all $\lambda, \mu \in M$. We define the *formal group ring* $R[[M]]_F$ to be the quotient

$$R[[M]]_F = R[[x_M]] / \mathcal{J}_F.$$

The class of x_λ in $R[[M]]_F$ will be denoted by the same letter.

2.5. By definition, $R[[x_M]]$ is a complete Hausdorff ring with respect to the topology induced by the kernel of the augmentation $R[[x_M]] \rightarrow R$. Since \mathcal{J}_F is clearly contained in this kernel, the augmentation map $R[[x_M]] \rightarrow R$ factors through the quotient $R[[M]]_F$. Therefore, $R[[M]]_F$ is a *complete Hausdorff* ring with respect to the \mathcal{I}_F -adic topology, where \mathcal{I}_F denotes the kernel of the augmentation $R[[M]]_F \rightarrow R$.

Let $f: R \rightarrow R'$ be a morphism of rings respecting the formal group laws, i.e. sending every coefficient of F to the corresponding coefficient of F' . Then, for every abelian group M , f induces a ring homomorphism $f_\star: R[[M]]_F \rightarrow R'[[M]]_{F'}$ sending

$x_\lambda \in R[[M]]_F$ to $x_\lambda \in R'[[M]]_{F'}$ for every $\lambda \in M$. This morphism sends \mathcal{I}_F to $\mathcal{I}_{F'}$ and, hence, is continuous.

Let now $f: M \rightarrow M'$ be a morphism of abelian groups. It induces a continuous ring homomorphism $\hat{f}: R[[M]]_F \rightarrow R[[M']]_F$ sending x_λ to $x_{f(\lambda)}$. Moreover, if f is surjective, then so is \hat{f} .

Finally, let $f: F \rightarrow F'$ be a morphism of formal group laws over R , i.e. a formal power series $f \in R[[x]]$ such that $f(x +_F y) = f(x) +_{F'} f(y)$. Then, f induces a continuous ring homomorphism $f^*: R[[M]]_{F'} \rightarrow R[[M]]_F$ sending x_λ to $f(x_\lambda)$.

We have therefore proved:

2.6. Lemma. *Via the above constructions $(-)_*$, $(\hat{-})$ and $(-)^*$, the assignment taking (R, M, F) to the topological ring $R[[M]]_F$ is covariant with respect to ring morphisms $R \rightarrow R'$ and morphisms of abelian groups $M \rightarrow M'$, and is contravariant with respect to morphisms of formal group laws $F \rightarrow F'$.*

2.7. Let \mathbb{L} be the Lazard ring of coefficients of the universal formal group law U . For any formal group law F over R , there is a unique ring homomorphism $\vartheta_F: \mathbb{L} \rightarrow R$ sending the universal formal group law U on \mathbb{L} to F . By universality, the morphism

$$R \otimes_{\mathbb{L}} \mathbb{L}[[M]]_U \rightarrow R[[M]]_F, \quad r \otimes z \mapsto r \cdot (\vartheta_F)_*(z)$$

is an isomorphism.

2.8. Since $R[[M]]_F$ is an R -algebra, we may consider the formal group law F as an element of $R[[M]]_F[[x, y]]$. Let $a, b \in \mathcal{I}_F$. The specialization at $x = a$ and $y = b$ defines a pairing

$$\boxplus: \mathcal{I}_F \times \mathcal{I}_F \rightarrow \mathcal{I}_F.$$

Similarly, we define $\boxminus: \mathcal{I}_F \rightarrow \mathcal{I}_F$ using the inverse of F and $\boxdot: \mathbb{Z} \times \mathcal{I}_F \rightarrow \mathcal{I}_F$ by applying \boxplus or \boxminus iteratively. From the associativity, commutativity, inverse properties of F and the continuity of the quotient map $R[[x_M]] \rightarrow R[[M]]_F$ it follows that $x_{\lambda+\mu} = x_\lambda \boxplus x_\mu$ and

$$x_{-\lambda} = x_{-\lambda} \boxplus (x_\lambda \boxminus x_\lambda) = (x_{-\lambda} \boxplus x_\lambda) \boxminus x_\lambda = 0 \boxminus (x_\lambda) = \boxminus(x_\lambda).$$

2.9. A formal group law F on R induces via a ring homomorphism $R \rightarrow R'$ a formal group law on R' . In particular, if M and N are abelian groups, then using $R \rightarrow R[[M]]_F = R'$, we can define $R[[M]]_F[[N]]_F$, which is naturally an $R[[M]]_F$ -algebra. By functoriality, $R[[M \oplus N]]_F$ is also an $R[[M]]_F$ -algebra. We define two morphisms of $R[[M]]_F$ -algebras

$$\phi: R[[M \oplus N]]_F \rightarrow R[[M]]_F[[N]]_F \quad \text{and} \quad \psi: R[[M]]_F[[N]]_F \rightarrow R[[M \oplus N]]_F$$

as follows:

The map $R[[x_{M \oplus N}]] \rightarrow R[[x_M]][[y_N]]$ sending $x_{(\lambda, \gamma)}$ to $x_\lambda +_F y_\gamma$ extends to a continuous map $f: R[[x_{M \oplus N}]] \rightarrow R[[x_M]][[y_N]]$. Consider the composition

$$R[[x_{M \oplus N}]] \xrightarrow{f} R[[x_M]][[y_N]] \xrightarrow{g} R[[M]]_F[[y_N]] \xrightarrow{h} R[[M]]_F[[N]]_F$$

where g is induced by $R[[x_M]] \rightarrow R[[M]]_F$ and h is the quotient map. We have

$$\begin{aligned} & h \circ g \circ f(x_{(\lambda, \gamma)} +_F x_{(\mu, \delta)}) \stackrel{(1)}{=} h \circ g((x_\lambda +_F y_\gamma) +_F (x_\mu +_F y_\delta)) = \\ & \stackrel{(2)}{=} h \circ g((x_\lambda +_F x_\mu) +_F (y_\gamma +_F y_\delta)) \stackrel{(3)}{=} h((x_\lambda \boxplus x_\mu) +_F (y_\gamma +_F y_\delta)) = \\ & \stackrel{(4)}{=} (x_\lambda \boxplus x_\mu) +_F (y_\gamma \boxplus y_\delta) \stackrel{(5)}{=} x_{\lambda+\mu} +_F y_{\gamma+\delta} \stackrel{(6)}{=} h(x_{\lambda+\mu} +_F y_{\gamma+\delta}) = \\ & \stackrel{(7)}{=} h \circ g(x_{\lambda+\mu} +_F y_{\gamma+\delta}) \stackrel{(8)}{=} h \circ g \circ f(x_{(\lambda+\mu, \gamma+\delta)}), \end{aligned}$$

where (1) and (8) hold by definition and continuity of f , (2) by associativity and commutativity of F , (3) and (7) by definition of g , (4) and (6) by definition of h and (5) by definition of \boxplus . It shows that $h \circ g \circ f$ factors through $R[[M \oplus N]]_F$ as

$$h \circ g \circ f: R[[x_{M \oplus N}]] \rightarrow R[[M \oplus N]]_F \xrightarrow{\phi} R[[M]]_F[[N]]_F,$$

where ϕ is a morphism of $R[[M]]_F$ -algebras.

In the opposite direction, we proceed as follows: We extend the functorial morphism $R[[M]]_F \rightarrow R[[M \oplus N]]_F$ to a morphism $R[[M]]_F[[y_N]] \rightarrow R[[M \oplus N]]_F$ by sending y_γ to $x_{(0, \gamma)}$. It induces a morphism $R[[M]]_F[[y_N]] \rightarrow R[[M \oplus N]]_F$ on completions, which factors through a continuous morphism $\psi: R[[M]]_F[[N]]_F \rightarrow R[[M \oplus N]]_F$.

2.10. Theorem. *The morphisms of $R[[M]]_F$ -algebras ϕ and ψ defined above are inverses to each other. In other words, we have an isomorphism*

$$R[[M \oplus N]]_F \simeq R[[M]]_F[[N]]_F.$$

Proof. We have

$$\psi \circ \phi(x_{(\lambda, \gamma)}) = \psi(x_\lambda +_F y_\gamma) = \psi(x_\lambda) +_F \psi(y_\gamma) = x_{(\lambda, 0)} +_F x_{(0, \gamma)} = x_{(\lambda, \gamma)}$$

where the second equality holds by continuity of ψ . The other composition can be checked on y_γ , since we are dealing with morphisms of $R[[M]]_F$ -algebras, and we have $\phi \circ \psi(y_\gamma) = \phi(x_{(0, \gamma)}) = x_0 +_F y_\gamma = 0 +_F y_\gamma = y_\gamma$. \square

2.11. Lemma. *Let $M = \mathbb{Z}$. Then sending x_m to $m \cdot_F x$ defines a ring isomorphism*

$$R[[\mathbb{Z}]]_F \simeq R[[x]].$$

In particular, if M is a free abelian group of rank one, then $R[[M]]_F$ is isomorphic to $R[[x]]$.

Proof. The morphism $\phi: R[[x_{\mathbb{Z}}]] \rightarrow R[[x]]$ sending x_n to $n \cdot_F x$ extends to a morphism $\hat{\phi}: R[[x_{\mathbb{Z}}]] \rightarrow R[[x]]$. By continuity, $\hat{\phi}$ satisfies $\hat{\phi}(x_m +_F x_n) = \hat{\phi}(x_m) +_F \hat{\phi}(x_n)$ as well as $\hat{\phi}(-_F(x_m)) = -_F \hat{\phi}(x_m)$. Thus $\hat{\phi}(x_{m+n}) = \hat{\phi}(x_m +_F x_n)$ and $\hat{\phi}$ factors through a $\bar{\phi}: R[[\mathbb{Z}]]_F \rightarrow R[[x]]$. A map $\hat{\psi}$ in the opposite direction is defined similarly using completeness, by sending x to x_1 . By continuity, checking that $\bar{\phi} \circ \hat{\psi} = \text{id}$ and $\hat{\psi} \circ \bar{\phi} = \text{id}$ can be done on generators. Namely, $\bar{\phi} \circ \hat{\psi}(x) = \bar{\phi}(x_1) = x$ by definition, and $\hat{\psi} \circ \bar{\phi}(x_n) = \hat{\psi}(n \cdot_F x) = n \cdot_F \hat{\psi}(x) = n \cdot_F x_1 = x_n$. \square

2.12. Corollary. *Let $\phi: M \rightarrow \mathbb{Z}^{\oplus n}$ be an isomorphism. Then ϕ induces an isomorphism*

$$R[[M]]_F \simeq R[[x_1, \dots, x_n]].$$

Proof. It follows from Lemma 2.11 and Theorem 2.10 by induction on n . \square

2.13. Remark. Note that the right hand side is independent of F , although the isomorphism depends on F . Also note that if R is an integral domain, so is $R[[\mathbb{Z}^{\oplus n}]]_F$.

2.14. Example. Similarly, one can prove that $R[[\mathbb{Z}/n]]_F \simeq R[[x]]/(n \cdot_F x)$ by sending x_m to $m \cdot_F x$. Observe that $n \cdot_F x = nx + x^2q$. In particular, if n is invertible in R , then $(n \cdot_F x) = (x)$ and $R[[\mathbb{Z}/n]]_F \simeq R$.

Let us now examine what happens at a finite level in $R[[M]]_F$.

2.15. Let $R[M]_F$ denote the subring of $R[[M]]_F$ defined as the image of the subring $R[x_M]$ by means of the composition $R[x_M] \rightarrow R[[x_M]] \rightarrow R[[M]]_F$. Then the ring $R[[M]]_F$ is the completion of $R[M]_F$ at the ideal $\mathcal{I}_F \cap R[M]_F$. By the functoriality of $R[[M]]_F$ (see 2.6) the assignment

$$(R, M, F) \mapsto R[M]_F$$

is a *covariant* functor with respect to morphism of rings $R \rightarrow R'$ and morphisms of abelian groups $M \rightarrow M'$. Moreover, if $M \rightarrow M'$ is surjective, then so is $R[M]_F \rightarrow R[M']_F$.

2.16. Example. If $M = \mathbb{Z}$, the isomorphism of Lemma 2.11 maps $R[\mathbb{Z}]_F$ to polynomials in $n \cdot_F x$. In particular, unless F and the formal inverse have a finite number of nonzero coefficients only, $R[\mathbb{Z}]_F$ does *not* map to $R[x]$.

2.17. Remark. The morphism ϕ from 2.9 doesn't necessarily send $R[M \oplus N]_F$ to $R[M]_F[N]_F$. The morphism ψ from 2.9 sends $R[M]_F[N]_F$ into $R[M \oplus N]_F$ but is not necessarily surjective.

2.18. Example (cf. 8.3). The additive formal group law over R is given by $F(x, y) = x + y$. In this case we have ring isomorphisms

$$R[[M]]_F \simeq \prod_{i=0}^{\infty} S_R^i(M) \quad \text{and} \quad R[M]_F \simeq \bigoplus_{i=0}^{\infty} S_R^i(M)$$

where $S_R^i(M)$ is the i -th symmetric power of M over R and the isomorphisms are induced by sending x_λ to $\lambda \in S_R^1(M)$.

2.19. Example (cf. 8.4). The multiplicative periodic formal group law over R is given by $F(x, y) = x + y - \beta xy$, where β is an invertible element in R . Consider the group ring

$$R[M] := \left\{ \sum_j r_j e^{\lambda_j} \mid r_j \in R, \lambda_j \in M \right\}.$$

Let $\text{tr} : R[M] \rightarrow R$ be the trace map, i.e. a R -linear map sending any e^λ to 1. Let $R[M]^\wedge$ be the completion of $R[M]$ at $\ker(\text{tr})$. Then we have ring isomorphisms

$$R[[M]]_F \simeq R[M]^\wedge \quad \text{and} \quad R[M]_F \simeq R[M]$$

induced by $x_\lambda \mapsto \beta^{-1}(1 - e^\lambda)$ and $e^\lambda \mapsto (1 - \beta x_\lambda) = (1 - \beta x_{-\lambda})^{-1}$.

2.20. Example (cf. 8.5). The multiplicative non-periodic formal group law over R is given by $F(x, y) = x + y - vxy$, where v is not invertible in R . Specializing $R[[M]]_F$ at $v = 0$ or $v = \beta$ where β is invertible in R , we obtain the formal group rings of the previous examples.

2.21. Remark. Let \mathfrak{h} be an oriented cohomology theory as defined in [13, Def. 1.1.2] and let F be the associated formal group law (see Section 8). Let M be the group of characters of a split torus T over k . Then the formal group ring $R[[M]]_F$ can be viewed as an algebraic substitute of the completed equivariant cohomology ring $\mathfrak{h}_T(\text{pt})^\wedge$ or the cohomology ring of the classifying space $\mathfrak{h}(BT)$. Its finite counterpart $R[M]_F$ could then play the role of the equivariant cohomology ring $\mathfrak{h}_T(\text{pt})$ itself.

3. DIFFERENTIAL OPERATORS Δ_α AND C_α

In the present section we introduce two linear operators on $R[[M]]_F$. The first operator Δ_α^F is a generalized version of the operator Δ_α from [7, §3 and §4]. Indeed, one recovers the results of loc. cit. when $R = \mathbb{Z}$ and F is the additive formal group law from Example 2.18. A version of the second operator C_α^F was already used in [5, §5] for topological complex cobordism.

3.1. Consider a reduced root system as in [7, §1], i.e. a free \mathbb{Z} -module M of finite rank, a finite subset of M whose elements are called *roots* and a map associating

a coroot $\alpha^\vee \in M^\vee$ to a root α , satisfying certain axioms.¹ The reflection map $\lambda \mapsto \lambda - \alpha^\vee(\lambda)\alpha$ is denoted by s_α .

3.2. The Weyl group W associated to a reduced root system is the subgroup of linear automorphisms of M generated by reflections s_α . It acts linearly on M and thus by R -algebra automorphisms on $R[[M]]_F$ using the functoriality in M of $R[[M]]_F$. We have the following obvious equalities for any $u \in R[[M]]_F$:

$$s_\alpha(u) = s_{-\alpha}(u) \quad \text{and} \quad s_\alpha^2(u) = u.$$

3.3. **Lemma.** *For any $n \in \mathbb{Z}$, the element $x - x +_F(n \cdot_F y)$ is uniquely divisible by y in $R[[x, y]]$.*

Proof. Since y is a regular element in $R[[x, y]]$, we just need to prove divisibility. Note that for any power series $g(x, y)$, the series $g(x, 0) - g(x, y)$ is divisible by y . Apply it to $g(x, y) = x +_F(n \cdot_F y)$. \square

3.4. **Corollary.** *Assume that R is integral. For any $u \in R[[M]]_F$, the element $u - s_\alpha(u)$ is uniquely divisible by x_α .*

Proof. First note that $R[[M]]_F$ is an integral domain by 2.13, so we just need to prove divisibility. Since

$$s_\alpha(x_\lambda) = x_{\lambda - \alpha^\vee(\lambda)\alpha} = x_\lambda \boxplus (-\alpha^\vee(\lambda) \boxtimes x_\alpha)$$

it holds by the previous lemma when $u = x_\lambda$. Then, by the formula

$$(3) \quad uv - s_\alpha(uv) = (u - s_\alpha(u))v + u(v - s_\alpha(v)) - (u - s_\alpha(u))(v - s_\alpha(v))$$

the result holds by induction on the degree of monomials for any element in $R[[M]]_F$. Finally, it holds by density on the whole $R[[M]]_F$. \square

For a given root α we define a linear operator Δ_α^F on $R[[M]]_F$ as follows:

3.5. **Definition.** First, using Corollary 3.4 we define a linear operator Δ_α^U on $\mathbb{L}[[M]]_U$, where U is the universal formal group law over the Lazard ring \mathbb{L} , as

$$\Delta_\alpha^U(u) = \frac{u - s_\alpha(u)}{x_\alpha}, \quad \text{where } u \in \mathbb{L}[[M]]_U.$$

Finally, identifying $R[[M]]_F$ with $R \otimes_{\mathbb{L}} \mathbb{L}[[M]]_U$ via the isomorphism from 2.7 we define the desired operator Δ_α^F on $R[[M]]_F$ as $\text{id}_R \otimes \Delta_\alpha^U$. We will simply write $\Delta_\alpha(u)$ when the formal group law F over R is understood.

3.6. **Remark.** Observe that if R is integral, then the operator Δ^F can be defined directly using the same formula as for Δ^U . If R is torsion, then it is not the case. Indeed, take $R = \mathbb{Z}/2$, $M = \mathbb{Z}$, F to be an additive law and $\alpha = 2$ to be a root. Then $x_\alpha = x_{1+1} = x_1 + x_1 = 2x_1 = 0$ in $R[[M]]_F$.

3.7. **Remark.** We could instead have defined Δ_α as $\Delta_\alpha(u) = \frac{s_\alpha(u) - u}{x_{-\alpha}}$. This exchanges Δ_α with $-\Delta_{-\alpha}$ so it is easy to switch from one convention to the other. Both conventions give the same classical operator when the formal group law is additive.

3.8. **Proposition.** *The following formulas hold for any $u, v \in R[[M]]_F$, $\lambda \in M$ and $w \in W$ (Compare to [7, §3]²).*

- (1) $\Delta_\alpha(1) = 0, \quad \Delta_\alpha(u)x_\alpha = u - s_\alpha(u),$
- (2) $\Delta_\alpha^2(u)x_\alpha = \Delta_\alpha(u) + \Delta_{-\alpha}(u), \quad \Delta_\alpha(u)x_\alpha = \Delta_{-\alpha}(u)x_{-\alpha},$
- (3) $s_\alpha\Delta_\alpha(u) = -\Delta_{-\alpha}(u), \quad \Delta_\alpha s_\alpha(u) = -\Delta_\alpha(u),$

¹This is often called a root datum instead of a root system in the literature, but we follow [7].

²There is a sign mistake in Equation (3) of [7, §3]. One should read $\Delta_\alpha = -\Delta_{-\alpha}$

- (4) $\Delta_\alpha(uv) = \Delta_\alpha(u)v + u\Delta_\alpha(v) - \Delta_\alpha(u)\Delta_\alpha(v)x_\alpha = \Delta_\alpha(u)v + s_\alpha(u)\Delta_\alpha(v)$,
(5) $w\Delta_\alpha w^{-1}(u) = \Delta_{w(\alpha)}(u)$.

Proof. All formulas can be proved in $\mathbb{L}\llbracket M \rrbracket_U$ (which is an integral domain) and then specialized to any other $R\llbracket M \rrbracket_F$. Formula (4) follows from equation (3) above. Formula (5) follows from the fact that $\alpha^\vee(w^{-1}(\lambda)) = w(\alpha)^\vee(\lambda)$. All other formulas follow by definition. \square

From now on, we fix a basis of simple roots $(\alpha_1, \dots, \alpha_n)$, with associated simple reflections s_1, \dots, s_n . This defines a length function on W . Let I denote a sequence (i_1, \dots, i_l) of l integers in $[1, n]$ and let $w(I) = s_{i_1} \cdots s_{i_l}$ be the corresponding product of simple reflections. The decomposition I is *reduced* if w has length l . We define the linear operator Δ_I as the composition

$$\Delta_I = \Delta_{\alpha_{i_1}} \circ \cdots \circ \Delta_{\alpha_{i_l}}.$$

The operators Δ_I have the following important property:

3.9. Theorem. *Let F be a formal group law of the form $F(x, y) = x + y - v \cdot xy$ for some $v \in R$. Let I and I' be two reduced decompositions of w in simple reflections. Then $\Delta_I = \Delta_{I'}$.*

Proof. See [8, Theorem 2 p. 86]. The proof assumes $v = 1$ but it works for any other value. \square

When the formal group law is of the above type, the previous theorem justifies the notation Δ_w instead of Δ_I , but in general, we have to keep the dependence on the decomposition.

3.10. Remark. Theorem 3.9 was proved in [4] (see loc. cit. Thm.3.7) for topological oriented theories. Moreover, the result of [4] says that the formal group law has to be of the above form for such an independence of the decomposition to hold.

For a given root α we define another linear operator C_α^F on $R\llbracket M \rrbracket_F$ as follows:

3.11. Definition. Let $g(x, y)$ be the power series defined by

$$x +_F y = x + y - xy \cdot g(x, y)$$

and let e_α be the element $g(x_\alpha, x_{-\alpha})$ in $R\llbracket M \rrbracket_F$. We set

$$C_\alpha^F(u) = ue_\alpha - \Delta_\alpha^F(u), \quad \text{where } u \in R\llbracket M \rrbracket_F.$$

We will simply write $C_\alpha(u)$ when the formal group law F over R is understood. For a sequence of integers I in $[1, n]$, we define the operator C_I in the same way as Δ_I .

3.12. Proposition. *The following formulas hold for any $u, v \in R\llbracket M \rrbracket_F$, $\lambda \in M$ and $w \in W$.*

- (1) $C_\alpha(1) = e_\alpha, \quad C_\alpha(x_{-\alpha}) = 2,$
- (2) $C_\alpha(u)x_\alpha x_{-\alpha} = ux_\alpha + s_\alpha(u)x_{-\alpha}, \quad C_\alpha(ux_{-\alpha}) = u + s_\alpha(u),$
- (3) $C_\alpha s_\alpha(u) = C_{-\alpha}(u), \quad s_\alpha C_\alpha(u) = C_\alpha(u),$
- (4) $C_\alpha(uv) = C_\alpha(u)v + s_\alpha(u)C_\alpha(v) - s_\alpha(u)ve_\alpha = C_\alpha(u)v - s_\alpha(u)\Delta_\alpha(v),$
- (5) $wC_\alpha w^{-1}(u) = C_{w(\alpha)}(u),$
- (6) $C_\alpha \Delta_\alpha = \Delta_\alpha C_\alpha = \Delta_\alpha C_{-\alpha} = 0.$

Proof. All these formulas can easily be derived from the ones of Proposition 3.8 using the definition of C_α and the fact that $e_\alpha x_\alpha x_{-\alpha} = x_\alpha + x_{-\alpha}$. \square

3.13. Proposition. *Let W_α be the subgroup of order 2 generated by s_α in W . Let $R[[M]]_F^{W_\alpha}$ denote the subring of fixed elements of $R[[M]]_F$ under the action of W .*

If the ring R has no 2-torsion, then the operators Δ_α and C_α are $R[[M]]_F^{W_\alpha}$ -linear. In particular, they are $R[[M]]_F^W$ -linear.

Proof. For the operator Δ_α , by formula (4) of 3.8 it suffices to show that $\Delta_\alpha(u) = 0$ for $u \in R[[M]]_F^{W_\alpha}$. According to formula (3) of 3.8, it is equivalent to $\Delta_\alpha(u) = -\Delta_\alpha(u)$ which holds if $R[[M]]_F$ has no 2-torsion.

The same facts for C_α then follow from its definition in terms of Δ_α . \square

3.14. Lemma. *Assume R has no 2-torsion. Then for any root α , the element x_α is regular in $R[[M]]_F$ and the set of elements u such that $\Delta_\alpha(u) = 0$ is $R[[M]]_F^{W_\alpha}$.*

Proof. Let $\omega_1, \dots, \omega_n$ be the fundamental weights corresponding to a choice of simple roots $\alpha = \alpha_1, \dots, \alpha_n$ of the root system. Possibly extending M , we can assume that the ω_i are a basis of M . By the isomorphism $R[[M]]_F \simeq R[[x_{\omega_1}, \dots, x_{\omega_n}]]$ of Corollary 2.12, x_α becomes $\sum_i \langle \alpha, \alpha_i \rangle x_{\omega_i} + u$, $u \in \mathcal{I}^2$. One easily checks that this element is regular if at least one of the $\langle \alpha, \alpha_i \rangle$ is regular in R . In particular, this is the case if $\langle \alpha_1, \alpha_1 \rangle = 2$ is regular in R . By equation (1) of Proposition 3.8, the last part of the claim is then clear. \square

3.15. Proposition. *Let $f : R \rightarrow R'$ be a ring morphism sending a formal group law F over R to a formal group law F' over R' . Then, the operators Δ and C satisfy*

$$f_* \Delta_\alpha^F = \Delta_\alpha^{F'} f_* \quad \text{and} \quad f_* C_\alpha^F = C_\alpha^{F'} f_*$$

where $f_* : R[[M]]_F \rightarrow R'[[M]]_{F'}$ is the morphism of Lemma 2.6.

Proof. This is clear by construction. \square

4. ENDOMORPHISMS OF A FORMAL GROUP RING

In the present section we introduce and study the subalgebra $\mathcal{D}(M)_F$ of R -linear endomorphisms of a formal group ring $R[[M]]_F$ generated by the Δ operators of the previous section. The main result (Theorem 4.11) says that operators $\Delta_{I_w}^F$, where w runs through all elements of the Weyl group and I_w is any chosen reduced decomposition of w , form a basis of $\mathcal{D}(M)_F$ as an $R[[M]]_F$ -module.

4.1. Let $R[[M]]_F$ be a formal group ring corresponding to a formal group law F over a ring R and an abelian group M (see Def. 2.4). Let \mathcal{I}_F be the kernel of the augmentation map $R[[M]]_F \rightarrow R$ as in 2.5. By convention, we set $\mathcal{I}_F^i = R[[M]]_F$ for any $i \leq 0$. We define the associated graded ring

$$\mathcal{G}r_R^*(M, F) = \bigoplus_{i=0}^{\infty} \mathcal{I}_F^i / \mathcal{I}_F^{i+1}.$$

4.2. Lemma. *The morphism of graded R -algebras*

$$\phi : S_R^*(M) \rightarrow \mathcal{G}r_R^*(M, F)$$

defined by sending λ to x_λ is an isomorphism.

Proof. The map is well-defined since $x_{\lambda+\mu} - (x_\lambda + x_\mu)$ is in \mathcal{I}_F^2 , and it is obviously surjective. Let us define a map in the other direction. Recall that by definition $R[[M]]_F = R[[x_M]] / \mathcal{J}_F$. Let $\tilde{\mathcal{I}}$ denote the kernel of the augmentation map $R[[x_M]] \rightarrow R$ (see 2.5). Then we have

$$\mathcal{I}_F = \tilde{\mathcal{I}} / \mathcal{J}_F \quad \text{and} \quad \mathcal{I}_F^i / \mathcal{I}_F^{i+1} = \tilde{\mathcal{I}}^i / (\mathcal{J}_F \cap \tilde{\mathcal{I}}^i + \tilde{\mathcal{I}}^{i+1}).$$

Since $\tilde{\mathcal{I}}^i / \tilde{\mathcal{I}}^{i+1}$ is given by monomials of degree i , we may define a map of R -modules

$$\psi_i : \tilde{\mathcal{I}}^i / \tilde{\mathcal{I}}^{i+1} \rightarrow S_R^i(M)$$

by sending a monomial of degree i in some x_λ -s to the symmetric product of the λ -s involved. This map passes to the quotient since $(\mathcal{I}_F \cap \tilde{\mathcal{I}}^i + \tilde{\mathcal{I}}^{i+1})/\tilde{\mathcal{I}}^{i+1}$ is generated as an R -module by elements of the form $\rho \cdot (x_\lambda + x_\mu - x_{\lambda+\mu})$ where ρ is a monomial of degree $i-1$. The sum $\oplus_i \psi_i$ passes through the quotient as well and, hence, gives the desired inverse map. \square

4.3. Consider now a reduced root system M as in Section 3 and let $R[[M]]_F$ be the associated formal group ring. Consider the operators Δ_α^F introduced in Def. 3.5. By their very definition, the operators Δ_α^F send \mathcal{I}_F^i to \mathcal{I}_F^{i-1} . Hence, they induce R -linear operators of degree -1 on the graded ring $\mathcal{G}r_R^*(M, F)$, denoted by $\mathcal{G}r\Delta_\alpha^F$. Similarly, we define graded versions of operators C_α^F (see Def. 3.11) on $\mathcal{G}r_R^*(M, F)$, denoted by $\mathcal{G}rC_\alpha^F$.

4.4. **Proposition.** *The isomorphism ϕ of Lemma 4.2 exchanges the operator $\mathcal{G}r\Delta_\alpha^F$ (resp. $\mathcal{G}rC_\alpha^F$) on $\mathcal{G}r_R^*(M, F)$ with $\Delta_\alpha^{F_a}$ (resp. $-\Delta_\alpha^{F_a}$) on the symmetric algebra $S_R^*(M)$, where F_a denotes the additive formal group law as in Example 2.18 and the $\Delta_\alpha^{F_a} = -C_\alpha^{F_a}$ are the classical operators of [7].*

Proof. Induction on the degree using (4) of Prop. 3.8. \square

4.5. **Definition.** Let $\mathcal{D}(M)_F$ be the subalgebra of R -linear endomorphisms of $R[[M]]_F$ generated by the Δ_α^F for all roots α and by multiplications by elements of $R[[M]]_F$. Note that by formula (1) of Proposition 3.8, $\mathcal{D}(M)_F$ contains s_α and it contains C_α^F by its definition. Let $\mathcal{D}(M)_F^{(i)}$ be the sub $R[[M]]_F$ -module of $\mathcal{D}(M)_F$ generated by the $u\Delta_{\alpha_1}^F \cdots \Delta_{\alpha_n}^F$, where $u \in \mathcal{I}_F^m$ and $m - n \geq i$. This defines a filtration on $\mathcal{D}(M)_F$

$$\mathcal{D}(M)_F \supseteq \cdots \supseteq \mathcal{D}(M)_F^{(i)} \supseteq \mathcal{D}(M)_F^{(i+1)} \supseteq \cdots \supseteq 0$$

with the property that $\mathcal{D}(M)_F = \cup_i \mathcal{D}(M)_F^{(i)}$. We define the associated graded $R[[M]]_F$ -module

$$\mathcal{G}r^*\mathcal{D}(M)_F = \bigoplus_{i=-\infty}^{\infty} \mathcal{D}(M)_F^{(i)} / \mathcal{D}(M)_F^{(i+1)}.$$

4.6. **Proposition.** *The filtration on $\mathcal{D}(M)_F$ above has the following properties:*

(1) *For any root α , we have*

$$\Delta_\alpha^F \mathcal{D}(M)_F^{(i)} \subseteq \mathcal{D}(M)_F^{(i-1)}, \quad \mathcal{D}(M)_F^{(i)} \Delta_\alpha^F \subseteq \mathcal{D}(M)_F^{(i-1)}$$

and similarly for C_α^F

(2) *For any integer j , we have $\mathcal{I}_F^j \mathcal{D}(M)_F^{(i)} \subseteq \mathcal{D}(M)_F^{(i+j)}$*

(3) *For any operator $D \in \mathcal{D}(M)_F^{(i)}$, we have $D(\mathcal{I}_F^j) \subseteq \mathcal{I}_F^{i+j}$*

(4) $\cap_i \mathcal{D}(M)_F^{(i)} = \{0\}$

Proof. The claim (1) follows from equation (4) in Prop. 3.8 (resp. equation (4) in Prop. 3.12). The claim (2) is obvious. The claim (3) follows from the fact that $\Delta_\alpha^F(\mathcal{I}_F^i) \subseteq \mathcal{I}_F^{i-1}$. To prove (4) observe that any D in $\cap_i \mathcal{D}(M)_F^{(i)}$ sends $R[[M]]_F$ to $\cap_i \mathcal{I}_F^i = 0$ by (3) so is the zero operator. \square

4.7. By claim (3) of Prop. 4.6 the graded module $\mathcal{G}r^*\mathcal{D}(M)_F$ acts by graded endomorphisms on the graded ring $\mathcal{G}r_R^*(M, F)$, which is isomorphic to $S_R^*(M)$ by Lemma 4.2. Let F_a denote the additive formal group law. Then the associated subalgebra $\mathcal{D} = \mathcal{D}(M)_{F_a}$ coincides with one considered in [7, §3]. By Prop. 4.4, the class of Δ_α^F (resp. C_α^F) in $\mathcal{G}r^{(-1)}\mathcal{D}(M)_F$ acts by $\Delta_\alpha^{F_a}$ (resp. $-\Delta_\alpha^{F_a}$) on $S_R^*(M)$ and the class of the multiplication by x_α in $\mathcal{G}r^{(1)}\mathcal{D}(M)_F$ acts by the multiplication by α on $S_R^*(M)$.

4.8. Proposition. *The graded module $\mathcal{G}r^*\mathcal{D}(M)_F$ over the graded ring $\mathcal{G}r^*_R(M, F) \simeq S^*_R(M)$ is isomorphic to \mathcal{D} .*

Proof. It is a sub-module of the R -linear endomorphisms of $\mathcal{G}r^*R[[M]]_F$ generated by the same elements as \mathcal{D} . \square

4.9. Let A be a ring and let \mathcal{I} be an ideal of A , such that A is complete and Hausdorff for the \mathcal{I} -adic topology. Let M be an A -module together with a \mathcal{I} -filtration $(M^{(i)})_{i \in \mathbb{Z}}$, i.e. $\mathcal{I}M^{(i)} \subseteq M^{(i+1)}$ for any i . Consider the associated graded ring $\mathcal{G}r^*A$ and the associated graded $\mathcal{G}r^*A$ -module $\mathcal{G}r^*M$.

The filtration is called *exhaustive* (see [2, III, §2, 1]) if any $\lambda \in M$ belongs to some $M^{(i)}$, i.e. $M = \cup_i M^{(i)}$. For an exhaustive filtration let $\nu(\lambda)$ denote the largest integer such that $\lambda \in M^{\nu(\lambda)}$ and let $\bar{\lambda}$ denote the class of λ in $M^{(\nu(\lambda))}/M^{(\nu(\lambda)+1)}$ if $\nu(\lambda)$ exists and 0 if $\nu(\lambda)$ does not exist. Note that $\bar{\lambda}$ is nonzero if and only if λ is not in $\cap_i M^{(i)}$.

4.10. Lemma. *Let M be an A -module with an exhaustive \mathcal{I} -filtration. Let $\lambda_1, \dots, \lambda_n$ be elements of M . Then*

- (1) *Assume A is complete for the \mathcal{I} -adic topology and M is Hausdorff. If $\bar{\lambda}_1, \dots, \bar{\lambda}_n$ generate $\mathcal{G}r^*M$ as a $\mathcal{G}r^*A$ -module, then $\lambda_1, \dots, \lambda_n$ generate M as an A -module.*
- (2) *Assume A is Hausdorff. If $\bar{\lambda}_1, \dots, \bar{\lambda}_n$ are independent elements in the $\mathcal{G}r^*A$ -module $\mathcal{G}r^*M$, then $\lambda_1, \dots, \lambda_n$ are independent in the A -module M .*
- (3) *Assume A is Hausdorff complete and M is Hausdorff. If $\mathcal{G}r^*M$ is a finitely generated free $\mathcal{G}r^*A$ -module, then M is a finitely generated free A -module.*

Proof. Let $A[i]$ be A itself considered as an A -module with the shifted filtration $\mathcal{I}^{j+\nu(\lambda_i)}$ (by convention, $\mathcal{I}^j = A$ for $j \leq 0$). The claims (1) and (2) then follow from [3, III, §2, 8, Cor. 1 and 2] applied to $X = \bigoplus_i A[i]$, $Y = M$ and $\bigoplus_i A[i] \rightarrow M$ the map sending $(a_i)_i$ to $\sum_i a_i \lambda_i$. The last claim (3) is an immediate consequence of (1) and (2). \square

We now come to the main result of this section:

4.11. Theorem. *Let $\mathcal{D}(M)_F$ be the algebra of operators defined above. For each element $w \in W$, choose a reduced decomposition I_w in simple reflections. Then the operators $\Delta_{I_w}^F$ (resp. $C_{I_w}^F$) form a basis of $\mathcal{D}(M)_F$ as an $R[[M]]_F$ -module.*

Proof. When the formal group law is additive, this is proved in [7, §4, Cor. 1] for the $\Delta_w^{F_a}$ (and $C_w^{F_a} = (-1)^{l(w)} \Delta_w^{F_a}$). The filtered module $\mathcal{D}(M)_F$ is Hausdorff by point (4) in Proposition 4.6, and $R[[M]]_F$ is complete. We can therefore apply point (3) of Lemma 4.10 to deduce the general case from the corresponding fact on the associated graded objects, which is the additive case by Proposition 4.8. \square

5. TORSION INDICES AND AUGMENTED OPERATORS

In the present section we study the augmentation $\epsilon\mathcal{D}(M)_F$ of the algebra of operators $\mathcal{D}(M)_F$. The main result is Proposition 5.4 which is what becomes of Theorem 4.11 when the augmentation ϵ is applied. Similarly, we then examine the filtration on $\epsilon\mathcal{D}(M)_F$ obtained by applying ϵ to the one on $\mathcal{D}(M)_F$ introduced in 4.5.

5.1. Consider the *torsion index* t of the root system as defined in [7, §5] and its prime divisors, called *torsion primes*. It has been computed for simply connected root systems of all types (see [7, §7, Prop. 8] and [18, 19]). The results of these computations are summarized in the following table:

Type	A_l	$B_l, l \geq 3$	C_l	$D_l, l \geq 4$	G_2	F_4	E_6	E_7	E_8
Torsion primes	\emptyset	2	\emptyset	2	2	2, 3	2, 3	2, 3	2, 3, 5
Torsion index	1	$2^{e(l)}$	1	$2^{e(l-1)}$	2	$2 \cdot 3$	$2 \cdot 3$	$2^2 \cdot 3$	$2^6 \cdot 3^2 \cdot 5$

The exponent $e(l)$ is equal to $l - \lfloor \log_2 \binom{l+1}{2} \rfloor$ except for certain values of l equal or slightly larger than a power of 2 which are explicitly given in [19, Thm. 0.1]. Since $\mathfrak{t}(\mathcal{R}_1 \times \mathcal{R}_2) = \mathfrak{t}(\mathcal{R}_1) \cdot \mathfrak{t}(\mathcal{R}_2)$, we may compute torsion indices of semi-simple root systems by reducing to irreducible ones.

5.2. Let M be the character group of a root system of G and let F be a formal group law over a ring R . Let $\epsilon: R[[M]]_F \rightarrow R$ be the augmentation map with the kernel \mathcal{I}_F . Let N be the length of the longest element w_0 of the Weyl group W of G . Let F_a be an additive formal group law.

By definition of the torsion index (see [7, §5]) there exists an element a of the symmetric algebra $S_R^N(M)$ such that

$$\Delta_{w_0}^{F_a}(a) = \mathfrak{t}.$$

Let $\phi: S_R^*(M) \xrightarrow{\cong} \mathcal{G}r_R^*(M, F)$ be an isomorphism from Lemma 4.2. Let $u_0 \in \mathcal{I}_F^N$ be an element such that $\bar{u}_0 = \phi(a)$. Therefore, for any reduced decomposition I_0 of w_0 we have $\epsilon \Delta_{I_0}^F(u_0) = \mathfrak{t}$ or, in other words, by Prop. 4.4

$$\Delta_{I_0}^F(u_0) = \mathfrak{t} + \mathcal{I}_F \quad \text{and} \quad C_{I_0}^F(u_0) = (-1)^N \mathfrak{t} + \mathcal{I}_F \quad \text{in } R[[M]]_F.$$

5.3. **Lemma.** *Let I be a sequence of simple reflections and let $l(I)$ be its length. Then*

- (1) for any $i \in \mathbb{Z}$, $\Delta_I^F(\mathcal{I}_F^i) \subseteq \mathcal{I}_F^{i-l(I)}$ and $C_I^F(\mathcal{I}_F^i) \subseteq \mathcal{I}_F^{i-l(I)}$;
- (2) if $l(I) \leq i$ and I is not reduced, then

$$\Delta_I^F(\mathcal{I}_F^i) \subseteq \mathcal{I}_F^{i-l(I)+1} \quad \text{and} \quad C_I^F(\mathcal{I}_F^i) \subseteq \mathcal{I}_F^{i-l(I)+1};$$

- (3) let $u_0 \in \mathcal{I}_F^N$ be the element chosen above. If $l(I) \leq N$ then

$$\epsilon \Delta_I^F(u_0) = (-1)^N \epsilon C_I^F(u_0) = \begin{cases} \mathfrak{t} & \text{if } I \text{ is reduced and } l(I) = N \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Claim (1) follows from Prop. 4.6, (1). To check (2), it therefore suffices to check that $\mathcal{G}r \Delta_I^F$ is zero. It follows from the corresponding fact in the additive case [7, §4, Prop. 3, (a)] after applying Prop. 4.4. The ‘‘otherwise’’ case of (3) follows from (2). The case where I is reduced and has length N follows from the definition of u_0 . \square

5.4. **Proposition.** *Consider the R -module $\epsilon \mathcal{D}(M)_F$ of all R -linear forms ϵD , where $D \in \mathcal{D}(M)_F$. Assume the torsion index \mathfrak{t} is regular in R . Then for any choice of a collection of reduced decomposition I_w for each $w \in W$, the $(\epsilon \Delta_{I_w})_{w \in W}$ (resp. the $(\epsilon C_{I_w})_{w \in W}$) form an R -basis of $\epsilon \mathcal{D}(M)_F$.*

Proof. By Theorem 4.11, the $\epsilon \Delta_{I_w}$ generate $\epsilon \mathcal{D}(M)_F$. We prove that the $\epsilon \Delta_{I_w}$ such that $l(I_w) \leq i$ are independent by induction on i . It is obviously true when $i < 0$. Let $\sum_{l(w) \leq i} r_w \epsilon \Delta_{I_w} = 0$. By induction, it suffices to show that the coefficients r_w with $l(w) = i$ are zero. For any v and w of length i , the concatenation $I_w + I_{v^{-1}w_0}$ is a reduced decomposition of w_0 if and only if $v = w$. Thus, by Lemma 5.3.(1) and (3) we have $\sum_{l(w) \leq i} r_w \epsilon \Delta_{I_w} (\Delta_{I_{v^{-1}w_0}}(u_0)) = r_v \mathfrak{t}$ which implies that $r_v = 0$ by regularity of \mathfrak{t} . The same proof works for the C_{I_w} . \square

5.5. **Lemma.** *Assume \mathfrak{t} is regular in R .*

- (1) For any reduced decomposition I_w of w and any element $u \in \mathcal{I}_F^i \setminus \mathcal{I}_F^{i+1}$, the element $u \Delta_{I_w}^F$ is in $\mathcal{D}(M)_F^{(i-l(w))} \setminus \mathcal{D}(M)_F^{(i-l(w)+1)}$.

(2) The group $\mathcal{D}(M)_F^{(i)}$ is the set of $D \in \mathcal{D}(M)_F$ such that $D(\mathcal{I}_F^j) \subseteq \mathcal{I}_F^{i+j}$.

Proof. It suffices to prove (1) after inverting the torsion index. By Lemma 5.3.(1), the operator $u\Delta_{I_w}^F$ is in $\mathcal{D}(M)_F^{(i-l(w))}$. Consider the element $y = \Delta_{I_{w^{-1}w_0}}^F(u_0)$. It is in $\mathcal{I}_F^{l(w)}$. By Lemma 5.3.(3) the element $s = \Delta_{I_w}^F(y)$ is invertible. Hence, $u\Delta_{I_w}^F(y) = us$ cannot be in \mathcal{I}_F^{i+1} , and therefore $u\Delta_{I_w}^F$ is not in $\mathcal{D}(M)_F^{(i-l(w)+1)}$.

To prove (2), first note that Prop. 4.6.(3) immediately gives one inclusion. Suppose D is such that $D(\mathcal{I}_F^j) \subseteq \mathcal{I}_F^{i+j}$. Using Theorem 4.11 we may write it as

$$D = \sum_w u_w \Delta_{I_w}^F.$$

Recall that $\nu(u_w)$ denotes the largest integer such that $u_w \in \mathcal{I}_F^\nu$ (see 4.9). Consider the set S of $w \in W$ such that $\nu(u_w) - l(w)$ is minimal and take an element $w_1 \in S$ such that $l(w_1)$ is maximal. To show that $D \in \mathcal{D}(M)_F^{(i)}$, it suffices to show that $\nu(u_{w_1}) - l(w_1) \geq i$ by minimality of elements of S .

Let $y_1 = \Delta_{I_{w_1^{-1}w_0}}^F(u_0)$ be the corresponding element in $\mathcal{I}^{l(w_1)}$. We have

$$D(y_1) = u_{w_1} \Delta_{I_{w_1}}^F(y_1) + \sum_{w \in S \setminus \{w_1\}} u_w \Delta_{I_w}(y_1) + \sum_{w \notin S} u_w \Delta_{I_w}(y_1).$$

By Lemma 5.3.(1) and (2), two sums on the right are in $\mathcal{I}_F^{\nu(u_{w_1})+1}$. As in the proof of (1), $u_{w_1} \Delta_{I_{w_1}}^F(y_1)$ is not in $\mathcal{I}_F^{\nu(u_{w_1})+1}$ so $D(y_1)$ is in $\mathcal{I}_F^{\nu(u_{w_1})} \setminus \mathcal{I}_F^{\nu(u_{w_1})+1}$. Since $D(y_1)$ is also in $\mathcal{I}_F^{l(w_1)+i}$, we obtain that $\nu(u_{w_1}) - l(w_1) \geq i$. \square

We now consider the filtration on $\epsilon\mathcal{D}(M)_F$ that is the image by ϵ of the filtration on $\mathcal{D}(M)_F$ defined in 4.5, i.e.

$$\epsilon\mathcal{D}(M)_F \supseteq \dots \supseteq \epsilon\mathcal{D}(M)_F^{(i)} \supseteq \epsilon\mathcal{D}(M)_F^{(i+1)} \supseteq \dots \supseteq 0.$$

5.6. Proposition. *Assume \mathfrak{t} is regular in R . This filtration satisfies the following.*

- (1) For any $i \in \mathbb{Z}$ and $\epsilon D \in \epsilon\mathcal{D}(M)_F^{(i)}$, we have $\epsilon D(\mathcal{I}_F^{-i+1}) = 0$.
- (2) Let E be a subset of W of elements of length l , and let the I_w be reduced decompositions of each of the $w \in E$. Then any nonzero R -linear combination

$$f = \sum_{w \in S} r_w \cdot \epsilon \Delta_{I_w}^F \quad (\text{resp. } \epsilon C_{I_w}^F)$$

is in $\epsilon\mathcal{D}(M)_F^{-l} \setminus \epsilon\mathcal{D}(M)_F^{-l+1}$. In particular, for any w

$$\epsilon \Delta_{I_w}^F \in \epsilon\mathcal{D}(M)_F^{-l(w)} \setminus \epsilon\mathcal{D}(M)_F^{-l(w)+1} \quad (\text{resp. with } C_{I_w}^F).$$

- (3) For any choice of reduced decompositions I_w for every element $w \in W$, the R -module $\epsilon\mathcal{D}(M)_F^{(-i)}$ is a free R -module with basis the $\epsilon\Delta_{I_w}$ (resp. ϵC_{I_w}) with $l(w) \leq i$.
- (4) For any $i \leq -N$, we have $\epsilon\mathcal{D}(M)_F^{(i)} = \epsilon\mathcal{D}(M)_F$ and for any $i > 0$, we have $\epsilon\mathcal{D}(M)_F^{(i)} = 0$, i.e. the filtration is of the form

$$\epsilon\mathcal{D}(M)_F = \epsilon\mathcal{D}(M)_F^{(-N)} \supseteq \epsilon\mathcal{D}(M)_F^{(-N+1)} \supseteq \dots \supseteq \epsilon\mathcal{D}(M)_F^{(0)} \supseteq 0.$$

Proof. Claim (1) follows from Lemma 5.5.(2). To prove (2), we first apply f to $\Delta_{I_{w^{-1}w_0}}^F(u_0)$. By Lemma 5.3.(3), we obtain tr_w , which has to be zero by (1) for the element f to be in \mathcal{I}_F^{l+1} . Therefore, each r_w is zero. The last two claims follow. \square

6. INVARIANTS AND THE CHARACTERISTIC MAP

We now come to the definition of an algebraic replacement $\mathcal{H}(M)_F$ for the oriented cohomology $\mathfrak{h}^*(G/B)$ when G is a split semi-simple simply connected algebraic group corresponding to the root system and B is a Borel of G . The identification of $\mathcal{H}(M)_F$ with $\mathfrak{h}^*(G/B)$ is the subject of section 13.

6.1. Definition. Let $\mathcal{H}(M)_F$ be the R -dual of $\epsilon\mathcal{D}(M)_F$ and let

$$\mathfrak{c}^F : R[[M]]_F \rightarrow \mathcal{H}(M)_F$$

denote the natural map obtained by duality, i.e. sending u to the evaluation at u . When the formal group law is clear from the context, we write \mathfrak{c} for \mathfrak{c}^F .

Again, when $F = F_a$ is additive, this $\mathcal{H}(M)_F$ corresponds to the one defined by Demazure in [7, §3].

6.2. Remark. Observe that when R has no 2-torsion, since the operators Δ_α^F are $R[[M]]_F$ -linear by Proposition 3.13, the characteristic map satisfies $\mathfrak{c}(fu) = \epsilon(f)\mathfrak{c}(u)$ when f is in $R[[M]]_F^W$.

6.3. Theorem. Let $(I_w)_{w \in W}$ be a choice of reduced decompositions. When the torsion index \mathfrak{t} is regular in R , there is a unique R -basis $z_{I_w}^\Delta$ of $\mathcal{H}(M)_F$ such that the characteristic map is given by

$$\mathfrak{c}(u) = \sum_w \epsilon \Delta_{I_w}^F(u) z_{I_w}^\Delta.$$

Similarly, there is a unique R -basis $z_{I_w}^C$ of $\mathcal{H}(M)_F$ such that the characteristic map is given by

$$\mathfrak{c}(u) = \sum_w \epsilon C_{I_w}^F(u) z_{I_w}^C.$$

Proof. We take the bases that are dual to the ones of Proposition 5.4. \square

6.4. Theorem. Assume that the torsion index \mathfrak{t} is invertible in R . Then for any choice of a collection of reduced decompositions I_w for every $w \in W$, the $\mathfrak{c}(\Delta_{I_w}(u_0))$ (resp. the $\mathfrak{c}(C_{I_w}(u_0))$) form an R -basis of the image of \mathfrak{c} . In particular, the characteristic map \mathfrak{c} is surjective.

Proof. We have $\mathfrak{c}(u_0) = \mathfrak{t} z_{I_0}^\Delta$ and

$$\mathfrak{c}(\Delta_{I_{w^{-1}w_0}}(u_0)) = \mathfrak{t} z_w^\Delta + \sum_{l(v) > l(w)} \epsilon \Delta_{I_v} \Delta_{I_{w^{-1}w_0}}(u_0) z_{I_v}^\Delta$$

by point (3) of Lemma 5.3. All the $z_{I_w}^\Delta$ are thus in the image of the characteristic map by decreasing induction on the length of w . The same proof works when replacing Δ by C . \square

Still assuming that \mathfrak{t} is regular in R , let $f : R \rightarrow R'$ be a ring morphism sending a formal group law F over R to a formal group law F' over R' . By restriction (through f), $\mathcal{H}_{R'}(M)_{F'}$ is a R -module. Let $f_{\mathcal{H}} : \mathcal{H}(M)_F \rightarrow \mathcal{H}_{R'}(M)_{F'}$ be the R -linear map sending the element $z_{I_w}^\Delta$ in $\mathcal{H}(M)_F$ to the corresponding one in $\mathcal{H}_{R'}(M)_{F'}$.

6.5. Proposition. We have the following:

- (1) The assignment $f \rightarrow f_{\mathcal{H}}$ defines a functor from the category of formal group laws (R, F) (\mathfrak{t} regular in R) to the category of pairs (R, M) where M is a module over R , and the morphisms are the obvious ones.
- (2) We have $f_{\mathcal{H}} \mathfrak{c}^F = \mathfrak{c}^{F'} f_*$.
- (3) The map $f_{\mathcal{H}}$ defined above is independent of the choice of the reduced decompositions I_w .

Proof. Point (1) holds by construction. Point (2) follows from the formula for the characteristic map given in Theorem 6.3. Since f_* is defined independently of the I_w , the last item (3) holds if the characteristic map is surjective. The general case follows by embedding $\mathcal{H}_R(M)_F$ into $\mathcal{H}_{R[1/\mathfrak{t}]}(M)_F$ where the characteristic map is surjective by Theorem 6.4. \square

6.6. Proposition. *Assume that \mathfrak{t} is invertible in R . Fix a reduced decomposition I_w for each $w \in W$. Then*

(1) *For any $x \in R[[M]]_F$, the system of linear equations in $R[[M]]_F$*

$$\Delta_{I_v}(x) = \sum_{w \in W} r_w \Delta_{I_v} \Delta_{I_w}(u_0)$$

for all v in W , has a unique solution $(r_w)_{w \in W}$.

(2) *If $(r_w)_{w \in W}$ is the solution of (1), then for any $D \in \mathcal{D}(M)_F$, we have the equality*

$$D(x) = \sum_{w \in W} r_w D \Delta_{I_w}(u_0).$$

(3) *If $(r_w)_{w \in W}$ is the solution of (1) and if R has no 2-torsion, then all the r_w are in fact in $R[[M]]_F^W$.*

The same is true when replacing Δ by C everywhere in points (1), (2) and (3).

Proof. Let A be a ring and \mathcal{I} be an ideal of A contained in its Jacobson radical. A matrix with coefficients in A is invertible if and only if the corresponding matrix with coefficients in A/\mathcal{I} is invertible. Since \mathcal{I}_F is contained in the radical of $R[[M]]_F$, it suffices to show that the matrix of the system in $R[[M]]_F/\mathcal{I}_F$ is invertible. If we order the v 's by increasing length and the w 's by decreasing length, Lemma 5.3 shows that the matrix is lower triangular with \mathfrak{t} on the diagonal, and it is therefore invertible. Point (2) follows from point (1) since the Δ_{I_w} (resp. the C_{I_w}) form a basis of $\mathcal{D}(M)_F$ as an $R[[M]]_F$ -module by Theorem 4.11. Let us prove point (3). For any simple root α and for any $v \in W$, we have

$$\begin{aligned} \Delta_\alpha \Delta_{I_v}(x) &= \sum_{w \in W} \Delta_\alpha(r_w \Delta_{I_v} \Delta_{I_w}(u_0)) \\ &= \sum_{w \in W} r_w \Delta_\alpha \Delta_{I_v} \Delta_{I_w}(u_0) + \sum_{w \in W} \Delta_\alpha(r_w) s_\alpha \Delta_{I_v} \Delta_{I_w}(u_0) \end{aligned}$$

using point (4) of Proposition 3.8. But we also have

$$\Delta_\alpha \Delta_{I_v}(x) = \sum_{w \in W} r_w \Delta_\alpha \Delta_{I_v} \Delta_{I_w}(u_0)$$

by point (2) with $D = \Delta_\alpha \Delta_{I_v}$. So

$$\sum_{w \in W} \Delta_\alpha(r_w) s_\alpha \Delta_{I_v} \Delta_{I_w}(u_0) = 0$$

for any $v \in W$. Applying the ring automorphism s_α , we obtain that the $s_\alpha \Delta_\alpha(r_w)$ are solution of the system (1) with $x = 0$ and are therefore 0 by uniqueness. Thus $\Delta_\alpha(r_w) = 0$ and r_w is fixed by s_α for any simple root α by Lemma 3.14 hence by the whole Weyl group W , since it is generated by the simple reflections.

Point (3) for C is proved exactly in the same way (still using Δ_α , not C_α). \square

6.7. Theorem. *Assume that the torsion index \mathfrak{t} is invertible in R and that R has no 2-torsion. Choose a reduced decomposition I_w for every $w \in W$. The elements $\Delta_{I_w}(u_0)$ (resp. $C_{I_w}(u_0)$) form a basis of $R[[M]]_F$ as an $R[[M]]_F^W$ -module.*

Proof. We need to show that any x can be decomposed in a unique way as $x = \sum_{w \in W} r_w \Delta_{I_w}(u_0)$. Note that this is the row $v = 1$ of the system (1) of Proposition 6.6. By $R[[M]]_F^W$ -linearity of any Δ_{I_v} (see Prop. 3.13), if that row is satisfied with coefficients r_w in $R[[M]]_F^W$, the rest of the system is satisfied, so this proves uniqueness. Existence of the decomposition then follows from points (1) and (3) of the proposition. The same proof goes through with the $C_{I_w}(u_0)$. \square

6.8. Remark. The previous theorem for the (uncompleted) symmetric algebra, i.e. the additive case, is [7, §6, Théorème 2, (c)], but the proof given there is incorrect: in the notation of loc. cit., the ideal I is only known a priori to be of the right form to apply the graded Nakayama lemma when it is tensored by \mathbb{Q} , a fact that cannot be assumed, this is the whole point of the theorem. When contacted by one of the authors, Demazure kindly and quickly supplied another proof which we adapted to our setting in Proposition 6.6. There is a slight difference: in the symmetric algebra case, the matrix of the system is upper triangular with diagonal \mathfrak{t} and is therefore invertible, whereas in our case, the lower triangular part does not vanish because our formal group law is not additive, but the strictly upper triangular part is in the radical because our ring is complete.

6.9. Theorem. *Assume that the torsion index \mathfrak{t} is invertible in R and that R has no 2-torsion. Then the kernel of the characteristic map $\mathfrak{c} : R[[M]]_F \rightarrow \mathcal{H}(M)_F$ is the ideal of $R[[M]]_F$ generated by elements in \mathcal{I}_F^W .*

Proof. By Remark 6.2, the ideal generated by \mathcal{I}_F^W is included in $\ker \mathfrak{c}$. Conversely, let $x \in \ker \mathfrak{c}$, and decompose it as $\sum_{w \in W} r_w \Delta_{I_w}(u_0)$ with the $r_w \in R[[M]]_F^W$ by the previous theorem. We then have $0 = \mathfrak{c}(x) = \sum_{w \in W} \epsilon(r_w) \mathfrak{c}(\Delta_{I_w}(u_0))$. But the $\mathfrak{c}(\Delta_{I_w}(u_0))$ form an R -basis of the image of \mathfrak{c} by Theorem 6.4, so $\epsilon(r_w) = 0$ for all $w \in W$ and x is the ideal generated by \mathcal{I}_F^W . \square

7. THE PRODUCT STRUCTURE OF THE COHOMOLOGY

In this section, we explain how to define a product on $\mathcal{H}(M)_F$ in order that the characteristic map be a ring homomorphism. Then, we study the structure of $\mathcal{H}(M)_F$.

7.1. Lemma. *For any $D \in \mathcal{D}(M)_F$,*

- (1) *there is a finite family (D_i, D'_i) of elements of $\mathcal{D}(M)_F$ such that $D(uv) = \sum_i D_i(u)D'_i(v)$ for any elements $u, v \in R[[M]]_F$.*
- (2) *Furthermore if D is in $\mathcal{D}(M)_F^{(m)}$, each D_i and D'_i can be chosen in $\mathcal{D}(M)_F^{(m_i)}$ and $\mathcal{D}(M)_F^{(m'_i)}$ respectively, such that $m_i + m'_i \geq m$.*

Proof. To show (1), the elements of $\mathcal{D}(M)_F$ satisfying a statement form a sub- $R[[M]]_F$ -algebra of $\mathcal{D}(M)_F$, and it is clear for generators of $\mathcal{D}(M)_F$ by equation (4) of Prop. 3.8. This same equation also proves (2) for generators by induction. \square

Let $R[[M]]_F^{\circ\vee}$ be the continuous R -dual of $R[[M]]_F$, i.e. the set of R -linear morphisms f from $R[[M]]_F$ to R such that $f(\mathcal{I}_F^i) = 0$ for some i (depending on f).

7.2. Lemma. *The R -module $R[[M]]_F^{\circ\vee}$ is flat. When \mathfrak{t} is regular in R , the R -module $\epsilon\mathcal{D}(M)_F$ is also flat.*

Proof. For any i , the R -module $\mathcal{I}_F^i/\mathcal{I}_F^{i-1}$ is a free R -module since it is isomorphic to $S_R^i(M)$. Thus, by induction, for any i , the quotient $R[[M]]_F/\mathcal{I}_F^i$ is a finitely generated free R -module, thus flat. Now $R[[M]]_F^{\circ\vee}$ is the direct limit of the $(R[[M]]_F/\mathcal{I}_F^i)^\vee$ (usual R -duals) and it is therefore flat. When \mathfrak{t} is regular, the module $\epsilon\mathcal{D}(M)_F$ is a finitely generated free R -module by 5.4, so it is flat. \square

7.3. Theorem. *Let \mathfrak{t} be regular in R . There is a unique ring structure on $\mathcal{H}(M)_F$ such that the characteristic map $\mathfrak{c} : R[[M]]_F \rightarrow \mathcal{H}(M)_F$ is a ring morphism.*

Proof. Since $R \rightarrow R[1/\mathfrak{t}]$ is an injection and $\mathcal{H}(M)_F$ is a free R -module, we can check uniqueness after inverting \mathfrak{t} , in which case it is obvious since the characteristic map is surjective. Let us now prove the existence of this product. The R -module $R[[M]]_F^{\circ\vee}$ has a natural structure of coalgebra induced by the collection of maps $(R[[M]]_F/\mathcal{I}^i)^\vee \rightarrow (R[[M]]_F/\mathcal{I}^i)^\vee \otimes (R[[M]]_F/\mathcal{I}^i)^\vee$ dual to the product, and taking direct limits. The inclusion $\epsilon\mathcal{D}(M)_F \subseteq R[[M]]_F^{\circ\vee}$ induces the diagonal map $\epsilon\mathcal{D}(M)_F \otimes_R \epsilon\mathcal{D}(M)_F \rightarrow R[[M]]_F^{\circ\vee} \otimes_R R[[M]]_F^{\circ\vee}$ which is injective by Lemma 7.2. To show that $\epsilon\mathcal{D}(M)_F$ is a subcoalgebra of $R[[M]]_F^{\circ\vee}$, it therefore suffices to show that the composition $\epsilon\mathcal{D}(M)_F \rightarrow R[[M]]_F^{\circ\vee} \rightarrow R[[M]]_F^{\circ\vee} \otimes_R R[[M]]_F^{\circ\vee}$ factors through the image of $\epsilon\mathcal{D}(M)_F \otimes_R \epsilon\mathcal{D}(M)_F$ in $R[[M]]_F^{\circ\vee} \otimes_R R[[M]]_F^{\circ\vee}$. This is ensured by Lemma 7.1. Now $\mathcal{H}(M)_F$ is the dual of a coalgebra, and is therefore an R -algebra. By construction, the characteristic map is a ring homomorphism. \square

7.4. We now filter the ring $\mathcal{H}(M)_F$ by setting that $\mathcal{H}(M)_F^{(i)}$ is the set of forms on $\epsilon\mathcal{D}(M)_F$ that are zero on $\epsilon\mathcal{D}(M)_F^{(-i+1)}$. By Proposition 5.6, we therefore have $\mathcal{H}(M)_F^{(0)} = \mathcal{H}(M)_F$ and $\mathcal{H}(M)_F^{(N+1)} = 0$. In other words, the filtration has the form

$$\mathcal{H}(M)_F = \mathcal{H}(M)_F^{(0)} \supseteq \mathcal{H}(M)_F^{(1)} \supseteq \cdots \supseteq \mathcal{H}(M)_F^{(N)} \supseteq 0.$$

We consider the augmentation map $\varepsilon : \mathcal{H}(M)_F \rightarrow R$ defined as the evaluation at $\epsilon \in \epsilon\mathcal{D}(M)_F$.

7.5. Proposition. *Assume \mathfrak{t} is regular in R . We have:*

- (1) $\mathcal{H}(M)_F^{(1)}$ is the kernel of ε .
- (2) The filtration is compatible with the product:

$$\mathcal{H}(M)_F^{(j)} \mathcal{H}(M)_F^{(m)} \subseteq \mathcal{H}(M)_F^{(j+m)}$$

- (3) The characteristic map \mathfrak{c} is a morphism of filtered rings.

Proof. Point (1) is clear since ϵ generates $\mathcal{D}(M)_F^{(0)}$. Let $h_1 \in \mathcal{H}(M)_F^{(j)}$ and $h_2 \in \mathcal{H}(M)_F^{(m)}$. The product $h = h_1 \cdot h_2$ is defined as $h(\epsilon D) = \sum_i h_1(\epsilon D_i) h_2(\epsilon D'_i)$ where the (D_i, D'_i) are as in Lemma 7.1. By the same lemma, if $D \in \mathcal{D}(M)_F^{(j+m+1)}$, they can be chosen such that $D_i \in \mathcal{D}(M)_F^{j_i}$ and $D'_i \in \mathcal{D}(M)_F^{m_i}$ with $j_i + m_i \geq j + m + 1$. Thus, for any i , either $j_i \geq j + 1$ or $m_i \geq m + 1$, so every term in the sum is zero and $h \in \mathcal{H}(M)_F^{(j+m)}$. For point (3), take an element u in \mathcal{I}_F^i . By definition, $\mathfrak{c}(u)$ is the evaluation at u in $\epsilon\mathcal{D}(M)_F^\vee = \mathcal{H}(M)_F$ so by Prop. 5.6.(1), $\mathfrak{c}(u)$ is in $\mathcal{H}(M)_F^{(i)}$. \square

7.6. Proposition. *Let I_w be choices of reduced decompositions for every $w \in W$. Assume the torsion index \mathfrak{t} is regular in R . Then*

- (1) For any i , the elements $z_{I_w}^\Delta$ (resp. $z_{I_w}^C$) with $l(w) \geq i$ generate $\mathcal{H}(M)_F^{(i)}$.
- (2) The elements $z_{I_{w_0}}^\Delta$ and $z_{I_{w_0}}^C$ of Theorem 6.3 do not depend on the choices of decompositions I_w and are equal up to the sign $(-1)^N$. The element $z_{I_{w_0}}^C$ is denoted by z_0 and for any u_0 as in section 5, we have $\mathfrak{c}(u_0) = (-1)^N \mathfrak{t}z_0$.

Proof. Point (1) follows from point (3) of Proposition 5.6 since the $z_{I_w}^\Delta$ (resp. $z_{I_w}^C$) are the dual basis to the $\epsilon\Delta_{I_w}$ (resp. ϵC_{I_w}). By Theorem 6.3, we have $\mathfrak{c}(u_0) = \mathfrak{t}z_{I_{w_0}}^\Delta$. But the left hand side is independent of the choices of the I_w and the right hand side is independent of the choice of u_0 , which proves the claim. We also have $\mathfrak{c}(u_0) = (-1)^N \mathfrak{t}z_0$. \square

7.7. We now consider operators on $\mathcal{H}(M)_F$. For any root α , multiplication on the right by C_α (resp. Δ_α) defines an endomorphism on $\epsilon\mathcal{D}(M)_F$, and therefore one on $\mathcal{H}(M)_F$ by duality. This operator is denoted by A_α (resp. B_α). In other words, for any $h \in \mathcal{H}(M)_F$ and $\epsilon D \in \epsilon\mathcal{D}(M)_F$, we have

$$A_\alpha(h)(\epsilon D) = h(\epsilon DC_\alpha) \quad \text{and} \quad B_\alpha(h)(\epsilon D) = h(\epsilon D\Delta_\alpha)$$

As for the operators Δ and C , we use the notation A_I and B_I for a sequence I of simple reflections.

7.8. Proposition. *When \mathfrak{t} is regular in R , these operators satisfy:*

- (1) $A_\alpha \circ \mathfrak{c}(u) = \mathfrak{c} \circ C_\alpha(u)$ and $B_\alpha \circ \mathfrak{c}(u) = \mathfrak{c} \circ \Delta_\alpha(u)$.
- (2) $A_\alpha \mathcal{H}(M)_F^{(i)} \subseteq \mathcal{H}(M)_F^{(i-1)}$ and $A_\alpha \mathcal{H}(M)_F^{(i)} \subseteq \mathcal{H}(M)_F^{(i-1)}$
- (3) For any reduced decomposition I_0 of w_0 , the element $A_{I_0}(z_0)$ (resp. $B_{I_0}(z_0)$) is invertible in $\mathcal{H}(M)_F$.
- (4) If I is nonreduced and of length N , then the element $A_I(z_0)$ (resp. $B_I(z_0)$) is in $\mathcal{H}(M)_F^{(1)} = \ker \varepsilon$.

Proof. Point (1) follows directly from the definitions of the characteristic map and of the operators. Point (2) follows from point (1) of Proposition 4.6. For point (3), first note that by Proposition 7.5, an element $h \in \mathcal{H}(M)_F$ is invertible if and only if $\varepsilon(h)$ is invertible in R . Now

$$\mathfrak{t} \varepsilon A_{I_0}(z_0) = \varepsilon A_{I_0}(\mathfrak{t}z_0) = \varepsilon A_{I_0} \mathfrak{c}(u_0) = \varepsilon \mathfrak{c}(C_{I_0}(u_0)) = \varepsilon C_{I_0}(u_0) = \mathfrak{t}.$$

Therefore, $\varepsilon A_{I_0}(z_0) = 1$. The same proof works for $\varepsilon B_{I_0}(z_0)$. For point (4), the same series of equalities is used, except that the last term is zero by Prop. 5.3. \square

7.9. Proposition. *Let I_w be a choice of reduced decompositions for all $w \in W$. Assume \mathfrak{t} is regular in R . Then the elements $A_{I_w}(z_0)$ (resp. $B_{I_w}(z_0)$) with $l(w) \leq N - i$ form an R -basis of $\mathcal{H}(M)_F^{(i)}$.*

Proof. We prove it for the $A_{I_w}(z_0)$, the proof for the $B_{I_w}(z_0)$ is similar. First note that by Proposition 7.6, z_0 is the unique element of $\mathcal{H}(M)_F$ such that $z_0(\epsilon C_{I_w}) = \delta_{w, w_0}$. Thus, if $l(w) + l(v) \leq N$, we have $A_{I_w}(z_0)(\epsilon C_{I_v}) = \delta_{w, v^{-1}w_0}$. In other words, if we decompose A_{I_w} on the basis of the z_{I_v} , the coordinate on z_{I_v} is $\delta_{w, v^{-1}w_0}$ if $l(v) + l(w) \leq N$. Thus, the $A_{I_w}(z_0)$ with $l(w) \leq N - i$ can be expressed as linear combinations of the z_{I_v} with $l(v) \geq i$ and the matrix of their expressions is unitriangular up to a correct ordering of rows and columns and is therefore invertible. \square

7.10. Proposition. *Assume \mathfrak{t} is regular in R and R' , respectively endowed with formal group laws F and F' . Let $f : R \rightarrow R'$ be a ring morphism sending F to F' . Then the morphism $f_{\mathcal{H}} : \mathcal{H}_R(M)_F \rightarrow \mathcal{H}_{R'}(M)_{F'}$ introduced before Proposition 6.5 is actually a morphism of filtered rings. It satisfies*

$$A_I^{F'} f_{\mathcal{H}} = f_{\mathcal{H}} A_I^F \quad \text{and} \quad B_I^{F'} f_{\mathcal{H}} = f_{\mathcal{H}} B_I^F$$

for any sequence I .

Proof. Both facts are clear by Point 2 of 6.5 when the characteristic map is surjective, and therefore when \mathfrak{t} is regular by extending scalars to $R[1/\mathfrak{t}]$. \square

Part 2. Bott-Samelson resolutions for oriented theories.

8. ALGEBRAIC COBORDISM AND ORIENTED COHOMOLOGY THEORIES

In the present section we recall the notion of *oriented* cohomology theory and the notion of algebraic cobordism Ω following the book of Levine and Morel [13]. As

main examples of oriented theories we consider the Chow ring CH , Grothendieck's \mathbf{K}^0 and connective K -theory \mathbf{k} .

8.1. According to [13, §1.1] an *oriented cohomology theory* \mathbf{h} is a contravariant functor from the category of smooth varieties over a field k of an arbitrary characteristic to the category of graded rings satisfying the standard cohomological axioms: localization, homotopy invariance and push-forwards for projective morphisms (see [13, Def.1.1.2]).

A universal theory Ω satisfying these properties was constructed in [13, §2] assuming the base field k has characteristic 0. It is called *algebraic cobordism*. An element of codimension i in $\Omega(X)$, i.e. in $\Omega^i(X)$, is additively generated by classes $[Y \rightarrow X]$ of projective maps of codimension i from smooth schemes Y .

For any map $f: X_1 \rightarrow X_2$ the induced functorial map $f^*: \Omega^i(X_2) \rightarrow \Omega^i(X_1)$ is called the *pull-back*. For a projective map $f: X_1 \rightarrow X_2$, there is a corresponding *push-forward* map $f_*: \Omega(X_1) \rightarrow \Omega(X_2)$ shifting the cohomological degree by $\dim X_2 - \dim X_1$ when X_1 and X_2 are equidimensional and given by $[Y_1 \rightarrow X_1] \mapsto [Y_1 \rightarrow X_1 \xrightarrow{f} X_2]$.

8.2. An oriented cohomology theory \mathbf{h} comes with a *formal group law* F over the coefficient ring $\mathbf{h}(\text{pt})$ such that

$$F(c_1^{\mathbf{h}}(\mathcal{L}_1), c_2^{\mathbf{h}}(\mathcal{L}_2)) = c_1^{\mathbf{h}}(\mathcal{L}_1 \otimes \mathcal{L}_2),$$

where \mathcal{L}_1 and \mathcal{L}_2 are lines bundles on X and $c_1^{\mathbf{h}}$ denotes the first Chern class in the cohomology theory \mathbf{h} (see [13, Cor.4.1.8]). For Ω , the associated formal group law U

$$U(x, y) = x + y + \sum_{i, j \geq 1} a_{ij} x^i y^j, \quad \text{where } a_{ij} \in \Omega(\text{pt}).$$

turns out to be universal. The ring of coefficients $\Omega(\text{pt})$ is generated by the coefficients a_{ij} and coincides with the classical *Lazard ring* \mathbb{L} .

There is a canonical map $pr_{\mathbf{h}}: \Omega \rightarrow \mathbf{h}$ sending the coefficients of U to the corresponding coefficients of F , and hence inducing a morphism of formal group laws.

8.3. **Example.** Consider the Chow ring $\text{CH}(X)$ of algebraic cycles on X modulo rational equivalence. According to [13, Thm. 4.5.1] the canonical map $pr_{\text{CH}}: \Omega \rightarrow \text{CH}$ induces an isomorphism $\Omega \otimes_{\mathbb{L}} \mathbb{Z} \xrightarrow{\cong} \text{CH}$ of oriented cohomology theories. In particular, $pr_{\text{CH}}: \Omega \rightarrow \text{CH}$ is surjective and its kernel is generated by $\mathbb{L}_{>0}$, the subgroup of elements of positive dimension in the Lazard ring. Observe that pr_{CH} restricted to the coefficient rings $\mathbb{L} \rightarrow \mathbb{Z} = \text{CH}(\text{pt})$ coincides with the augmentation map. The associated formal group law, denoted by F_0 , is called the *additive formal group law* and is given by $F_0(x, y) = x + y$.

8.4. **Example.** Consider the oriented cohomology theory $\mathbf{K}(X) = \mathbf{K}^0(X)[\beta, \beta^{-1}]$, where $\mathbf{K}^0(X)$ is the Grothendieck \mathbf{K}^0 of X . Observe that $\mathbf{K}(\text{pt}) = \mathbb{Z}[\beta, \beta^{-1}]$. According to [13, Cor.4.2.12] the canonical map $pr_{\mathbf{K}}: \Omega \rightarrow \mathbf{K}$ sending $[\mathbb{P}^1]$ to β induces an isomorphism $\Omega \otimes_{\mathbb{L}} \mathbb{Z}[\beta, \beta^{-1}] \rightarrow \mathbf{K}$ of oriented cohomology theories. The associated formal group law, denoted by F_{β} , is called a *multiplicative periodic formal group law* and is given by $F_{\beta}(x, y) = x + y - \beta xy$.

8.5. **Example.** Consider a ring homomorphism $\mathbb{L} \rightarrow \mathbb{Z}[v]$ given by $[\mathbb{P}^1] = -a_{11} \mapsto v$ and $a_{ij} \mapsto 0$ for $(i, j) \neq (1, 1)$. We define a new cohomology theory, called a *connective K-theory*, by $\mathbf{k} = \Omega \otimes_{\mathbb{L}} \mathbb{Z}[v]$ (see [13, §4.3.3]). Its formal group law is denoted by F_v and is given by $F_v(x, y) = x + y - vxy$. Observe that contrary to Example 8.4 the element v is non-invertible in the coefficient ring $\mathbf{k}(\text{pt}) = \mathbb{Z}[v]$.

8.6. Remark. Avoiding the localization axiom one obtains a more general notion of oriented cohomology theory studied by Merkurjev in [14]. In particular, he showed that taking the so-called ‘tilde operation’ of CH one obtains a theory $\tilde{\text{CH}}$, defined over a field of any characteristic, which serves as a good replacement for algebraic cobordism in all questions related with cohomological operations. Note that the Lazard ring \mathbb{L} injects into the coefficient ring $\tilde{\text{CH}}(\text{pt})$.

We come now to the following important property of an algebraic cohomology theory:

8.7. Definition. We say that a cohomology theory is *weakly birationally invariant* if for any proper birational morphism $f: Y \rightarrow X$ the push-forward of the fundamental class $f_*(1_Y)$ is invertible.

8.8. Example. Any *birationally invariant* theory, i.e. such that $f_*(1_Y) = 1_X$ for any proper birational $f: Y \rightarrow X$, is weakly birationally invariant. The Chow ring CH considered over an arbitrary field and the K -theory \mathbf{K} considered over a field of characteristic 0 provide examples of birationally invariant theories. According to [13, Thm.4.3.9] the connective K -theory \mathbf{k} defined over a field of characteristic 0 is universal among all *birationally invariant* theories. In particular, the kernel of the canonical map $\mathbb{L} \rightarrow \mathbf{k}(\text{pt})$ is generated by differences of classes $[X] - [X']$, where X and X' are birationally equivalent.

8.9. Lemma. *For any smooth variety X over a field of characteristic 0, the kernel of the natural map $\Omega^n(X) \rightarrow \text{CH}^n(X)$ consists of nilpotent elements if $n \geq 0$ and is trivial if $n = \dim X$.*

Proof. According to [13, Rem. 4.5.6] the kernel is additively generated by products of the form ab , $a \in \mathbb{L}^{-i}$, $b \in \Omega^{n+i}(X)$, $i > 0$. Now the claim follows from the fact that $\Omega^j(X) = 0$ when $j > \dim X$. \square

8.10. Corollary. *Over a field of characteristic 0, any oriented cohomology theory in the sense of Levine-Morel is weakly birationally invariant.*

Proof. Consider the algebraic cobordism Ω . Let $f: Y \rightarrow X$ be a proper birational map. The element $f_*^\Omega(1_Y) - 1_X$ is in the kernel of the map $\Omega^0(X) \rightarrow \text{CH}^0(X)$. By Lemma 8.9 the difference $f_*^\Omega(1_Y) - 1_X$ is nilpotent, therefore, $f_*^\Omega(1_Y)$ is invertible. The statement for an arbitrary theory follows by the universality of Ω . \square

9. A SEQUENCE OF SPLIT \mathbb{P}^1 -BUNDLES

In the present section we compute the oriented cohomology \mathbf{h} of a variety obtained as a sequence of split \mathbb{P}^1 -bundles. The main tool is the projective bundle theorem for \mathbf{h} . Observe that all formulas are given in terms of pull-backs and push-forwards of fundamental classes. This invariant description will play an important role in the sequel.

9.1. Lemma. *Let X be a smooth projective variety over a field k . Let $p: \mathbb{P}_X(\mathcal{E}) \rightarrow X$ be the projective bundle of a vector bundle \mathcal{E} of rank 2 over X . Assume that p has a section σ . Then there is a ring isomorphism*

$$\mathbf{h}(\mathbb{P}_X(\mathcal{E})) \simeq \mathbf{h}(X)[\xi]/(\xi^2 - y\xi), \text{ where } \xi = \sigma_*(1_X) \text{ and } y = p^*\sigma^*\xi.$$

Proof. Consider the canonical embedding $\mathcal{O}_{\mathcal{E}}(-1) \rightarrow p^*\mathcal{E}$, where $\mathcal{O}_{\mathcal{E}}(-1)$ is a tautological line bundle over $\mathbb{P}_X(\mathcal{E})$ (see [9, B.5.5]). Let \mathcal{L} denote the quotient $\mathcal{E}/\sigma^*\mathcal{O}_{\mathcal{E}}(-1)$. By Cartan formula (see [16, Def. 3.26.(3)]) we have

$$c_1^{\mathbf{h}}(\mathcal{E}) = c_1^{\mathbf{h}}(\sigma^*\mathcal{O}_{\mathcal{E}}(-1)) + c_1^{\mathbf{h}}(\mathcal{L}) \quad \text{and} \quad c_2^{\mathbf{h}}(\mathcal{E}) = c_1^{\mathbf{h}}(\sigma^*\mathcal{O}_{\mathcal{E}}(-1)) \cdot c_1^{\mathbf{h}}(\mathcal{L}).$$

According to the projective bundle theorem (see [16, Thm. 3.9 and formula (17)]) applied to p , there is a ring isomorphism

$$h(\mathbb{P}_X(\mathcal{E})) \simeq h(X)[t]/(t-a)(t-b),$$

where $a = p^*c_1^h(\sigma^*\mathcal{O}_{\mathcal{E}}(-1))$, $b = p^*c_1^h(\mathcal{L})$ and $t = c_1^h(\mathcal{O}_{\mathcal{E}}(-1))$.

Consider the elements $\xi = b -_F t$ and $y = b -_F a$, where F is the formal group law corresponding to h . Observe that by [15, 2.2.9] $\xi = c_1^h(\mathcal{O}_{\mathcal{E}}(1) \otimes p^*\mathcal{L}) = \sigma_*(1_X)$ and by the very definition $y = c_1^h(p^*\sigma^*\mathcal{O}_{\mathcal{E}}(1) \otimes p^*\mathcal{L}) = p^*\sigma^*\xi$.

Since changing t to ξ induces an automorphism of $h(\mathbb{P}_X(\mathcal{E}))$, we only need to prove the relation $\xi^2 - y\xi = 0$. Note that for any power series f , we may write $f(x) - f(y) = (x-y)f'(x, y)$ where $f'(x, y)$ is again a power series. Hence, we have $b -_F t = (b -_F t) - (t -_F t) = (t-b)f'_1(b, t)$ and $(b -_F t) - (b -_F a) = (t-a)f'_2(a, b, t)$ for some power series f'_1 and f'_2 . Therefore

$$\xi(\xi - y) = (b -_F t)((b -_F t) - (b -_F a)) = (t-b)(t-a)f'_1(b, t)f'_2(a, b, t) = 0$$

and the proof is finished. \square

9.2. Corollary. *In the notation of Lemma 9.1 we have the following formula for the push-forward of the fundamental class*

$$p_*(1_{\mathbb{P}_X(\mathcal{E})}) = \sigma^*(\xi^{-1} + (-_F\xi)^{-1}).$$

Proof. By [20, Thm. 5.30] we have

$$p_*(1_{\mathbb{P}_X(\mathcal{E})}) = (\sigma^*b -_F \sigma^*a)^{-1} + (\sigma^*a -_F \sigma^*b)^{-1},$$

where σ^*a and σ^*b are the roots of the bundle \mathcal{E} as in the proof of Lemma 9.1. Since $y = b -_F a = p^*\sigma^*\xi$ we obtain the desired formula. \square

9.3. Theorem. *More generally, let X be a variety obtained by means of a sequence of split \mathbb{P}^1 -bundles, i.e. there is a sequence of varieties X_i , $0 \leq i \leq N$, starting from a point $X_0 = \text{pt}$ and finishing at $X_N = X$ such that for each $1 \leq i \leq N$*

$$p_i: X_i \simeq \mathbb{P}_{X_{i-1}}(\mathcal{E}_i) \rightarrow X_{i-1}$$

is a projective bundle with a section σ_i , where \mathcal{E}_i is some vector bundle of rank 2 over X_{i-1} . Then there is a ring isomorphism

$$h(X_N) \simeq h(\text{pt})[\xi_1, \dots, \xi_N]/I,$$

where I is an ideal generated by elements $\{\xi_i^2 - y_i\xi_i\}_{i=1\dots N}$, $\xi_i = p^\sigma_{i*}(1_{X_{i-1}})$, $y_i = p^*\sigma_i^*\sigma_{i*}(1_{X_{i-1}})$ and p^* denotes the pull-back on X_N .*

Proof. Follows by induction using Lemma 9.1. \square

9.4. Remark. Observe that the element y_i appearing in the relations is not necessarily in $h(\text{pt})$. To obtain a complete answer in terms of generators ξ_1, \dots, ξ_N and relations with coefficients in $h(\text{pt})$ one has to express the elements y_i in terms of the ‘previous’ generators ξ_1, \dots, ξ_{i-1} . In the next section, we show how to obtain such an expression for Bott-Samelson varieties using the characteristic map.

9.5. Example. Let X be as in Lemma 9.1. By [9, 3.2.11] we have the following formula for the total Chern class of the tangent bundle of $\mathbb{P}_X(\mathcal{E})$:

$$c^h(\mathcal{T}_{\mathbb{P}_X(\mathcal{E})}) = c^h(p^*\mathcal{T}_X) \cdot c^h(\mathcal{O}_{\mathcal{E}}(1) \otimes p^*\mathcal{E}).$$

Since there is an exact sequence

$$0 \rightarrow \mathcal{O}_{\mathcal{E}}(1) \otimes p^*\sigma^*\mathcal{O}_{\mathcal{E}}(-1) \rightarrow \mathcal{O}_{\mathcal{E}}(1) \otimes p^*\mathcal{E} \rightarrow \mathcal{O}_{\mathcal{E}}(1) \otimes p^*\mathcal{L} \rightarrow 0,$$

by Cartan formula we obtain

$$c^h(\mathcal{O}_{\mathcal{E}}(1) \otimes p^*\mathcal{E}) = c^h(\mathcal{O}_{\mathcal{E}}(1) \otimes p^*\mathcal{L}) \cdot c^h(\mathcal{O}_{\mathcal{E}}(1) \otimes p^*\sigma^*\mathcal{O}_{\mathcal{E}}(-1)).$$

Since $\mathcal{O}_{\mathcal{E}}(1) \otimes p^* \sigma^* \mathcal{O}_{\mathcal{E}}(-1) = (\mathcal{O}_{\mathcal{E}}(1) \otimes p^* \mathcal{L}) \otimes (p^* \sigma^* \mathcal{O}_{\mathcal{E}}(-1) \otimes p^* \mathcal{L}^\vee)$, we obtain that

$$c^h(\mathcal{T}_{\mathbb{P}_X(\mathcal{E})}) = c^h(p^* \mathcal{T}_X) \cdot (1 + \xi) \cdot (1 + (\xi -_F y)).$$

More generally, for X_N from Theorem 9.3 we obtain by induction

$$(4) \quad c^h(\mathcal{T}_{X_N}) = \prod_{i=1}^N (1 + \xi_i)(1 + (\xi_i -_F y_i)).$$

10. AN ELEMENTARY STEP OF THE BOTT-SAMELSON RESOLUTION

We apply the results of the previous section to compute an oriented cohomology of a split $P/B = \mathbb{P}^1$ -bundle, where P is a semi-direct product of a split reductive linear algebraic group of semisimple rank one and a connected unipotent group, and B is a Borel subgroup of P . This example appears as an elementary step in the construction of a Bott-Samelson variety.

10.1. Consider a split connected solvable algebraic group B over a field k . Let T denote its split maximal torus. Let X be a scheme on which B acts on the right such that the quotient X/B exists and $X \rightarrow X/B$ is a principal B -bundle. Let M denote the group of characters of T . Each character $\lambda \in M$ extends to a one-dimensional representation V_λ of B and, hence, defines a line bundle $\mathcal{L}(\lambda)$ over X/B whose total space is the fiber product $X \times^B V_\lambda$, i.e. the quotient $(X \times_k V_\lambda)/B$ by means of the right B -action $(x, v)b = (xb, b^{-1}v)$ (see [1, Ch.1 §3]).

10.2. **Definition.** Let \mathfrak{h} be an oriented cohomology theory and let F be the corresponding formal group law over the coefficient ring $R = \mathfrak{h}(\text{pt})$. We define a ring homomorphism $c: R[[M]]_F \rightarrow \mathfrak{h}(X/B)$ from the formal group ring $R[[M]]_F$ to the cohomology ring $\mathfrak{h}(X/B)$ by sending a generator x_λ to the first Chern class $c_1^h(\mathcal{L}(\lambda))$ of the line bundle $\mathcal{L}(\lambda)$. This map is well-defined since all Chern classes are nilpotent and satisfy $c_1^h(\mathcal{L}(\lambda + \mu)) = c_1^h(\mathcal{L}(\lambda) \otimes \mathcal{L}(\mu)) = c_1^h(\mathcal{L}(\lambda)) +_F c_1^h(\mathcal{L}(\mu))$. It is called the *characteristic map*.

10.3. We follow the notation of [7, §2]: Let P be a semi-direct product of a connected unipotent group U and a reductive split group L of semi-simple rank 1. Let T be a maximal split torus of L and let M denote its group of characters. Let α be the one of two roots of L with respect to T , let U_α be the corresponding unipotent subgroup and let $B = T \cdot U_\alpha \cdot U$ be the Borel subgroup of P containing T . Let $s_\alpha(\lambda) = \lambda - \alpha^\vee(\lambda) \cdot \alpha$ denote the reflection corresponding to the root α .

Consider the fibered product $X' = X \times^B P$, i.e. the quotient $(X \times_k P)/B$ by means of the B -action $(x, h)b = (xb, b^{-1}h)$. By definition, X' is a principal P -bundle over X/B , all fibers of the canonical projection $p: X'/B \rightarrow X/B$ are isomorphic to $P/B \simeq \mathbb{P}^1$ and there is an obvious section $\sigma: X/B \rightarrow X'/B$ given by $x \mapsto (x, 1)$. According to [11, Exrc. 7.10.(c)] there exists a vector bundle \mathcal{E} of rank 2 over X/B such that X'/B can be identified with the projective bundle $\mathbb{P}_{X/B}(\mathcal{E})$.

10.4. **Remark.** Observe that in [7, §2] instead of the quotient X'/B the author considers the quotient of X' modulo the opposite Borel subgroup B' . This doesn't change much, since there is an obvious isomorphism $i: X'/B \rightarrow X'/B'$ induced by an automorphism of X' sending the class of (x, p) to the class of (x, pn_α) , where n_α is an element of P representing the reflection s_α . In particular, for any character λ we have $i^* \mathcal{L}'(\lambda) = \mathcal{L}'(s_\alpha(\lambda))$, where $\mathcal{L}'(\lambda)$ (resp. $\mathcal{L}'(s_\alpha(\lambda))$) is the line bundle over X'/B' (resp. over X'/B).

10.5. Applying Lemma 9.1 to the projective bundle $p: \mathbb{P}_{X/B}(\mathcal{E}) \rightarrow X/B$ with the section σ we obtain an isomorphism

$$\mathfrak{h}(X'/B) \simeq \mathfrak{h}(X/B)[\xi]/(\xi^2 - y\xi), \quad \text{where } \xi = \sigma_*(1_{X/B}) \text{ and } y = p^* \sigma^* \xi.$$

10.6. Lemma. *We have $\sigma^*\xi = c_1^h(\mathcal{L}(-\alpha)) = \mathbf{c}(x_{-\alpha})$.*

Proof. By definition of the push-forward we have $\xi = c_1^h(\mathcal{O}(D))$, where $\mathcal{O}(D)$ is a line bundle corresponding to the divisor $D = \sigma(X/B)$. The first equality then follows from [8, §2.5 Lem. 3]. The second one follows from the definition of the characteristic map \mathbf{c} . \square

10.7. Corollary. *We have $y = p^*\sigma^*\xi = p^*\mathbf{c}(x_{-\alpha})$.*

10.8. Lemma. *Consider the characteristic map $\mathbf{c}': R[[M]]_F \rightarrow \mathfrak{h}(X'/B)$. For any $u \in R[[M]]_F$, we have*

$$(1) \quad \sigma^*\mathbf{c}'(u) = \mathbf{c}(u) \quad \text{and} \quad (2) \quad \mathbf{c}'(u) = p^*\mathbf{c}(s_\alpha(u)) + p^*\mathbf{c}(\Delta_{-\alpha}(u)) \cdot \xi$$

Proof. By [8, §2.5 Lem. 1 and Prop.1], for any character $\lambda \in M$, we have two equalities

$$\sigma^*\mathcal{L}'(\lambda) = \mathcal{L}(\lambda) \quad \text{and} \quad \mathcal{L}'(s_\alpha(\lambda)) = p^*(\mathcal{L}(\lambda)) \otimes \mathcal{O}(D)^{\otimes \alpha^\vee(\lambda)},$$

where $\mathcal{L}'(\lambda)$ is the corresponding line bundle over X'/B and $\mathcal{O}(D)$ is a line bundle corresponding to the divisor $D = \sigma(X/B)$.

By the first equality, formula (1) holds for $u = x_\lambda$. It therefore holds for all $u \in R[[M]]_F$ and then for all $u \in R[[M]]_F$ by continuity.

By the second equality, we have for $u = x_\lambda$

$$\mathbf{c}'(x_{s_\alpha(\lambda)}) = p^*\mathbf{c}(x_\lambda) +_F (\alpha^\vee(\lambda)) \cdot_F \xi.$$

Note that for any power series $g(\xi)$ in a variable ξ we may write $g(\xi) = g(0) + \xi \cdot g'(\xi)$, where $g'(\xi) = (g(\xi) - g(0))/\xi$ is again a power series. Since, $\xi^2 = y\xi$, we have $g(\xi) = g(0) + \xi \cdot g'(y)$. Applying this to $g(\xi) = p^*\mathbf{c}(x_\lambda) +_F \alpha^\vee(\lambda) \cdot_F \xi$ and observing that $g(0) = p^*\mathbf{c}(x_\lambda)$ we obtain

$$\mathbf{c}'(x_{s_\alpha(\lambda)}) = p^*\mathbf{c}(x_\lambda) + \xi \cdot g'(y),$$

where $g(y) = p^*\mathbf{c}(x_\lambda +_F \alpha^\vee(\lambda) \cdot_F x_{-\alpha}) = p^*\mathbf{c}(x_{s_\alpha(\lambda)})$ by Cor. 10.7. Then by definition of the operator $\Delta_{-\alpha}$ we obtain $g'(y) = p^*(-\Delta_{-\alpha}(x_\lambda))$ and, hence, $\mathbf{c}'(x_{s_\alpha(\lambda)}) = p^*\mathbf{c}(x_\lambda) - p^*\mathbf{c}(\Delta_{-\alpha}(x_\lambda))\xi$. Since $\Delta_{-\alpha}s_\alpha = -\Delta_{-\alpha}$, applying s_α to u we obtain

$$\mathbf{c}'(x_\lambda) = p^*\mathbf{c}(s_\alpha(x_\lambda)) + p^*\mathbf{c}(\Delta_{-\alpha}(x_\lambda)) \cdot \xi.$$

For general u formula (2) then follows by induction on the degree of monomials and by continuity, using Prop. 3.8.(4) and $\xi^2 = y\xi$. \square

10.9. Remark. Following Remark 3.7, the second formula would look simpler with the other definition of Δ_α .

10.10. Proposition. *The following formulas hold for any $u \in R[[M]]_F$*

$$(1) \quad p_*(1_{X'/B}) = \mathbf{c}(C_\alpha(1)), \quad (3) \quad p^*(\mathbf{c}(C_\alpha(u))) = \mathbf{c}'(C_\alpha(u)), \\ (2) \quad p_*(\mathbf{c}'(u)) = \mathbf{c}(C_\alpha(u)), \quad (4) \quad p^*p_*(\mathbf{c}'(u)) = \mathbf{c}'(C_\alpha(u)).$$

Proof. The first formula of the proposition follows from Corollary 9.2 applied to the projective bundle $p: X'/B \rightarrow X/B$ where we identify $\sigma^*\xi$ with $\mathbf{c}(x_{-\alpha})$ according to Lemma 10.6.

By point (2) of Lemma 10.8 and the projection formula we have

$$p_*(\mathbf{c}'(u)) = \mathbf{c}(s_\alpha(u)) \cdot p_*(1_{X'/B}) + \mathbf{c}(\Delta_{-\alpha}(u)) \cdot p_*(\xi) =$$

By the first formula and the fact that $p_*(\xi) = p_*(\sigma_*(1_{X/B})) = 1_{X/B}$ it can be rewritten as

$$\mathbf{c}(s_\alpha(u)) \cdot \mathbf{c}(C_\alpha(1)) + \mathbf{c}(\Delta_{-\alpha}(u)) = \mathbf{c}(s_\alpha(u)C_\alpha(1) + \Delta_{-\alpha}(u))$$

Using the definition of C_α and point (3) of Proposition 3.12 we finally obtain

$$= \mathbf{c}(s_\alpha(u) \cdot e_\alpha + \Delta_{-\alpha}(u)) = \mathbf{c}(C_{-\alpha}(s_{-\alpha}(u))) = \mathbf{c}(C_\alpha(u)).$$

This proves the second formula.

The third formula follows from point (2) of Lemma 10.8 and the fact that $C_{-\alpha}\Delta_{-\alpha} = 0$ (see (6) of Prop. 3.12). The last formula is obtained by the composite of the second and the third one. \square

11. BOTT-SAMELSON VARIETIES

In the present section we compute an oriented cohomology of Bott-Samelson varieties. For definition and basic properties of Bott-Samelson varieties, we refer to papers [1], [17] and [5].

11.1. Let G be a split semisimple linear algebraic group over a field k , let T be a split maximal torus of G and let $T \subset B$ be a Borel subgroup. Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$, where n is the rank of G , be a set of simple roots of G . Let P_i be a minimal parabolic subgroup corresponding to a simple root α_i , i.e. the subgroup generated by unipotent subgroups U_{α_j} , $1 \leq j \leq n$ and $U_{-\alpha_i}$.

11.2. **Definition.** For an l -tuple of integers $I = (i_1, i_2, \dots, i_l)$ with $1 \leq i_j \leq n$, we define a variety X_I to be the fiber product

$$X_I = P_{i_1} \times^B P_{i_2} \times^B \dots \times^B P_{i_l}.$$

If $I = \emptyset$, then we set $X_\emptyset = \text{pt}$.

Observe that there is a natural right action of B on X_I , the quotient X_I/B exists and $X_I \rightarrow X_I/B$ is a principal B -bundle. The variety X_I/B is called a Bott-Samelson variety corresponding to I .

11.3. Let $J = (i_1, i_2, \dots, i_{l-1})$. Then $X_I = X_J \times^B P_{i_l}$, and we are in the situation of 10.3 for $P = P_{i_l}$, $\alpha = \alpha_{i_l}$, $X' = X_I$ and $X = X_J$. In particular, the projection map

$$p_I: X_I/B \rightarrow X_J/B$$

has a structure of a \mathbb{P}^1 -bundle with a section denoted by σ_I .

Observe that the structure map $\pi_I: X_I/B \rightarrow \text{pt}$ can be written as a composite of projection maps

$$X_I/B \rightarrow X_{(i_1, \dots, i_{l-1})}/B \rightarrow \dots \rightarrow X_{(i_1)}/B \rightarrow \text{pt}.$$

Hence, the variety X_I/B is obtained by means of a sequence of split \mathbb{P}^1 -bundles.

According to 10.1 for an oriented cohomology theory \mathfrak{h} with a formal group law F there is a well-defined characteristic map

$$\mathfrak{c}_I: R[[M]]_F \rightarrow \mathfrak{h}(X_I/B),$$

where $R[[M]]_F$ is the formal group ring of the group of characters M of T . Observe that $\mathfrak{c}_\emptyset = \epsilon$ is the augmentation map. Applying Theorem 9.3 and Corollary 10.7 we obtain an isomorphism

$$(5) \quad \mathfrak{h}(X_I/B) \simeq \mathfrak{h}(\text{pt})[\xi_1, \xi_2, \dots, \xi_l] / (\{\xi_j^2 - y_j \xi_j\}_{j=1 \dots l}),$$

where $y_j = p^* \mathfrak{c}_{(i_1, \dots, i_{j-1})}(x_{-\alpha_{i_j}})$ and p^* denotes the pull-back to $\mathfrak{h}(X_I/B)$.

11.4. **Theorem.** Let X_I/B be the Bott-Samelson variety corresponding to an l -tuple $I = (i_1, \dots, i_l)$. For any subset K of $[1, l]$ we define

$$\xi_K = \prod_{j \in K} \xi_j \quad \text{and} \quad \Theta_K = \Theta_1 \cdots \Theta_l, \quad \text{where } \Theta_j = \begin{cases} \Delta_{-\alpha_{i_j}} & \text{if } j \in K, \\ s_{\alpha_{i_j}} & \text{otherwise.} \end{cases}$$

Then the elements ξ_K , where K runs through all subsets of $[1, l]$, form a basis of the free $\mathfrak{h}(\text{pt})$ -module $\mathfrak{h}(X_I/B)$. As a ring $\mathfrak{h}(X_I/B)$ is a quotient of the polynomial ring $\mathfrak{h}(\text{pt})[\xi_1, \dots, \xi_l]$ modulo the relations

$$\xi_j^2 = \sum_{K \subseteq [1, j-1]} \epsilon_{\Theta_K}(x_{-\alpha_{i_j}}) \xi_K \xi_j.$$

Proof. The fact that $\mathfrak{h}(X_I/B)$ is a free $\mathfrak{h}(\text{pt})$ -module with a basis ξ_K follows by the projective bundle theorem, since X_I/B is obtained by means of a sequence of split \mathbb{P}^1 -bundles. To obtain the relations, we first use Lemma 10.8.(2) to prove by induction that

$$(6) \quad \mathfrak{c}_{(i_1, \dots, i_j)}(u) = \sum_{K \subseteq [1, j]} \epsilon_{\Theta_K}(u) \xi_K.$$

Using this we then compute y_j and plug the result into the formula (5). \square

The following lemma provides a formula for the push-forward of the structure map $\pi_I: X_I/B \rightarrow \text{pt}$:

11.5. Lemma. *Consider the variety X_I/B for a tuple $I = (i_1, \dots, i_l)$. For any $u \in R[[M]]_F$ we have*

$$\pi_{I*}(\mathfrak{c}_I(u)) = \epsilon_{C_I}(u),$$

where $C_I = C_{\alpha_1} \dots C_{\alpha_l}$ is the composite of operators defined in 3.11.

Proof. Decomposing the structure map as $\pi_I: X_I/B \xrightarrow{p_I} X_J/B \xrightarrow{\pi_J} \text{pt}$, where $J = (i_1, \dots, i_{l-1})$ and applying Prop. 10.10.(2) we obtain

$$\pi_{I*}(\mathfrak{c}_I(u)) = \pi_{J*} p_{I*}(\mathfrak{c}_I(u)) = \pi_{J*}(\mathfrak{c}_J(C_{\alpha_l}(u))).$$

Repeating this recursively for the structure map $\pi_J: X_J/B \rightarrow \text{pt}$ and so on, we obtain the desired formula. \square

12. BOTT-SAMELSON RESOLUTIONS

In the present section we discuss the relations between Bott-Samelson varieties and Schubert varieties on G/B .

12.1. According to Bruhat decomposition the variety of complete flags G/B is a finite disjoint union of affine spaces $G/B = \coprod_{w \in W} BwB/B$, where W is the Weyl group of G . The closure of a cell BwB/B is denoted by X_w and is called a Schubert variety. Observe that X_w is not smooth in general.

Let $w = s_{i_1} s_{i_2} \dots s_{i_l}$ be a (reduced) decomposition into a product of simple reflections (here s_i denotes s_{α_i}) with $l = l(w)$ being the length of w . The multiplication map induces a morphism

$$q_I: X_I/B \rightarrow G/B, \text{ where } I = (i_1, i_2, \dots, i_l).$$

By the results of [1, Ch. 3] (see also [17, §1.7] and [5, §4]) this map factors as $q_I: X_I/B \rightarrow X_w \hookrightarrow G/B$, where the first map is surjective birational and the second is a closed embedding. Furthermore, when J is I with the last entry removed, we have $q_I \circ \sigma_I = q_J$.

12.2. Lemma. *Consider the characteristic map $\mathfrak{c}_{G/B}: R[[M]]_F \rightarrow \mathfrak{h}(G/B)$. For any $u \in R[[M]]_F$ we have*

$$q_I^*(\mathfrak{c}_{G/B}(u)) = \mathfrak{c}_I(u).$$

Proof. By definition of the map q_I and the bundle $\mathcal{L}(\lambda)$ we have $q_I^*(\mathcal{L}(\lambda)) = \mathcal{L}_I(\lambda)$, where $\mathcal{L}(\lambda)$ (resp. $\mathcal{L}_I(\lambda)$) is the line bundle over G/B (resp. over X_I/B) corresponding to a character λ . Hence, $q_I^*(\mathfrak{c}_{G/B}(x_\lambda)) = \mathfrak{c}_I(x_\lambda)$. For a general u it follows by continuity. \square

Let w_0 be the element of maximal length in W and let $X = X_{I_0}/B$ be the Bott-Samelson variety where I_0 is a reduced decomposition of w_0 .

12.3. Lemma. *Assume that h is weakly birationally invariant. Then the pull-back $q_{I_0}^*$ is injective.*

Proof. Since our theory is weakly birationally invariant, the push-forward of the fundamental class $q_{I_0*}(1)$ is invertible. The lemma now follows by projection formula. \square

12.4. Theorem (cf. [5, Prop. 3]). *Assume that h is weakly birationally invariant. Consider the quotient map $p_i: G/B \rightarrow G/P_i$, where $1 \leq i \leq n$. Then for any $u \in R[[M]]_F$ we have the equality*

$$p_i^* p_{i*}(\mathbf{c}_{G/B}(u)) = \mathbf{c}_{G/B}(C_{\alpha_i}(u))$$

Proof. We choose a reduced decomposition I_0 of w_0 in simple reflections such that the last reflection is s_i . Let J be the same tuple without the last entry. Consider the Cartesian diagram

$$(7) \quad \begin{array}{ccc} X_{I_0}/B & \xrightarrow{p_{I_0}} & X_J/B \\ q_{I_0} \downarrow & & \downarrow p_i \circ q_J \\ G/B & \xrightarrow{p_i} & G/P_i \end{array}$$

Since the horizontal map p_i is smooth, the diagram is transversal. Since $q_{I_0}^*$ is injective by Lemma 12.3, it is enough to prove the equality after applying $q_{I_0}^*$ to both sides. By base change and by transversality of the diagram we have

$$q_{I_0}^* p_i^* p_{i*}(\mathbf{c}_{G/B}(u)) = p_{I_0}^*(p_i \circ q_J)^* p_{i*}(\mathbf{c}_{G/B}(u)) = p_{I_0}^* p_{I_0*} q_{I_0}^*(\mathbf{c}_{G/B}(u))$$

By Lemma 12.2 and by Prop. 10.10.(4) we obtain

$$= p_{I_0}^* p_{I_0*}(\mathbf{c}_{I_0}(u)) = \mathbf{c}_{I_0}(C_{\alpha_i}(u)) = q_{I_0}^*(\mathbf{c}_{G/B}(C_{\alpha_i}(u)))$$

and the theorem is proved. \square

Part 3. Algebraic and geometric comparison. Applications.

13. COMPARISON RESULTS

In the present section we explain why the ring $\mathcal{H}(M)_F$ is naturally isomorphic to the ring $h^*(G/B)$ (see 13.12) for most weakly birationally invariant theories.

13.1. Let I be a sequence of simple reflections. Recall that $w(I) = s_{i_1} \cdots s_{i_l}$ is the corresponding product of simple reflections. Let

$$\zeta_I = (q_I)_*(1) \in h^*(G/B)$$

denote the push-forward of the fundamental class of the Bott-Samelson variety corresponding to I .

13.2. Assumption. For each element $w \in W$, let I_w be a chosen reduced decomposition of w . The elements ζ_{I_w} , where w runs through all elements of the Weyl group W , form an R -basis of the cohomology $h^*(G/B)$.

13.3. Lemma. *Assumption 13.2 holds for*

- (1) *any oriented cohomology theory $h_1^* = h_0^* \otimes_{R_0} R_1$ obtained from another one h_0^* for which the assumption already holds;*
- (2) *Chow groups and Chow groups mod n (in arbitrary characteristic);*
- (3) *algebraic cobordism, and therefore all oriented cohomology theories over a base field of characteristic zero;*

(4) *K-theory (and K-theory mod n) in arbitrary characteristic.*

Proof. Point (1) is obvious since a free module stays free by base change. For Chow groups and K-theory, this is classical and for example proved in [8]. For algebraic cobordism, since G/B is a cellular space and classes of Schubert varieties form a \mathbb{Z} -basis of $\mathrm{CH}(G/B)$, the lemma follows from [21, Cor. 2.9]. \square

We consider the characteristic map $\mathbf{c}_{G/B}: R[[M]]_F \rightarrow \mathfrak{h}^*(G/B)$.

13.4. Lemma. *Let $\pi: G/B \rightarrow \mathrm{pt}$ be the structure map. For any sequence I and $u \in R[[M]]_F$ we have*

$$\pi_*(\zeta_I \cdot \mathbf{c}_{G/B}(u)) = \epsilon C_I(u).$$

Proof. The projection formula and Lemma 11.5 imply that we have

$$\pi_*(\zeta_I \cdot \mathbf{c}_{G/B}(u)) = \pi_*(q_{I*}(1) \cdot \mathbf{c}_{G/B}(u)) = \pi_* q_{I*} q_I^*(\mathbf{c}_{G/B}(u)) = \pi_{I*}(\mathbf{c}_I(u)) = \epsilon C_I(u).$$

\square

Recall from Definition 6.1 that we also have an algebraic characteristic map $\mathbf{c}: R[[M]]_F \rightarrow \mathcal{H}(M)_F$.

13.5. Lemma. *Assume the torsion index \mathfrak{t} is invertible in R and that Assumption 13.2 is satisfied. Then, there is a unique morphism $\theta: \mathcal{H}(M)_F \rightarrow \mathfrak{h}^*(G/B)$ of R -algebras such that $\mathbf{c}_{G/B} = \theta \circ \mathbf{c}$.*

Proof. We fix a reduced decomposition I_w for each $w \in W$. By 6.4, the map \mathbf{c} is surjective. We can therefore find a pre-image u_w of each element $z_{I_w}^C$ of Theorem 6.3. By the form of the characteristic map given there, these elements satisfy $\epsilon C_{I_w}(u_w) = \delta_{w,v}$. Let us set $\tau_{I_w} = \mathbf{c}_{G/B}(u_w)$. By the previous lemma, $(\tau_{I_w})_{w \in W}$ is a basis of $\mathfrak{h}^*(G/B)$ that is dual basis to the basis $(\zeta_{I_w})_{w \in W}$ with respect to the bilinear form $(\zeta, \tau) \mapsto \pi_*(\zeta \cdot \tau)$, and this form is therefore nondegenerate. We now define θ as the R -linear morphism sending $z_{I_w}^C$ to τ_{I_w} . By the formula $\pi_*(\zeta_{I_w} \cdot \mathbf{c}_{G/B}(u)) = \epsilon C_{I_w}(u)$, we must have

$$(8) \quad \mathbf{c}_{G/B}(u) = \sum_{w \in W} \epsilon C_{I_w}(u) \tau_{I_w}$$

and the isomorphism θ therefore satisfies $\mathbf{c}_{G/B} = \theta \circ \mathbf{c}$ since the $z_{I_w}^C$ satisfy the same formula with \mathbf{c} . The uniqueness of θ is immediate by surjectivity of \mathbf{c} . \square

13.6. Corollary. *When \mathfrak{t} is invertible in R and Assumption 13.2 holds, the characteristic map $\mathbf{c}_{G/B}$ is surjective.*

Proof. Use Theorem 6.4 and the previous lemma. \square

13.7. Corollary. *When \mathfrak{t} is invertible in R which has no 2-torsion and Assumption 13.2 holds, the kernel of the characteristic map $\mathbf{c}_{G/B}$ is the ideal of $R[[M]]_F$ generated by elements in \mathcal{I} fixed by W .*

Proof. Use Theorem 6.9 and the previous lemma. \square

Let $u_0 \in \mathcal{I}^N$ be chosen as in section 5.

13.8. Lemma. *Let I_w be a reduced decomposition of an element w . Then*

$$\zeta_{I_w} \cdot \mathbf{c}_{G/B}(u_0) = \begin{cases} 0 & \text{if } w \neq w_0 \\ \mathbf{c}_{G/B}(u_0) & \text{if } w = w_0 \end{cases}$$

Proof. We have $\zeta_{I_w} \cdot \mathbf{c}_{G/B}(u_0) = q_{I_w*}(\mathbf{c}_{I_w}(u_0))$ by the projection formula and the fact that $q_{I_w}^* \mathbf{c}_{G/B} = \mathbf{c}_{I_w}$. By formula (6), $\mathbf{c}_{I_w}(u_0) = 0$ if $l(I_w) < N$ i.e. if $w \neq w_0$. This proves the first part of the formula. The other case will follow if we show that $\zeta_{I_{w_0}} = 1 - \sum_{w \neq w_0} r_w \zeta_{I_w}$ for some coefficients r_w in R . For this, it suffices to show that the coefficient r_{w_0} in front of $\zeta_{I_{w_0}}$ in the decomposition of 1 on the basis of the ζ_{I_w} is 1. We have the commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{i_{U'}} & X_{I_0}/B \xleftarrow{\sigma} X_{I'}/B \\ \parallel & \square & \downarrow q_{I_0} \swarrow q_{I'} \\ U & \xrightarrow{i_U} & G/B \end{array}$$

where I_0 is any reduced decomposition of w_0 , I' is I_0 with its last entry removed, σ is the section of p_{I_0} , U is the big open cell in G/B and U' is its pre-image in X_{I_0}/B . By [8, §3.10 and 3.11], U' is indeed isomorphic to U and it is included in the complement of $\sigma X_{I'}/B$. Thus, $i_{U'}^* \sigma_* = 0$, using that $X_{I_0}/B \rightarrow X_{I'}/B$ is a projective bundle. By base change on the cartesian diagram, we obtain $i_U^* q_{I'}^* = 0$ and $i_U^* q_{I_0}^*(1) = 1$. Since any I_w can be completed to a decomposition of w_0 , this proves that $i_U^*(\zeta_{I_w}) = 0$ if $w \neq w_0$. Pulling back $1 = \sum_w r_w \zeta_{I_w}$ by i_U , we get $1 = r_{w_0}$ (by homotopy invariance, $\mathbf{h}^*(U) \simeq R$). \square

13.9. Lemma. *We have*

$$\mathfrak{t} \zeta_\emptyset = \mathbf{c}_{G/B}(u_0) \quad \text{and} \quad \mathfrak{t} \zeta_I = \mathbf{c}_{G/B}(C_{I^{\text{rev}}}(u_0)).$$

Proof. Let $I_{w_0} = (i_1, \dots, i_N)$. With the notation of Theorem 11.4, we have

$$\begin{aligned} \mathfrak{t} \zeta_\emptyset &= \mathfrak{t} q_{\emptyset*}(1) = \mathfrak{t} q_{I_{w_0}*} \sigma_{I_{w_0}*} \cdots \sigma_{(i_1)*}(1) = q_{I_{w_0}*}(\mathfrak{t} \xi_{[1,N]}) = q_{I_{w_0}*}(\mathbf{c}_{I_{w_0}}(u_0)) \\ &= q_{I_{w_0}*}(1) \mathbf{c}_{G/B}(u_0) = \mathbf{c}_{G/B}(u_0) \end{aligned}$$

where the two last equalities follow from the projection formula together with $q_{I_w}^* \mathbf{c}_{G/B} = \mathbf{c}_{I_w}$ and then Lemma 13.8. For the second formula, which coincides with the first one when I is empty, we use induction on the length of I . If $I = (i_1, \dots, i_l)$ and $J = (i_1, \dots, i_{l-1})$, then by base change via diagram (7) we obtain

$$\mathfrak{t} \cdot \zeta_I = \mathfrak{t} \cdot q_{I*}(1) = \mathfrak{t} \cdot q_{I*} p_I^*(1) = \mathfrak{t} \cdot p_{i_l}^*(p_{i_l} \circ q_J)_*(1) = p_{i_l}^* p_{i_l*}(\mathfrak{t} \cdot \zeta_J),$$

By the induction step and Theorem 12.4 we obtain

$$= p_{i_l}^* p_{i_l*}(\mathbf{c}_{G/B}(C_{J^{\text{rev}}}(u_0))) = \mathbf{c}_{G/B}(C_{\alpha_{i_l}} \circ C_{J^{\text{rev}}}(u_0)) = \mathbf{c}_{G/B}(C_{I^{\text{rev}}}(u_0))$$

and the proof is finished. \square

13.10. Lemma. *Under the assumptions of Proposition 13.5, we have $\zeta_\emptyset = \tau_{I_{w_0}}$. In particular, $\tau_{I_{w_0}}$ does not depend on the choices of the reduced decompositions I_w .*

Proof. By definition of the dual basis $(\tau_{I_w})_{w \in W}$, it suffices to show that we have $\pi_*(\zeta_{I_w} \cdot \zeta_\emptyset) = \delta_{w_0, w}$. By lemmas 13.9 and 13.4, we have

$$\pi_*(\zeta_{I_w} \cdot \zeta_\emptyset) = \pi_*(\zeta_{I_w} \cdot \mathfrak{t}^{-1} \mathbf{c}_{G/B}(u_0)) = \mathfrak{t}^{-1} \epsilon_{C_{I_w}}(u_0) = \delta_{w, w_0}. \quad \square$$

Let \tilde{A}_i denote the operator $p_i^*(p_i)_*$ on $\mathbf{h}^*(G/B)$ and let \tilde{A}_I be as usual for a sequence I .

13.11. Lemma. *The isomorphism θ of Lemma 13.5 satisfies $\theta A_I = \tilde{A}_I \theta$ and therefore sends the $A_{I_w}(z_0)$ to the $\tilde{A}_{I_w}(\zeta_\emptyset)$.*

Proof. It follows from the surjectivity of the characteristic map, the fact that $\theta \circ \mathbf{c} = \mathbf{c}_{G/B}$, Theorem 12.4 and Proposition 7.8 point (1). \square

We can now show that the isomorphism θ of Lemma 13.5 descends to the case where \mathfrak{t} is regular but not necessarily invertible.

13.12. Theorem. *Let \mathfrak{h}^* be a weakly birationally invariant theory satisfying Assumption 13.2 and such that \mathfrak{t} is regular in $R = \mathfrak{h}^*(\text{pt})$. Then there is a unique isomorphism of R -algebras $\theta : \mathcal{H}(M)_F \rightarrow \mathfrak{h}^*(G/B)$ such that the characteristic map $\mathfrak{c}_{G/B}$ is given by $\theta \circ \mathfrak{c}$. It satisfies $\theta \circ A_I = \hat{A}_I \circ \theta$.*

Proof. We know from 7.9 that the $A_{I_w}(z_0)$ are a basis of $\mathcal{H}(M)_F$ and from Assumption 13.2 that the ζ_{I_w} are a basis of $\mathfrak{h}^*(G/B)$. We define θ as the isomorphism sending $A_{I_w}(z_0)$ to ζ_{I_w} . By Lemma 13.11, this definition coincides with the one of Lemma 13.5 when \mathfrak{t} is invertible in R . We therefore have a commutative diagram

$$\begin{array}{ccc} \mathcal{H}(M)_F \otimes_R R[\mathfrak{t}^{-1}] & \xrightarrow{\theta \otimes id} & \mathfrak{h}^*(G/B) \otimes_R R[\mathfrak{t}^{-1}] \\ \uparrow & & \uparrow \\ \mathcal{H}(M)_F & \xrightarrow{\theta} & \mathfrak{h}^*(G/B) \end{array}$$

where $\theta \otimes id$ is the θ of Lemma 13.5 and the vertical maps are injective since $\mathcal{H}(M)_F$ and $\mathfrak{h}^*(G/B)$ are free R -modules and R injects in $R[\mathfrak{t}^{-1}]$. Since all other maps are ring morphisms, θ is one too, and the required equalities follow from the same ones in the case where \mathfrak{t} is invertible. The morphism θ is unique since it is unique after inverting \mathfrak{t} . \square

13.13. Remark. Note that the morphism θ is independent of the choices of the I_w used in its construction. This is obvious by surjectivity of the characteristic map after inverting \mathfrak{t} .

This completes the identification of $\mathcal{H}(M)_F$, the algebraic model for the cohomology ring introduced in section 5, with the actual cohomology $\mathfrak{h}^*(G/B)$.

14. FORMULAS FOR PUSH-FORWARDS

In the present section we assume that \mathfrak{t} and 2 are regular in R . Note that given a formal group law F , both $R[[M]]_F$ and $\mathcal{H}(M)_F$ can be defined without any reference to a cohomology theory. The goal of this section is to define an ‘‘algebraic push-forward’’ $\pi_{\mathcal{H}} : \mathcal{H}(M)_F \rightarrow R$ and to prove formulas related to it. Of course, when the formal group law comes from a cohomology theory, we prove that the morphism $\pi_{\mathcal{H}}$ corresponds to its geometric counterpart $\pi_* : \mathfrak{h}(G/B) \rightarrow R$ through the isomorphism $\theta : \mathcal{H}(M)_F \xrightarrow{\sim} \mathfrak{h}^*(G/B)$ from Theorem 13.12. The general idea of the proofs is to use that there is a universal formal group law U over the Lazard ring \mathbb{L} , and therefore that such formulas can be proved in $\mathcal{H}_{\mathbb{L}}(M)_U$, where they hold for geometric reasons, using algebraic cobordism. Then, they hold in any $\mathcal{H}(M)_F$ by specialization (even if it has no geometric origin).

14.1. Recall that since (U, \mathbb{L}) is the universal formal group law, for any given formal group law F over a ring R , there is a unique morphism $f : \mathbb{L} \rightarrow R$ sending U to F . By Lemma 2.6 and Propositions 6.5 and 7.10, there is a commutative diagram

$$\begin{array}{ccc} \mathbb{L}[[M]]_U & \xrightarrow{c^U} & \mathcal{H}_{\mathbb{L}}(M)_U \\ \downarrow f_* & & \downarrow f_{\mathcal{H}} \\ R[[M]]_F & \xrightarrow{c^F} & \mathcal{H}_R(M)_F \end{array}$$

Furthermore, the morphism $f_{\mathcal{H}}$ commutes with the operators A_I and B_I and sends z_0^U to z_0^R by definition. Therefore, it sends $A_I^U(z_0^U)$ to $A_I^F(z_0^R)$ for any sequence I .

Let $(I_w)_{w \in W}$ be a choice of reduced decompositions. Then, by Prop. 7.9, the elements $(A_{I_w}(z_0))_{w \in W}$ form an R -basis of $\mathcal{H}(M)_F$.

14.2. Definition. Let $\pi_{\mathcal{H}} : \mathcal{H}(M)_F \rightarrow R$ be the R -linear morphism defined by $\pi_{\mathcal{H}}(A_{I_w}(z_0)) = \epsilon C_{I_w}(1)$.

14.3. Proposition. *We have*

- (1) *for any morphism $f : R \rightarrow R'$ of rings sending a formal group law F to F' , we have $\pi_{\mathcal{H}}^{F'} = \pi_{\mathcal{H}}^F \circ f_{\mathcal{H}}$.*
- (2) *Assume that \mathfrak{h} is an oriented cohomology theory satisfying the assumptions of Theorem 13.12, in which the isomorphism $\theta : \mathcal{H}(M)_F \xrightarrow{\sim} \mathfrak{h}^*(G/B)$ is defined. Then the morphism $\pi_{\mathcal{H}}$ satisfies $\pi_{\mathcal{H}} \circ \theta = \theta \circ \pi_*$. In other words, the morphism $\pi_{\mathcal{H}}$ is an algebraic replacement for the push-forward π_* .*

Proof. Since $f_{\mathcal{H}}(A_{I_w}^F(z_0^F)) = A_{I_w}^{F'}(z_0^{F'})$, (1) follows from the definition of $\pi_{\mathcal{H}}$. Since the basis of $\mathfrak{h}(G/B)$ formed by the ζ_{I_w} and the basis of $\mathcal{H}(M)_F$ formed by the $A_{I_w}(z_0)$ corresponds to each other through the isomorphism θ , (2) follows from Lemma 13.4. \square

14.4. Corollary. *For any sequence I the morphism $\pi_{\mathcal{H}}$ of the proposition satisfies*

$$\pi_{\mathcal{H}}(A_{I^{\text{rev}}}(z_0)\mathfrak{c}(u)) = \epsilon C_I(u).$$

In particular, it is independent of the choice $(I_w)_{w \in W}$ of reduced decompositions.

Proof. By Proposition 14.3.(2) and by Lemma 13.4, the formula holds for $\mathcal{H}_{\mathbb{L}}(M)_U$. In general, it reduces to the case where \mathfrak{t} is invertible. When $v \in R[[M]]_F^W$, we have $\epsilon C_I(vu) = \epsilon(v)\epsilon C_I(u)$ and $\mathfrak{c}(vu) = \epsilon(v)\mathfrak{c}(u)$ so by decomposing u on the $C_{I_w}(u_0)$, which form an $R[[M]]_F^W$ -basis of $R[[M]]_F$, it suffices to prove the formula for $u = C_{I_w}(u_0)$. For those, it follows by specialization using $f_{\mathcal{H}}$ from $\mathcal{H}_{\mathbb{L}}(M)_U$. \square

14.5. Remark. Note that the elements $\pi_{\mathcal{H}}(A_{I^{\text{rev}}}(z_0))$ are particularly important because they represent the images of desingularized Schubert varieties in the cohomology of the point.

14.6. Proposition. *The morphism $\pi_{\mathcal{H}}$ satisfies*

$$\pi_{\mathcal{H}}(A_{I_w^{\text{rev}}}(z_0) \cdot z_{I_w}^C) = \delta_{v,w}.$$

In other words, the bilinear form $\pi_{\mathcal{H}}(-, -)$ is nondegenerate and the bases $(z_{I_w}^C)_{w \in W}$ and $(A_{I_w^{\text{rev}}}(z_0))_{w \in W}$ are dual to each other.

Proof. The formula can be computed after extending scalars to $R[\mathfrak{t}^{-1}]$. Then, since the characteristic map \mathfrak{c} is surjective, the elements $z_{I_w}^C$ have preimages u_w , such that $\epsilon C_{I_w}(u_w) = \delta_{v,w}$, by the expression of \mathfrak{c} given in Theorem 6.3. The result then follows from formula (14.4). \square

Let I_0 be a reduced decomposition of the longest element w_0 . When \mathfrak{t} is invertible, the element $y_0 = C_{I_0^{\text{rev}}}(u_0)/\mathfrak{t}$ is invertible (by Lemma 5.3), and $\mathfrak{c}(y_0) = A_{I_0^{\text{rev}}}(z_0)$ by Proposition 7.8.

14.7. Lemma. *When \mathfrak{t} is invertible in R , the morphism $\pi_{\mathcal{H}}$ satisfies*

$$\pi_{\mathcal{H}}(\mathfrak{c}(u)) = \epsilon C_{I_0}(uy_0^{-1}).$$

Proof. We have

$$\pi_{\mathcal{H}}(\mathfrak{c}(u)) = \pi_{\mathcal{H}}(\mathfrak{c}(uy_0^{-1})\mathfrak{c}(y_0)) = \pi_{\mathcal{H}}(\mathfrak{c}(uy_0^{-1}A_{I_0^{\text{rev}}}(u_0))) = \epsilon C_{I_0}(uy_0^{-1}). \quad \square$$

14.8. Proposition. *The operators C satisfy the formula*

$$\epsilon C_I(uC_J(u_0)) = \epsilon C_{J^{\text{rev}}}(uC_{I^{\text{rev}}}(u_0))$$

for any pair of sequences I and J . In particular ($u = 1, J = \emptyset$), for any sequence I , we have $\epsilon C_I(u_0) = \epsilon C_{I^{\text{rev}}}(u_0)$.

Proof. The formula can be proved after extension to $R[\mathfrak{t}^{-1}]$. It then follows from the computation of $\pi_{\mathcal{H}}$ in formula (14.4) applied to both ends of

$$A_{I^{\text{rev}}}(z_0)\mathfrak{c}(uC_J(u_0)) = \mathfrak{t}\mathfrak{c}(C_{I^{\text{rev}}}(u_0)uC_J(u_0)) = \mathfrak{c}(uC_{I^{\text{rev}}}(u_0))A_J(z_0). \quad \square$$

14.9. Remark. The formulas given in this section are purely algebraic, but their proofs use geometric properties of cobordism. It would be interesting to have algebraic proofs derived directly from the various formulas in section 3.

15. ALGORITHM FOR MULTIPLYING IN $\mathfrak{h}^*(G/B)$

Fixing a choice of reduced decompositions I_w for all $w \in W$, we define τ_{I_w} as $\theta(z_{I_w}^C)$.

15.1. Proposition. *Under the assumptions of Theorem 13.12, the bilinear form $(b, a) \mapsto \pi_*(b \cdot a)$ is non degenerate on $\mathfrak{h}^*(G/B)$ and for any choice of reduced decompositions I_w , the basis (ζ_{I_w}) and (τ_{I_w}) are dual to each other.*

Proof. The form can be computed after scalar extension to $R[\mathfrak{t}^{-1}]$, in which case it follows from the proof of Lemma 13.5. \square

Now the multiplication algorithm goes as follows: Substituting $u = C_{I_w^{\text{rev}}}(u_0)$ in (8) and using that $\mathfrak{t}\zeta_{I_w} = \mathfrak{c}_{G/B}(C_{I_w^{\text{rev}}}(u_0))$ we obtain the transition matrix from the basis (ζ_{I_w}) to the basis (τ_{I_w}) . Substituting $u = C_{I_w^{\text{rev}}}(u_0)C_{I_w^{\text{rev}}}(u_0)$ in (8) we obtain the decomposition of the product $\zeta_{I_w}\zeta_{I_w'}$ on the basis (τ_{I_w}) , and we rewrite it in terms of the ζ_{I_w} by using the transition matrix. This also explains how to decompose any ζ_I (I not necessarily reduced) on a given basis of ζ_{I_w} . Again, it suffices to substitute $u = C_I(u_0)$ in (8) and use the transition matrix.

15.2. Remark. Note that this algorithm is an improvement over the one in [5] or [12] in the following sense. In both of these articles, it is explained how to decompose a product of two generators ζ_I and ζ_J as a linear combination of other such generators. But starting from a basis (i.e. the ζ_{I_w} for a choice of a reduced decomposition for each $w \in W$), it is not explained how to obtain a linear combination containing only generators ζ_I with I among the I_w and an algorithm for redecomposing any given ζ_I on a chosen basis is not given either. This is crucial to compute multiplication tables.

16. LANDWEBER-NOVIKOV OPERATIONS

In this section we provide an algorithm for computing the Landweber-Novikov operations S^{LN} on $\Omega^*(G/B)$.

16.1. Let us recall briefly the definition of S^{LN} (details can be found in [13, § 4.1.9]). Consider a graded polynomial ring $\mathbb{Z}[\mathfrak{t}] = \mathbb{Z}[t_1, t_2, \dots, t_k, \dots]$ in infinite number of variables; for a multi-index $I = (i_1, i_2, \dots, i_k)$ we set

$$\mathfrak{t}^I = t_1^{i_1} t_2^{i_2} \dots t_k^{i_k}.$$

Let $\lambda_{(\mathfrak{t})}$ denote the formal power series

$$\lambda_{(\mathfrak{t})}(x) = x + \sum_{i=1}^{\infty} t_i x^{i+1}.$$

Consider a twisted theory $\tilde{\Omega}$ of $\Omega[\mathfrak{t}] = \Omega \otimes_{\mathbb{Z}} \mathbb{Z}[\mathfrak{t}]$ (see [14, §4]). By definition $\tilde{\Omega}(X) = \Omega(X)[\mathfrak{t}]$ for any X , its Chern class is given by the formula

$$(9) \quad c_1^{\tilde{\Omega}}(\mathcal{L}) = \lambda_{(\mathfrak{t})}(c_1^{\Omega}(\mathcal{L})),$$

and its formal group law is given by

$$F(x, y) = \lambda_{(\mathfrak{t})}(U(\lambda_{(\mathfrak{t})}^{-1}(x), \lambda_{(\mathfrak{t})}^{-1}(y))).$$

By the universality of Ω there is a natural transformation $\Omega \rightarrow \tilde{\Omega}$ given by

$$a \mapsto \sum_I S_I^{LN}(a)t^I, \quad a \in \Omega(X)$$

where the components S_I^{LN} are called *Landweber-Novikov operations* on $\Omega(X)$.

16.2. By functoriality of $R[[M]]_F$ in R (see 2.6) the map $\mathbb{L} \simeq \Omega(\text{pt}) \rightarrow \tilde{\Omega}(\text{pt}) \simeq \mathbb{L}[\mathfrak{t}]$ induces a homomorphism

$$\mathbb{L}[[M]]_U \rightarrow \mathbb{L}[\mathfrak{t}][[M]]_F,$$

while functoriality with respect to formal group laws induces a homomorphism

$$\begin{aligned} \mathbb{L}[\mathfrak{t}][[M]]_F &\rightarrow \mathbb{L}[\mathfrak{t}][[M]]_{U_{\Omega[\mathfrak{t}]}} \\ x_\mu &\mapsto \lambda_{(\mathfrak{t})}(x_\mu). \end{aligned}$$

By (9) we have the following commutative diagram:

$$\begin{array}{ccccc} \mathbb{L}[[M]]_U & \longrightarrow & \mathbb{L}[\mathfrak{t}][[M]]_F & \longrightarrow & \mathbb{L}[\mathfrak{t}][[M]]_{U_{\Omega[\mathfrak{t}]}} \\ \mathfrak{c}^\Omega \downarrow & & \mathfrak{c}^{\tilde{\Omega}} \downarrow & & \mathfrak{c}^{\Omega[\mathfrak{t}]} \downarrow \\ \Omega(G/B) & \longrightarrow & \tilde{\Omega}(G/B) & \longrightarrow & \Omega(G/B)[\mathfrak{t}]. \end{array}$$

16.3. An action of the Landweber-Novikov operation S_I^{LN} on a basis element ζ_{I_w} can be computed as follows: First, we compute the image of $u = C_{I_w^{\text{rev}}}(u_0)$ under the composition of top horizontal arrows. Second, we extract the coefficient at the monomial \mathfrak{t}^I of this image. Finally, we apply the characteristic map \mathfrak{c}^Ω to that coefficient. The result will give $S_I^{LN}(\zeta_{I_w})$.

Indeed, by definition $S_I^{LN}(\zeta_{I_w})$ is equal to the coefficient at \mathfrak{t}^I of the image of ζ_{I_w} under the composition of bottom horizontal arrows. Since $\mathfrak{t} \zeta_I = \mathfrak{c}_{G/B}(C_{I^{\text{rev}}}(u_0))$ and by commutativity of the diagram we are done.

17. EXAMPLES OF COMPUTATIONS

In the present section we list the multiplication tables for rings $\Omega^*(G/B)$, where G has rank 2. The results are obtained by means of the algorithm described in Section 15 and realized in Macaulay 2 packages [6]. The answers for the other oriented cohomology theories are easily derived by a specialization of the coefficients of the universal formal group law. For instance, the answer for connective K -theory is obtained by specializing the coefficient a_1 to v and all others to zero.

17.1. We use the presentation of the Lazard ring

$$\mathbb{L} = \mathbb{Z}[a_1, a_2, \dots],$$

where the first generators a_i are the following linear combinations of the coefficients a_{ij} of the universal formal group law U :

$$\begin{aligned} a_1 &= a_{11}; \\ a_2 &= a_{12}; \\ a_3 &= a_{22} - a_{13}; \\ a_4 &= a_{14}; \\ a_5 &= -9a_{15} + a_{24} + 2a_{33}. \end{aligned}$$

Note that the a_i (resp. the a_{ij}) are of cohomological degree $-i$ (resp. a_{1-i-j}). For root systems of rank 2, the longest element is of length at most $N = 6$ (in the G_2

case), and we therefore need to compute the universal formal group law up to order 7, which thus only involves a_1, \dots, a_6 . In fact, a_6 does not appear in the formulas: it is not difficult to show that a_N will not appear in the multiplication formulas for a root system with longest element of length N .

17.2. We use the upper case letter ζ_I for the element $\zeta_I = q_{I^*}(1)$. For brevity ζ_I with $I = (i_1, \dots, i_l)$ is denoted just by $\zeta_{i_1 \dots i_l}$, and when I is the empty sequence, ζ_I is denoted by pt . Note that when $l(w) + l(w') \geq N = \dim G/B$ one has

$$\zeta_{I_w} \zeta_{I_{w'}} = \delta_{w, w_0 w'} \text{pt}$$

so we list the remaining cases only.

17.3 (A_2 case).

$$\begin{aligned} \zeta_{121} &= 1 + a_2 \zeta_1; \\ \zeta_{12}^2 &= \zeta_2; \\ \zeta_{21}^2 &= \zeta_1; \\ \zeta_{12} \zeta_{21} &= \zeta_1 + \zeta_2 + a_1 \text{pt}. \end{aligned}$$

This agrees with the computations of Hornbostel-Kiritchenko in [12].

17.4 (B_2 case).

$$\begin{aligned} \zeta_{1212} &= 1 + 2a_2 \zeta_{12} + (a_3 - a_1 a_2) \zeta_2; \\ \zeta_{121}^2 &= \zeta_{21}; \\ \zeta_{212}^2 &= 2\zeta_{12} + a_1 \zeta_2; \\ \zeta_{121} \zeta_{212} &= \zeta_{12} + \zeta_{21} + a_1 \zeta_1 + a_1 \zeta_2 + (2a_2 + a_1^2) \text{pt}; \\ \zeta_{121} \zeta_{12} &= \zeta_1 + \zeta_2 + a_1 \text{pt}; \\ \zeta_{121} \zeta_{21} &= \zeta_1; \\ \zeta_{212} \zeta_{12} &= \zeta_2; \\ \zeta_{212} \zeta_{21} &= 2\zeta_1 + \zeta_2 + 2a_1 \text{pt}. \end{aligned}$$

17.5 (G_2 case).

$$\begin{aligned} \zeta_{121212} &= 1 + 4a_2 \zeta_{1212} + (10a_3 - 10a_1 a_2) \zeta_{212} \\ &\quad - (4a_4 + 9a_1 a_3 + 3a_2^2 - 9a_1^2 a_2) \zeta_{12} \\ &\quad - (54a_5 - 459a_1 a_4 - 1188a_2 a_3 - 108a_1^2 a_3 + 1080a_1 a_2^2 + 108a_1^3 a_2) \zeta_2; \\ \zeta_{12121}^2 &= 3\zeta_{2121} + 3a_1 \zeta_{121} + (13a_2 + 2a_1^2) \zeta_{21} + (2a_3 + 7a_1 a_2 + a_1^3) \zeta_1; \\ \zeta_{21212}^2 &= \zeta_{1212} + 5a_2 \zeta_{12} + (6a_3 - 5a_1 a_2) \zeta_2; \\ \zeta_{12121} \zeta_{21212} &= \zeta_{1212} + \zeta_{2121} + a_1 \zeta_{121} + a_1 \zeta_{212} + (8a_2 + a_1^2) \zeta_{12} + (8a_2 + a_1^2) \zeta_{21} \\ &\quad + (4a_3 + 8a_1 a_2 + a_1^3) \zeta_1 + (10a_3 + 6a_1 a_2 + a_1^3) \zeta_2 \\ &\quad + (-4a_4 + a_1 a_3 + 13a_2^2 + 15a_1^2 a_2 + a_1^4) \text{pt}; \\ \zeta_{12121} \zeta_{1212} &= \zeta_{121} + 3\zeta_{212} + 4a_1 \zeta_{12} + 3a_1 \zeta_{21} + (8a_2 + 4a_1^2) \zeta_1 + (13a_2 + 5a_1^2) \zeta_2 \\ &\quad + (a_3 + 16a_1 a_2 + 5a_1^3) \text{pt}; \\ \zeta_{12121} \zeta_{2121} &= 2\zeta_{121} + 2a_1 \zeta_{21} + (4a_2 + a_1^2) \zeta_1; \\ \zeta_{21212} \zeta_{1212} &= 2\zeta_{212} + a_1 \zeta_{12} + 4a_2 \zeta_2; \\ \zeta_{21212} \zeta_{2121} &= \zeta_{121} + \zeta_{212} + a_1 \zeta_{12} + a_1 \zeta_{21} + (5a_2 + a_1^2) \zeta_1 + (8a_2 + a_1^2) \zeta_2 \\ &\quad + (3a_3 + 6a_1 a_2 + a_1^3) \text{pt}; \\ \zeta_{12121} \zeta_{121} &= 3\zeta_{21} + 2a_1 \zeta_1; \end{aligned}$$

$$\begin{aligned}
\zeta_{12121}\zeta_{212} &= 2\zeta_{12} + \zeta_{21} + 2a_1\zeta_1 + 3a_1\zeta_2 + (4a_2 + 3a_1^2)\text{pt}; \\
\zeta_{21212}\zeta_{121} &= \zeta_{12} + 2\zeta_{21} + 2a_1\zeta_1 + 2a_1\zeta_2 + (4a_2 + 2a_1^2)\text{pt}; \\
\zeta_{21212}\zeta_{212} &= \zeta_{12}; \\
\zeta_{12121}^2 &= 2\zeta_{12} + a_1\zeta_2; \\
\zeta_{2121}^2 &= 2\zeta_{21} + a_1\zeta_1; \\
\zeta_{1212}\zeta_{2121} &= 2\zeta_{12} + 2\zeta_{21} + 3a_1\zeta_1 + 4a_1\zeta_2 + (4a_2 + 4a_1^2)\text{pt}; \\
\zeta_{12121}\zeta_{12} &= \zeta_1 + 3\zeta_2 + 3a_1\text{pt}; \\
\zeta_{12121}\zeta_{21} &= \zeta_1; \\
\zeta_{21212}\zeta_{12} &= \zeta_2; \\
\zeta_{21212}\zeta_{21} &= \zeta_1 + \zeta_2 + a_1\text{pt}; \\
\zeta_{1212}\zeta_{121} &= 2\zeta_1 + 3\zeta_2 + 4a_1\text{pt}; \\
\zeta_{1212}\zeta_{212} &= \zeta_2; \\
\zeta_{2121}\zeta_{121} &= \zeta_1; \\
\zeta_{2121}\zeta_{212} &= \zeta_1 + 2\zeta_2 + 2a_1\text{pt}.
\end{aligned}$$

REFERENCES

- [1] R. Bott and H. Samelson. Applications of the theory of Morse to symmetric spaces. *Amer. J. Math.*, 80:964–1029, 1958.
- [2] N. Bourbaki. *Éléments de mathématique. Algèbre. Chapitres 1 à 3*. Hermann, Paris, 1970.
- [3] N. Bourbaki. *Éléments de mathématique. Algèbre commutative. Chapitres 1 à 3*. Masson, Paris, 1985.
- [4] P. Bressler and S. Evens. The Schubert calculus, braid relations, and generalized cohomology. *Trans. Amer. Math. Soc.*, 317(2):799–811, 1990.
- [5] P. Bressler and S. Evens. Schubert calculus in complex cobordism. *Trans. Amer. Math. Soc.*, 331(2):799–813, 1992.
- [6] B. Calmès and V. Petrov. CohomologyOfBorelVarieties, a Macaulay 2 package. <http://www.math.uni-bielefeld.de/~bcalmes/M2packages/cohbovar.html>, 2009.
- [7] M. Demazure. Invariants symétriques entiers des groupes de Weyl et torsion. *Invent. Math.*, 21:287–301, 1973.
- [8] M. Demazure. Désingularisation des variétés de Schubert généralisées. *Ann. Sci. École Norm. Sup. (4)*, 7:53–88, 1974. Collection of articles dedicated to Henri Cartan on the occasion of his 70th birthday, I.
- [9] W. Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3*. Springer-Verlag, Berlin, second edition, 1998.
- [10] D. R. Grayson and M. E. Stillman. Macaulay 2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [11] R. Hartshorne. *Algebraic geometry*, volume 52 of *GTM*. Springer-Verlag, New York-Heidelberg, 1977.
- [12] J. Hornbostel and V. Kiritchenko. Schubert calculus for algebraic cobordism. arXiv:0903.3936v1, 2009.
- [13] M. Levine and F. Morel. *Algebraic cobordism*. Springer Monographs in Mathematics. Springer, Berlin, 2007.
- [14] A. Merkurjev. *Algebraic oriented cohomology theories*, volume 300 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, algebraic number theory and algebraic geometry edition, 2002.
- [15] I. Panin. Oriented cohomology theories of algebraic varieties II. *Homology, Homotopy and Applications*, to appear.
- [16] I. Panin. Oriented cohomology theories of algebraic varieties. *K-Theory*, 30(3):265–314, 2003.
- [17] T. Springer. *Schubert varieties and generalizations*, volume 514 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.* Kluwer Acad. Publ., Dordrecht, representation theories and algebraic geometry edition, 1998.
- [18] B. Totaro. The torsion index of E_8 and other groups. *Duke Math. J.*, 129(2):219–248, 2005.
- [19] B. Totaro. The torsion index of the spin groups. *Duke Math. J.*, 129(2):249–290, 2005.
- [20] A. Vishik. Symmetric operations in algebraic cobordism. *Adv. Math.*, 213(2):489–552, 2007.
- [21] A. Vishik and N. Yagita. Algebraic cobordisms of a pfister quadric. *J. Lond. Math. Soc. (2)*, 76(3):586–604, 2007.

UNIVERSITÉ D'ARTOIS, LABORATOIRE DE MATHÉMATIQUES DE LENS, FRANCE
E-mail address: `baptiste.calmes@univ-artois.fr`

MAX-PLANK-INSTITUT FÜR MATHEMATIK, BONN, GERMANY
E-mail address: `victorapetrov@googlemail.com`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, CANADA
E-mail address: `kirill@uottawa.ca`