

# ON THE ALGEBRAIC $K$ -THEORY OF SOME HOMOGENEOUS VARIETIES.

ALEXEY ANANYEVSKIY

ABSTRACT. In this paper the  $K$ -theory of the inner twisted forms of the homogeneous varieties  $G/H$  with the connected reductive algebraic groups  $H \subset G$  of the same rank is computed. We provide an explicit isomorphism with the  $K$ -theory of certain central simple algebras associated to the considered variety, as a consequence one has that  $K_0(G/H)$  is a free abelian group of rank  $[W(G) : W(H)]$ . The result is used for computing the  $K$ -theory of some affine homogeneous varieties including the octonionic projective plane and the quaternionic projective spaces.

## 1. INTRODUCTION.

It is known that  $K$ -theory of homogeneous projective varieties could be expressed by means of  $K$ -theory of central simple algebras. The most fundamental result concerns the case of projective space and states that there is an isomorphism

$$K_*(k)[t]/t^{n+1} \xrightarrow{\sim} K_*(\mathbb{P}^n).$$

The case of Severi-Brauer varieties was treated by Quillen [Q, §8], and one has

$$\bigoplus_{i=0}^n K_*(A^{\otimes i}) \xrightarrow{\sim} K_*(\mathbb{P}_\gamma^n),$$

where  $A$  is a central simple algebra defined by the cocycle  $\gamma$ . Swan [Sw] computed the  $K$ -theory of a smooth projective quadric and showed that there is an analogous isomorphism involving some Clifford algebras. It was shown by Panin [P1] that one has an analogous isomorphism for every homogeneous projective variety and one can express its  $K$ -theory in terms of the  $K$ -theory of the certain separable algebras. In the present paper we provide a unified approach to the  $K$ -theory of the homogeneous varieties and compute it for the inner forms of  $G/H$  with the connected reductive  $H \subset G$  of the same rank. The main result is the theorem 4 which claims that there is an isomorphism

$$K_*((G/H)_\gamma) \xrightarrow{\sim} \bigoplus_{i=1}^r K_*(A(\lambda_i)_\gamma),$$

with  $r = [W(G) : W(H)]$  and the separable algebras  $A(\lambda_i)_\gamma$  associated to  $(G/H)_\gamma$  in some canonical way. In the last section of the present paper it is shown that the known results concerning the projective varieties could be derived from this theorem, although it deals with the affine varieties.

An essential role in our computations plays the well-known equivalence between the category of the equivariant vector bundles over the homogeneous

variety and the category of the finite dimensional representations of the stabilizer of a rational point. Another important ingredient is the spectral sequence constructed by Merkurjev [M] that allows to pass from the equivariant  $K$ -theory to the ordinary one. It turns out that when the groups have the same rank the spectral sequence degenerates and provides the very explicit answer. In order to show that the sequence degenerates we use the theorem proved by Steinberg [St] which states that in our case the representation ring  $R(H)$  is a free module over  $R(G)$ . We give a new proof of the last theorem which provides us some good basis consisting of the irreducible representations such that we can handle it in the twisted case.

Note that there is a decent classification of the connected reductive subgroups containing the maximal torus [BT, § 3]. They correspond to the quasi-closed (for  $\text{char } k = 0$  one can say closed) symmetric subsets in the root system of the group  $G$ , so one can explicitly write down the varieties covered by theorem 4. For example we can compute by hand the  $K$ -theory for the variety  $G(E_6)/G(A_2 + A_2 + A_2)$ , the inclusion provided by  $3A_2 \subset E_6$ . The  $K_0$  in this case is a free abelian group of rank 240.

In the article everything is settled over the field  $k$  of an arbitrary characteristic. Algebraic groups are supposed to be the smooth algebraic varieties over the field  $k$ . The text is organized in the following way. In the second section we recall some well-known facts on the representation theory of the reductive groups, including the combinatorics concerning roots, weights and the Weyl group.

In the next section we introduce some useful combinatorics arising from the reductive subgroup of maximal rank. We define a linear order on the dominant weights and prove key lemmas providing the technical tool for the new approach to the Steinberg theorem.

In section 4 we show that with the given order one can choose some set resembling the Gröbner basis and could carry out the division relative to the chosen elements. Using the above idea we construct the basis for the representation ring in theorem 2 and show that there is a natural freedom in the choice of basis. The introduced division algorithm provides an explicit method for calculation of the multiplicative structure on the obtained free module.

The fifth section contains some examples, from the vivid two-dimensional case involving  $G_2$  to the non-obvious series of  $C_n$  root systems.

In section 6 we recall the basic notions from the equivariant  $K$ -theory and present the spectral sequence constructed by Merkurjev. The following section deals with the split case of the homogeneous varieties, the degeneration of the spectral sequence is demonstrated and the isomorphism for  $K$ -theory is constructed.

Section 8 deals with the twisted forms, separable algebras are introduced and the main result is proved by means of the splitting principle ??.

In the last section we use the developed technique towards concrete examples. First of all the relations with the known results are presented and  $K$ -theory for the twisted flags is computed. Then we turn to the case of characteristic zero and show that  $K$ -theory for any homogeneous variety with the

stabilizer connected and having the maximal rank could be computed without assumption about reductiveness. Also some affine homogeneous examples are considered, including the octonionic projective plane and the quaternionic projective spaces.

I want to express my sincere gratitude to I.Panin for numerous discussions and useful suggestions concerning the subject of this paper.

## 2. REPRESENTATIONS OF REDUCTIVE ALGEBRAIC GROUPS.

In this section we fix the notations and recall some well-known facts concerning the representation theory of split reductive algebraic groups. A comprehensive survey of this theme could be found in [Ja], the semisimple case is covered also in [H2].

Let  $G$  be a connected split reductive algebraic group and let  $T \subset G$  be a split maximal torus of  $G$ . Let  $W(G, T) = N_G(T)/Z_G(T)$  be the Weyl group of  $G$ . Since all split maximal tori are conjugate,  $W(G, T)$  does not depend on the choice of torus  $T$  so we will as usual denote it by  $W(G)$ . Let

$$X^*(T) = \text{Hom}(T, G_m) \cong \mathbb{Z}^{rk(T)}, \quad \text{Ch} = \text{Hom}(Z(G), G_m)$$

be the character groups of torus  $T$  and center  $Z(G)$  respectively. Recall that the Weyl group  $W(G)$  obviously acts on  $X^*(T)$  and that there is a natural Weyl-equivariant  $\text{Ch}$  grading on  $X^*(T)$ .

Let  $\text{Rep}_k(G)$  be the category of finite dimensional  $k$ -rational representations of  $G$  and let  $R(G) = K_0(\text{Rep}_k(G))$  be the representation ring of  $G$ . Recall that as an additive group the ring  $R(G)$  is a free abelian group generated by the isomorphism classes of irreducible representations. The following result is well-known (for example, see [Ja, Cor. 2.7]).

**Theorem 1.** *Let  $G$  be a connected split reductive algebraic group and let  $T \subset G$  be a split maximal torus of  $G$ . Then there is a ring isomorphism  $R(T) \cong \mathbb{Z}[X^*(T)]$  where the last one denotes the group ring. Moreover, the restriction of representations induces  $R(G) \cong \mathbb{Z}[X^*(T)]^{W(G)}$ .*

We need some more combinatorial data on the connection between representations and characters group ring.

There is a root system  $\Phi$  in  $X^*(T)$ , so there is a bilinear form on

$$V = X^*(T) \otimes_{\mathbb{Z}} \mathbb{R} \cong \mathbb{R}^{rk(G)}$$

such that the Weyl group is generated by the reflections  $\{w_\alpha, \alpha \in \Phi\}$ . The hyperplanes  $H_\alpha$  orthogonal to the roots  $\alpha \in \Phi$  divide  $V$  into chambers which are the fundamental domains for the Weyl group action. The hyperplanes adjacent to the chamber are called walls of this chamber. Fix a set of simple roots  $\Pi \subset \Phi$  and denote by

$$\mathcal{C}(G) = \{v \in V \mid (v, \alpha) \geq 0, \alpha \in \Pi\}$$

the fundamental Weyl chamber. The walls of the fundamental Weyl chamber  $\mathcal{C}(G)$  coincide with the hyperplanes orthogonal to the simple roots. Let

$$\Lambda_G^+ = \mathcal{C}(G) \cap X^*(T)$$

be the cone of dominant weights. Note that in the semisimple case group is simply connected iff  $\Lambda_G^+ \cong \mathbb{N}_0^+$ .

Let  $\lambda \in \Lambda_G^+$  be a dominant weight. Theorem 1 states that there is a bijection between such weights and irreducible  $G$ -modules, so we will denote by  $V_G(\lambda)$  the corresponding  $G$ -module.

At last, recall that there is a partial order on  $X^*(T)$  which is defined by the set of simple roots  $\Pi$ :  $\mu \preceq_{\Pi} \lambda$  if and only if  $\lambda - \mu$  is the sum of positive roots. The interaction between this ordering and Weyl action is stated in the next lemma [H1, Lemma 13.2A].

**Lemma 1.** *Let  $\lambda \in \Lambda_G^+$ ,  $w \in W(G)$  then  $w(\lambda) \preceq_{\Pi} \lambda$ .*

### 3. SUBGROUP COMBINATORICS.

In this section we introduce the necessary combinatorics that we need in order to prove theorem 2. The main goal is to order dominant weights of the subgroup and show that there are several weights with good properties relative to the order.

Let  $G$  be a connected split semisimple simply connected group of rank  $r$ , let  $T \subset G$  be a split maximal torus of  $G$  and let  $T \subset H \subset G$  be a connected split reductive subgroup of maximal rank. Evidently, in this setting there is an inclusion of the Weyl groups  $W(H) \subset W(G)$ . Hence we have the corresponding combinatorial data introduced in the previous section: lattice  $X^*(T) \subset V$  in the euclidean space, root system  $\Phi \subset X^*(T)$  and actions of the Weyl groups  $W(H) \subset W(G)$  on the  $V$ . The following lemma shows that we could choose the compatible fundamental Weyl chambers and the corresponding cones of dominant weights

$$\begin{array}{ccc} \mathcal{C}(G) & \hookrightarrow & \mathcal{C}(H) \\ \uparrow & & \uparrow \\ \Lambda_G^+ & \hookrightarrow & \Lambda_H^+ \end{array}$$

Let  $k = [W(G) : W(H)]$  be the Weyl group index.

**Lemma 2.** *Any Weyl chamber of group  $H$  is the union of  $k$  Weyl chambers of group  $G$ .*

*Proof.* It is clear that any wall for  $W(H)$  action is a wall for  $W(G)$  action, so in order to get chambers of group  $G$  we need to subdivide  $H$  chambers. The number of  $G$  subchambers is independent on  $H$  chamber. Since the number of chambers coincides with the order of Weyl group, the number of subchambers equals to the index  $k$ .  $\square$

So we choose some chambers  $\mathcal{C}(G) \subset \mathcal{C}(H)$  and elements

$$e = w_1, w_2, \dots, w_k \in W(G),$$

such that

$$\mathcal{C}(H) = \bigcup_{1 \leq i \leq k} w_i \mathcal{C}(G), \quad \Lambda_H^+ = \bigcup_{1 \leq i \leq k} w_i \Lambda_G^+.$$

Let  $\omega_1, \dots, \omega_r$  be the fundamental weights corresponding to  $\mathcal{C}(G)$  and let  $\Pi, \Pi'$  be the sets of simple roots for  $G, H$  respectively. By  $\Phi^+$  denote the set of positive roots of  $G$  relative to  $\Pi$ .

**Definition 1.** Let  $\mu \in X^*(T)$  be some weight. Define

$$H(\mu) = \{H_\alpha, \alpha \in \Phi^+ | \exists i : (\mu, \alpha) \cdot (\omega_i, \alpha) < 0\} = \{H_\alpha, \alpha \in \Phi^+ | (\mu, \alpha) < 0\}$$

to be the set of walls which separate  $\mu$  from  $\mathcal{C}(G)$ . Let  $H(w\mathcal{C}(G)) = H(\mu)$  for some interior weight  $\mu \in w\mathcal{C}(G)^\circ$ .

*Remark 1.* The set  $H(\mu)$  somehow measures the spherical distance from  $\mu$  to  $\mathcal{C}(G)$ , the furthest weights are separated by the most hyperplanes. Also note that  $\#H(w\mathcal{C}(G)) = l(w)$ , the usual length of an element of Weyl group, which is defined to be the number of simple reflections in the shortest word representing  $w$ .

**Lemma 3.** Let  $\mu \in \Lambda_H^+$  be some dominant weight. Then there exists such  $i$  that  $\mu \in w_i\Lambda_G^+$  and  $H(w_i\mathcal{C}(G)) = H(\mu)$ .

*Proof.* If  $\mu$  belongs to the interior of some chamber we of course should take that chamber. Otherwise we can choose an arbitrary  $\nu \in \mathcal{C}(G)^\circ$  and draw a segment connecting the points corresponding to  $\mu$  and  $\nu$ . Since  $\nu$  is interior for  $\mathcal{C}(G)$  this segment does not belong to hyperplanes  $H_\alpha$  and we should take the chamber  $w_i\mathcal{C}(G)$  which interior it crosses first, starting from the  $\mu$ . There are no hyperplanes separating the chosen chamber from  $\mu$  so  $H(w_i\mathcal{C}(G)) = H(\mu)$ .  $\square$

**Lemma 4.** Let  $\mu, \lambda \in w_i\Lambda_G^+$  for some  $i$  and  $w \in W(G)$ . Suppose that there exists a hyperplane  $H_\alpha$  such that  $(\lambda, \alpha) \cdot (w\mu, \alpha) < 0$ . Then  $(w\mu, \lambda) < (\mu, \lambda)$ .

*Proof.* First of all we multiply weights by  $w_i^{-1}$  and consider

$$\mu' = w_i^{-1}\mu, \lambda' = w_i^{-1}\lambda, w' = w_i^{-1}ww_i, \alpha' = w_i^{-1}\alpha.$$

It follows that  $\mu', \lambda' \in \Lambda_G^+$  and

$$(\lambda', \alpha') \cdot (w'\mu', \alpha') = (w_i^{-1}\lambda, w_i^{-1}\alpha) \cdot (w_i^{-1}w\mu, w_i^{-1}\alpha) = (\lambda, \alpha) \cdot (w\mu, \alpha),$$

and by the same vein

$$(w'\mu', \lambda') = (w\mu, \lambda), (\mu', \lambda') = (\mu, \lambda).$$

So from now on we suppose  $w_i = e$ .

Note that by lemma 1  $\mu - w\mu$  equals to the sum of positive roots and  $\lambda$  is the sum of fundamental weights with nonnegative coefficients, so in general  $(w\mu - \mu, \lambda) \leq 0$ , and we need to show that the difference  $w\mu - \mu$  is not orthogonal to  $\lambda$ . First of all  $w\mu \notin \Lambda_G^+$  (i.e. not equals  $\mu$ ) since there are no hyperplanes crossing  $\mathcal{C}(G)$ .

We can find a sequence  $\alpha_1, \alpha_2, \dots, \alpha_n \in \Phi^+$  such that the following conditions hold, where  $s_i = w_{\alpha_i}w_{\alpha_{i-1}} \dots w_{\alpha_1}$  and  $s_0 = e$ .

- (a)  $w = s_n = w_{\alpha_n}w_{\alpha_{n-1}} \dots w_{\alpha_1}$ .
- (b) For every  $1 \leq i < n$  the hyperplane  $H_{\alpha_i}$  is a wall of  $s_{i-1}\mathcal{C}(G)$  and separates it and  $\mathcal{C}(G)$  from  $s_n\mathcal{C}(G)$

So this presentations divides  $w$  into the sequence of flips, and each flip drives the chamber further from the  $\mathcal{C}(G)$ .

We claim that the roots  $\alpha_i$  should be the roots corresponding to hyperplanes in  $H(w\mathcal{C}(G))$  written in the appropriate order. Indeed, there exists some hyperplane  $H_{\alpha_1} \in H(w\mathcal{C}(G))$  which is the wall of  $\mathcal{C}(G)$ , otherwise  $w\mathcal{C}(G) = \mathcal{C}(G)$

and  $w = e$ , contradicting  $\mu \neq w\mu$ . Note that

$$H(s_1\mathcal{C}(G)) = \{H_{\alpha_1}\} \subset H(w\mathcal{C}(G)),$$

and whenever  $s_1 \neq w$  we can find  $H_{\alpha_2} \in H(w\mathcal{C}(G)) \setminus H(s_1\mathcal{C}(G))$  satisfying the condition (b), i.e. it should be the wall of  $s_1\mathcal{C}(G)$ , the separating part is valid since we look at the separating hyperplanes. Now one has

$$H(s_2\mathcal{C}(G)) = \{H_{\alpha_1}, H_{\alpha_2}\} \subset H(w\mathcal{C}(G)).$$

If  $s_2 \neq w$  we can find  $\alpha_3 \in H(w\mathcal{C}(G)) \setminus H(s_2\mathcal{C}(G))$  and so on.

For the above roots  $\alpha_i$  one has

$$(w\mu - \mu, \lambda) = \left( \sum_{i=1}^n s_i\mu - s_{i-1}\mu, \lambda \right) = \sum_{i=1}^n c_i(\alpha_i, \lambda),$$

where  $c_i = -2 \frac{(s_{i-1}\mu, \alpha_i)}{(\alpha_i, \alpha_i)}$ . From the condition (b) it follows that  $H_{\alpha_i}$  does not separate  $s_{i-1}\mathcal{C}(G)$  from  $\mathcal{C}(G)$ , so  $(s_{i-1}\mu, \alpha_i) \geq 0$ , hence  $c_i$  is nonpositive.

In general  $(\alpha_i, \lambda) \geq 0$  so it is sufficient to show that there exists some  $\alpha_i$  such that  $(\alpha_i, \lambda) \neq 0$  and  $(s_{i-1}\mu, \alpha_i) \neq 0$ . The first condition is equivalent to  $\lambda \notin H_{\alpha_i}$  and the second means that  $s_i\mu \neq s_{i-1}\mu$ . Now suppose that there is no such  $\alpha_i$ , then we can get from  $\mu$  to  $w\mu$  by reflections  $w_{\alpha_i}$  such that  $(s_{i-1}\mu, \alpha_i) = 0$ , i.e. through the hyperplanes which contain  $\lambda$ . Then  $w\mu$  and  $\lambda$  lie in the same chamber and there are no hyperplanes separating them. So by contradiction we can find such  $i$  that the corresponding term  $c_i(\alpha_i, \lambda) < 0$  and this finishes the proof.  $\square$

Now we are ready to introduce a good order on  $\Lambda_H^+$  which uses  $W(G)$  action and hence somehow connects  $W(G)$  orbits with  $H$  weights.

**Definition 2.** Let  $\mu_1, \mu_2 \in \Lambda_H^+$ , we say that  $\mu_1 \preceq' \mu_2$  if and only if one of the following conditions holds:

- (1)  $\mu_1 = \mu_2$
- (2)  $(\mu_1, \mu_1) < (\mu_2, \mu_2)$
- (3)  $(\mu_1, \mu_1) = (\mu_2, \mu_2)$  and  $H(\mu_1) \supsetneq H(\mu_2)$

*Remark 2.* The meaning of the above definition is that the dominant weight is smaller if the vector is shorter or the spherical distance to  $\mathcal{C}(G)$  is greater.

**Lemma 5.**

- (1)  $\preceq'$  defines a partial order on  $\Lambda_H^+$ .
- (2) For any  $\mu \in \Lambda_H^+$  there are only finitely many  $\mu'$  such that  $\mu' \not\preceq' \mu$ .
- (3) Let  $\mu_1, \mu_2 \in \Lambda_H^+$  and  $\mu_1 \prec_{\Gamma'} \mu_2$ . Then  $\mu_1 \prec' \mu_2$ .

*Proof.*

- (1) is checked by hand.
- (2) Follows from the fact that there are finitely many weights  $\mu'$  such that  $(\mu', \mu') \leq (\mu, \mu)$ .
- (3) There exists  $\beta \in X^*(T)$  such that  $\beta$  equals a sum of positive roots and  $\mu_2 = \mu_1 + \beta$ . Then  $(\mu_2, \mu_2) = (\mu_1, \mu_1) + (\beta, \beta) + 2(\mu_1, \beta)$ . The last term is nonnegative since the scalar product of simple root and dominant weight is nonnegative and so is the scalar product of positive root and dominant weight. So  $\mu_1 \preceq' \mu_2$  follows from examining their lengths.

□

**Definition 3.** Let  $\preceq$  be an arbitrary linear extension of order  $\preceq'$ , i.e. such linear order that from  $\mu_1 \preceq' \mu_2$  it follows that  $\mu_1 \preceq \mu_2$ .

*Remark 3.* Part (3) of the previous lemma is valid for  $\preceq$  too and part (2) transforms into the property that there are only finitely many  $\mu'$  such that  $\mu' \preceq \mu$ .

The next lemma introduces basic and in some sense minimal and indecomposable elements  $\lambda_i \in \Lambda_H^+$ , one for each chamber  $w_i\mathcal{C}(G)$ .

**Lemma 6.** For every  $i$  there exists an element  $\lambda_i \in w_i\Lambda_G^+$  such that

- (1)  $H(\lambda_i) = H(w_i\mathcal{C}(G))$ .
- (2) For every  $\mu \in w_i\Lambda_G^+$ ,  $H(\mu) = H(w_i\mathcal{C}(G))$  one has  $\mu - \lambda_i \in w_i\Lambda_G^+$ .
- (3) For every  $\mu \in w_i\Lambda_G^+$  one has  $\lambda_i + w_jw_i^{-1}\mu \preceq \lambda_i + \mu$ .

*Proof.* The set of weights  $\{\mu \in w_i\Lambda_G^+ | H(\mu) \neq H(w_i\mathcal{C}(G))\}$  is just the intersection of  $w_i\Lambda_G^+$  with the union of chamber  $w_i\mathcal{C}(G)$  walls which separate it from the  $\mathcal{C}(G)$ . Indeed, the only chance for the weight to have the lesser number of walls separating it from  $\mathcal{C}(G)$  is to belong to such wall, and every weight lying on this wall has the lesser number of separating hyperplanes.

Now we use the fact that  $G$  is simply connected so  $w_i\Lambda_G^+ \cong \mathbb{N}_0^r$ . The walls of the chamber correspond to the hyperplanes where some coordinate equals 0, so the weights, which have the same  $H(\mu)$  as the chamber, correspond to the points with certain coordinates, say  $1, \dots, l$ , strictly greater than 0. Let  $\lambda_i$  be the element corresponding to the point with first  $l$  coordinates equal 1 and others equal 0. From the above it follows that we get (1) and (2).

First of all note that all weights really lie in the  $\Lambda_H^+$ , so we can try to compare them. Examine their lengths:

$$\begin{aligned} (\lambda_i + w_jw_i^{-1}\mu, \lambda_i + w_jw_i^{-1}\mu) &= (\lambda_i, \lambda_i) + 2(\lambda_i, w_jw_i^{-1}\mu) + (w_jw_i^{-1}\mu, w_jw_i^{-1}\mu) = \\ &= (\lambda_i, \lambda_i) + 2(\lambda_i, w_jw_i^{-1}\mu) + (\mu, \mu) \\ (\lambda_i + \mu, \lambda_i + \mu) &= (\lambda_i, \lambda_i) + 2(\lambda_i, \mu) + (\mu, \mu) \end{aligned}$$

From lemma 1 one has  $(\lambda_i, w_jw_i^{-1}\mu) \leq (\lambda_i, \mu)$  and, consequently,

$$(\lambda_i + w_jw_i^{-1}\mu, \lambda_i + w_jw_i^{-1}\mu) \leq (\lambda_i + \mu, \lambda_i + \mu).$$

Now look at  $H(\lambda_i)$ . Observe that  $H(\lambda_i) = H(\lambda_i + \mu)$ . Indeed,  $\lambda_i + \mu \in w_i\Lambda_G^+$  and from the first part of the lemma it follows that  $H(\lambda_i) \supset H(\lambda_i + \mu)$ . The opposite inclusion follows from the fact that since  $\mu$  and  $\lambda_i$  lie in the same chamber there are no hyperplanes  $H_\alpha$  separating them, i.e. one has  $(\lambda_i, \alpha) \cdot (\mu, \alpha) \geq 0$ . For every  $H_\alpha \in H(\lambda_i)$  one has  $(\lambda_i, \alpha) < 0$ , so  $(\mu, \alpha) \leq 0$  and  $(\lambda_i + \mu, \alpha) < 0$ , then  $H_\alpha \in H(\lambda_i + \mu)$ .

First suppose that there exists some  $H_\alpha \in H(\lambda_i)$  such that  $(w_jw_i^{-1}\mu, \alpha) > 0$ . Since  $H_\alpha \in H(\lambda_i)$  one has  $(\lambda_i, \alpha) < 0$  and

$$(\lambda_i, \alpha) \cdot (w_jw_i^{-1}\mu, \alpha) < 0.$$

Then we are in the setting of lemma 4 with a slight change of notation, so  $(\lambda_i, w_jw_i^{-1}\mu) < (\lambda_i, \mu)$  hence  $\lambda_i + w_jw_i^{-1}\mu \preceq \lambda_i + \mu$ .

Otherwise for all  $H_\alpha \in H(\lambda_i)$  one has  $(w_j w_i^{-1} \mu, \alpha) \leq 0$  and since  $(\lambda_i, \alpha) < 0$  one has  $(\lambda_i + w_j w_i^{-1} \mu, \alpha) < 0$ , so  $H(\lambda_i) \subset H(w_j w_i^{-1} \mu + \lambda_i)$  and

$$H(\lambda_i + \mu) \subset H(w_j w_i^{-1} \mu + \lambda_i).$$

The last two sets coincide only if  $w_j w_i^{-1} \mu + \lambda_i \in w_i \Lambda_G^+$  and the second part of the lemma in this case yields  $w_j w_i^{-1} \mu \in w_i \Lambda_G^+$  that means  $\mu = w_j w_i^{-1} \mu$ . In any case one has  $\lambda_i + w_j w_i^{-1} \mu \preceq \lambda_i + \mu$ .  $\square$

*Remark 4.* Note that from the above construction one gets  $\lambda_1 = 0 \in X^*(T)$ .

#### 4. RESTRICTION OF REPRESENTATIONS.

In this section we study the representation restriction homomorphism on the representation rings and prove theorem 2.

Let  $T \subset H \subset G$  be the same groups as in the previous section. From theorem 1 we get the following commutative diagram.

$$\begin{array}{ccccc} R(G) & \hookrightarrow & R(H) & \hookrightarrow & R(T) \\ \downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\ \mathbb{Z}[X^*(T)]^{W(G)} & \hookrightarrow & \mathbb{Z}[X^*(T)]^{W(H)} & \hookrightarrow & \mathbb{Z}[X^*(T)] \end{array}$$

Recall that  $Z(G) \subset T$ , hence all the rings above are  $Ch = X^*(Z(G))$ -graded. We are interested in  $R(G)$ -module structure on  $R(H)$  and its connection with the grading.

We need the following easy lemma from commutative algebra.

**Lemma 7.** *Let  $S \subset R$  be domains, let  $\lambda_1, \dots, \lambda_k \in R$  generate  $R$  as  $S$ -module and let  $[Q(R) : Q(S)] = k$ . Then  $R$  is a free  $S$ -module with basis  $\lambda_1, \dots, \lambda_k$ .*

*Proof.*  $R$  is finitely generated as  $S$ -module hence it is integral over  $S$ . Then  $R \otimes_S Q(S)$  is integral over field  $S \otimes_S Q(S) = Q(S)$ , hence itself is a field so  $R \otimes_S Q(S) \cong Q(R)$ .

We have the following short exact sequence induced by  $\lambda_1, \dots, \lambda_k$ .

$$N \twoheadrightarrow S^k \twoheadrightarrow R,$$

hence

$$N \otimes_S Q(S) \twoheadrightarrow Q(S)^k \twoheadrightarrow R \otimes_S Q(S).$$

The last term is isomorphic to  $Q(R)$  and comparing dimensions one can see that  $N \otimes_S Q(S) = 0$  hence  $N = 0$  and we get the claim of the lemma.  $\square$

We consider  $X^*(T)$  as an additive group, so we will write the element of  $\mathbb{Z}[X^*(T)]$  corresponding to weight  $\mu$  in such way:  $x^\mu$ .

**Definition 4.** For  $\mu \in X^*(T)$  we will denote by  $(x^\mu)^{W(H)}$  the sum in the  $\mathbb{Z}[X^*(T)]$  of all elements corresponding to the weights in  $W(H)$ -orbit of  $\mu$  and for any monomial  $ax^\mu \in \mathbb{Z}[X^*(T)]$  by  $(ax^\mu)^{W(H)} = a(x^\mu)^{W(H)}$  we denote the similar orbit but with a coefficient.



With the above notation one has the unique decomposition of

$$f = \sum (a_{\mu_j} x^{\mu_j})^{W(H)} \in \mathbb{Z}[X^*(T)]^{W(H)}$$

into the sum of monomial orbits with distinct  $\mu_j \in \Lambda_H^+$ . Recall that we have a linear order  $\preceq$  on  $\mu_j$  introduced in the previous section.

**Definition 5.** Let  $f = \sum (a_{\mu_j} x^{\mu_j})^{W(H)}$ , then define the degree  $\deg(f) = \max_j \mu_j$  to be the maximal  $\mu_j$  in the decomposition and the leading orbit  $\text{lo}(f) = (a_{\deg(f)} x^{\deg(f)})^{W(H)}$  to be the orbit of the maximal monomial.

We will use the analogous notation for group  $G$ .

**Theorem 2.** *Let  $G$  be a split semisimple simply connected group and let  $H$  be a connected split reductive subgroup of the maximal rank (i.e.  $H$  contains the split maximal torus  $T$  of  $G$ ). Then  $R(H)$  is a free  $R(G)$ -module of rank  $[W(G) : W(H)]$  and there is a Ch-homogeneous basis.*

*Proof.* First of all we will deal with the weight realization of the rings of representations, i.e. with the following sequence.

$$\mathbb{Z}[X^*(T)]^{W(G)} \hookrightarrow \mathbb{Z}[X^*(T)]^{W(H)} \hookrightarrow \mathbb{Z}[X^*(T)]$$

In the previous section in lemma 6 we have constructed some  $\lambda_i$  and we claim that the orbits  $(x^{\lambda_i})^{W(H)}$  form a homogeneous basis of  $\mathbb{Z}[X^*(T)]^{W(H)}$  over  $\mathbb{Z}[X^*(T)]^{W(G)}$ .

- (a) Homogeneity. It is the easiest part since it follows at once from the equivariant  $e$  of  $W(H)$  action.
- (b)  $(x^{\lambda_i})^{W(H)}$  generate  $\mathbb{Z}[X^*(T)]^{W(H)}$  as  $\mathbb{Z}[X^*(T)]^{W(G)}$ -module. We will show by induction on  $\deg(f)$  that  $f \in \mathbb{Z}[X^*(T)]^{W(H)}$  could be expressed as linear combination of  $(x^{\lambda_i})^{W(H)}$  with  $\mathbb{Z}[X^*(T)]^{W(G)}$  coefficients. Note that  $\lambda_1 = 0 \in X^*(T)$  and

$$(x^{\lambda_1})^{W(H)} = 1 \in \mathbb{Z}[X^*(T)]^{W(H)},$$

so we have the constants. Now suppose that we can express as linear combinations all  $f \in \mathbb{Z}[X^*(T)]^{W(H)}$  such that  $\deg(f) \prec \mu_0$  and we need to write such an expression for  $(x^{\mu_0})^{W(H)}$ .

By lemma 3 we have some chamber  $w_l \mathcal{C}(G)$  such that  $\mu_0 \in w_l \Lambda_G^+$  and  $H(\mu_0) = H(w_l \mathcal{C}(G))$ , hence, by lemma 6  $\nu = \mu_0 - \lambda_l \in w_l \Lambda_G^+$ . Choose the subset  $\{w_j\}$  of  $\{w_i\}$  such that one has all the distinct  $w_j w_l^{-1} \nu$ . Then by subdividing  $W(G)$ -orbit into  $W(H)$ -orbits we have the following equality.

$$\text{lo} \left( (x^\nu)^{W(G)} (x^{\lambda_l})^{W(H)} \right) = \text{lo} \left( \left( \sum_j (x^{w_j w_l^{-1} \nu})^{W(H)} \right) (x^{\lambda_l})^{W(H)} \right).$$

From lemma 1 and lemma 5 the last one equals to

$$\text{lo} \left( \left( \sum_j x^{w_j w_l^{-1} \nu} x^{\lambda_l} \right)^{W(H)} \right) = \text{lo} \left( \left( \sum_j x^{w_j w_l^{-1} \nu + \lambda_l} \right)^{W(H)} \right),$$

and finally, by lemma 6, one gets

$$\text{lo} \left( \left( \sum_j x^{w_j w_i^{-1} \nu + \lambda_i} \right)^{W(H)} \right) = (x^{\nu + \lambda_i})^{W(H)} = (x^{\mu_0})^{W(H)}.$$

Hence

$$\text{deg} \left( (x^{\mu_0})^{W(H)} - (x^\nu)^{W(G)} (x^{\lambda_i})^{W(H)} \right) < \mu_0$$

and we can use the induction.

(c)  $(x^{\lambda_i})^{W(H)}$  are linearly independent. From the sequence

$$\mathbb{Z}[X^*(T)]^{W(G)} \hookrightarrow \mathbb{Z}[X^*(T)]^{W(H)} \hookrightarrow \mathbb{Z}[X^*(T)],$$

one gets the sequence of fraction fields

$$\begin{array}{ccccc} Q(\mathbb{Z}[X^*(T)]^{W(G)}) & \hookrightarrow & Q(\mathbb{Z}[X^*(T)]^{W(H)}) & \hookrightarrow & Q(\mathbb{Z}[X^*(T)]) \\ \parallel & & \parallel & & \parallel \\ Q(\mathbb{Z}[X^*(T)])^{W(G)} & \hookrightarrow & Q(\mathbb{Z}[X^*(T)])^{W(H)} & \hookrightarrow & Q(\mathbb{Z}[X^*(T)]) \end{array}$$

and the degree of field extension equals to  $[W(G) : W(H)]$ . Hence, by lemma 7 one gets the claim of the theorem.  $\square$

**Corollary 1.** *In the notation of theorem 2 one has a basis consisting of the irreducible representations of  $H$ .*


*Proof.* One can take  $V_H(\lambda_i)$  and since their leading orbits coincide with the basis constructed in the theorem one gets the claim.  $\square$

*Remark 5.* We can choose various chambers  $\mathcal{C}(G)$  and the different choices produce different bases. Also the proof of the theorem gives an explicit algorithm for calculating the coefficients of decomposition with respect to the chosen basis, so, for example, in every particular case one can write down the multiplication table for the basis, yet it seems that there is no elegant general formula.

## 5. EXAMPLES: $A_1 + A_1 \subset G_2$ , $B_4 \subset F_4$ AND $C_1 + C_{n-1} \subset C_n$ .

In this section we compute some examples of bases. Every reductive subgroup containing the maximal torus is defined by some quasi-closed root subset [BT, § 3], so we use the root system notation. Every maximal root subsystem of full rank corresponds to some node in the Dynkin diagram and the subsystem diagram is just the extended Dynkin diagram with the chosen node removed. We label simple roots in a way of [Bo].

5.1.  $A_1 + A_1 \subset G_2$ . In this example we take the the subsystem in  $G_2$  defined by the short simple root and the maximal one. The corresponding Dynkin diagram is the next one, with the white node removed.



We label the roots in a way shown above and  $\alpha_{max} = 3\alpha_1 + 2\alpha_2$ . The fundamental chamber  $\mathcal{C}(G_2)$  is the chamber spanned by the fundamental weights

$$\omega_1 = \alpha_{max} = 3\alpha_1 + 2\alpha_2, \quad \omega_2 = 2\alpha_1 + \alpha_2.$$

The fundamental chamber  $\mathcal{C}(A_1 + A_1)$  should contain  $\mathcal{C}(G_2)$  so it is the quarter of the plane bounded by  $\alpha_1$  and  $\alpha_{max}$ . Note, by the way, that the considered group  $G(A_1 + A_1)$  is not simply connected since there are no weights  $\frac{1}{2}\alpha_1$  and  $\frac{1}{2}\alpha_{max}$  in our lattice and one can see that

$$G(A_1 + A_1) = SL_2 \otimes SL_2 = (SL_2 \times SL_2)/\mu_2$$

with  $\mu_2$  embedded diagonally.

The chamber  $\mathcal{C}(A_1 + A_1)$  subdivides into the  $G_2$  chambers in the following way:

$$\mathcal{C}(A_1 + A_1) = \mathcal{C}(G_2) \cup w_{\alpha_2}\mathcal{C}(G_2) \cup w_{\alpha_1+\alpha_2}w_{\alpha_2}\mathcal{C}(G_2)$$

So the theorem 2 tells us that we should take in each subchamber the shortest of the furthest by spherical distance weights, i.e. the generator for the furthest wall, hence one has

$$0 \in \mathcal{C}(G), \quad 3\alpha_1 + \alpha_2 \in w_{\alpha_2}\mathcal{C}(G), \quad \alpha_1 \in w_{\alpha_1+\alpha_2}w_{\alpha_2}\mathcal{C}(G_2)$$

and the corresponding sums over  $W(A_1 + A_1)$  would form the basis. The basis from the theorem is the follows:

$$1, \quad x^{3\alpha_1+\alpha_2} + x^{\alpha_2} + x^{-\alpha_2} + x^{-3\alpha_1-\alpha_2}, \quad x^{\alpha_1} + x^{-\alpha_1}.$$


One could compute the corresponding basis consisting of irreducible modules from corollary 1 having the following weight subspaces:

$$\begin{aligned} V(0) &= 1 \\ V(3\alpha_1 + \alpha_2) &= x^{2\alpha_1+\alpha_2} + x^{\alpha_1+\alpha_2} + x^{-\alpha_1-\alpha_2} + x^{-2\alpha_1-\alpha_2} + \\ &\quad + x^{3\alpha_1+\alpha_2} + x^{\alpha_2} + x^{-\alpha_2} + x^{-3\alpha_1-\alpha_2} \\ V(\alpha_1) &= 1 + x^{\alpha_1} + x^{-\alpha_1} \end{aligned}$$

In fact after identifying  $G(A_1 + A_1) = SL_2 \otimes SL_2$  one can write the above representations in more natural way, denoting by  $W_1, W_2$  the regular representations of the factors one has

$$V(0) = S^0W_1 \otimes S^0W_2, \quad V(3\alpha_1 + \alpha_2) = S^3W_1 \otimes W_2, \quad V(\alpha_1) = S^2W_1 \otimes S^0W_2.$$

5.2.  $B_4 \subset F_4$ . In this case we remove the  $\alpha_4$  node from the extended Dynkin diagram of type  $F_4$ . One can show that it corresponds to  $Spin_9 \subset G(F_4)$ .



With the above labeling one has

$$\alpha_{max} = 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 2\alpha_4.$$

We will shorten the above notation to  $(2, 3, 4, 2)$ . The fundamental weights defining  $\mathcal{C}(F_4)$  are

$$\omega_1 = (2, 3, 4, 2), \quad \omega_2 = (3, 6, 8, 4), \quad \omega_3 = (2, 4, 6, 3), \quad \omega_4 = (1, 2, 3, 2).$$

Choose the simple roots for  $B_4$  in the following way:

$$\alpha'_1 = w_{\alpha_4}(-\alpha_{max}) = (0, 1, 2, 2), \quad \alpha'_2 = \alpha_1, \quad \alpha'_3 = \alpha_2, \quad \alpha'_4 = \alpha_3,$$

hence we have the fundamental weights defining  $\mathcal{C}(B_4) \sim \{\omega'_1, \omega'_2, \omega'_3, \omega'_4\}$ :

$$\omega'_1 = (1, 2, 3, 2) = \omega_4, \quad \omega'_2 = (2, 3, 4, 2) = \omega_1,$$

$$\omega'_3 = (2, 4, 5, 2), \quad \omega'_4 = (1, 2, 3, 1).$$

Since these weights belong to the considered lattice the chosen  $G(B_4)$  is simply connected, so it really is  $Spin_9$ . Now we compute the subdividing of  $\mathcal{C}(B_4)$ :


$$\mathcal{C}(B_4) = \mathcal{C}(F_4) \cup w_{\alpha_4}\mathcal{C}(F_4) \cup w_{\alpha_3+\alpha_4}w_{\alpha_4}\mathcal{C}(F_4),$$

$$w_{\alpha_4}\mathcal{C}(B_4) \sim \{\omega_1, \omega_2, \omega_3, \omega'_4\}, \quad w_{\alpha_3+\alpha_4}w_{\alpha_4}\mathcal{C}(B_4) \sim \{\omega_1, \omega_2, \omega'_3, \omega'_4\}.$$

Theorem 2 suggests to look at the elements appeared after flips, since they are the spherically furthest, so the basis would consist of  $W(B_4)$  orbits of  $0, \omega'_4, \omega'_3$ .

Another basis comes from the corollary 1 that claims  $V(0), V(\omega'_3), V(\omega'_4)$  to be a basis. These representations are just the trivial one, the  $\Lambda^3 W$  for the regular  $W$  and the spin one.

5.3.  $C_1 + C_{n-1} \subset C_n$ . In this case we remove the  $\alpha_1$  node from the extended Dynkin diagram of type  $C_n$  and it corresponds to  $(Sp_2 \times Sp_{2n-2}) \subset Sp_{2n}$  with the quotient variety  $HP^{n-1} = Sp_{2n}/(Sp_2 \times Sp_{2n-2})$  being the quaternionic projective space in notation of section 9.3.



One has

$$\alpha_{\max} = 2\alpha_1 + 2\alpha_2 + \dots + 2\alpha_{n-1} + \alpha_n = (2, 2, \dots, 2, 1).$$

The fundamental weights for  $\mathcal{C}(C_n)$  are

$$\omega_i = (1, 2, \dots, i-1, i, i, \dots, i, \frac{i}{2}).$$

Choosing  $\alpha_2, \alpha_3, \dots, \alpha_n, \alpha_{\max}$  to be the simple roots of  $C_1 + C_{n-1}$  one gets the following  $\mathcal{C}(C_1 + C_{n-1})$ :

$$\omega'_i = (0, 1, \dots, i-1, i, i, \dots, i, \frac{i}{2}), \quad \omega'_n = (1, 1, \dots, 1, \frac{1}{2}).$$

The subdividing of  $\mathcal{C}(C_1 + C_{n-1})$  is straightforward:

$$\begin{aligned} \mathcal{C}(C_1 + C_{n-1}) = & \mathcal{C}(C_n) \cup w_{\alpha_1}\mathcal{C}(C_n) \cup w_{\alpha_1+\alpha_2}w_{\alpha_1}\mathcal{C}(C_n) \cup \dots \\ & \dots \cup (w_{\alpha_1+\dots+\alpha_{n-1}} \dots w_{\alpha_1+\alpha_2}w_{\alpha_1})\mathcal{C}(C_n), \end{aligned}$$

$$(w_{\alpha_1+\dots+\alpha_i} \dots w_{\alpha_1+\alpha_2}w_{\alpha_1})\mathcal{C}(C_n) \sim \{\omega'_1, \omega'_2, \dots, \omega'_i, \omega_{i+1}, \dots, \omega_n\}.$$

The corollary 1 claims  $V(0), V(\omega'_1), \dots, V(\omega'_{n-1})$  to be the basis and this representations are just  $\Lambda^i W$  for regular representation  $W$  of group  $Sp_{2n-2}$ .

6. REPRESENTATIONS, VECTOR BUNDLES AND EQUIVARIANT  $K$ -THEORY.

In this section we recall some results on the equivariant  $K$ -theory. An extensive exposition and further references could be found in [M].

Let  $G$  be an algebraic group, let  $H \subset G$  be a closed subgroup and let  $X = G/H$  be the corresponding smooth homogeneous  $G$ -variety. There is a well-known tensor equivalence [M, Example 2]

$$\text{Rep}_k(H) \xrightarrow{\sim} \text{Vect}^G(X)$$

between the categories  $\text{Rep}_k(H)$  of finite dimensional  $k$ -rational representations of  $H$  and  $\text{Vect}^G(X)$  of  $G$ -equivariant vector bundles over  $X$ . The inverse for the above equivalence is given by the fiber over the extinguished point  $eH$  of  $X$ . Further we will use the following notation.

**Definition 6.** Let  $V_H(\lambda)$  be the irreducible representation of  $H$  with the highest weight  $\lambda \in \Lambda_H^+$ , then denote by  $\mathcal{V}_H(\lambda)$  the corresponding vector bundle over  $G/H$ . For an irreducible representation  $V_G(\mu)$  of group  $G$  with the highest weight  $\mu \in \Lambda_G^+$  one can use the restriction of representations, get the representation of  $H$  (not necessary irreducible) and then take the corresponding vector bundle  $\mathcal{V}_G(\mu)$ . Occasionally we will write  $V_G(\lambda)$  and  $\mathcal{V}_G(\lambda)$  for a  $\lambda \in \Lambda_H^+$  and it means that one should find  $\mu \in \Lambda_G^+$  from  $W(G)$ -orbit of  $\lambda$  and then take the corresponding  $V_G(\mu)$  and  $\mathcal{V}_G(\mu)$ .

*Remark 6.* Note that after forgetting about the  $G$ -action the last bundle becomes trivial, i.e. the composition

$$\text{Rep}_k(G) \xrightarrow{\text{Res}} \text{Rep}_k(H) \xrightarrow{\sim} \text{Vect}^G(X) \longrightarrow \text{Vect}(X)$$

takes  $G$  representations to trivial bundles.

Set

$$K_n(G; X) = K_n(\text{Vect}^G(X)).$$

The above equivalence yields

$$K_n(G; X) \cong K_n(\text{Rep}(H)),$$

in particular

$$K_0(G; X) \cong R(H).$$

Note that  $R(H)$  is a  $R(G)$ -module, hence every  $K_n(G, X)$  also is. The following proposition, being a straightforward consequence of [M, Theorem 10], compares  $K_n(G; X)$  with  $K_n(X)$ .

**Proposition 1.** *Let  $G$  be a split simply connected semisimple group. Then there is a spectral sequence*

$$E_{p,q}^2 = \text{Tor}_p^{R(G)}(\mathbb{Z}, K_q(G; X)) \implies K_{p+q}(X).$$

## 7. K-THEORY OF A HOMOGENEOUS VARIETY.

In this section we calculate  $K$ -theory of a homogeneous variety  $X = G/H$  for connected split reductive algebraic groups  $H \subset G$  of the same rank.

**Lemma 8.**  $K_n(\text{Rep}_k(H)) \cong R(H) \otimes_{\mathbb{Z}} K_n(k)$ .

*Proof.* Note that  $\text{char } k$  not necessary equals 0, so the reductive group  $H$  not necessary geometrically reductive, i.e. the category  $\text{Rep}_k(H)$  may be not semisimple. But, nevertheless, all objects of  $\text{Rep}_k(H)$  have finite length and, thanks to Devissage property of  $K$ -theory, one has

$$K_n(\text{Rep}_k(H)) \cong K_n(\text{Rep}_k(H)_{ss}),$$

where  $\text{Rep}_k(H)_{ss}$  stands for the subcategory of semisimple representations. By Shur's Lemma we can pass to the endomorphisms of irreducible representations

$$K_n(\text{Rep}_k(H)_{ss}) \cong \bigoplus_i K_n(\text{End}(V_i)),$$

and, since  $\text{End}(V_i) = k$ , the last one equals  $\bigoplus K_n(k)$  with the sum over isomorphism classes of irreducible representations, which could be identified with  $R(H) \otimes_{\mathbb{Z}} K_n(k)$ .  $\square$

**Proposition 2.** *Let  $G$  be a connected split simply connected semisimple algebraic group and let  $H \subset G$  be a connected split reductive subgroup of the same rank. Then the spectral sequence in proposition 1 degenerates, i.e.*

$$\text{Tor}_p^{R(G)}(\mathbb{Z}, K_n(G; X)) = \begin{cases} K_n(X), & \text{if } p = 0; \\ 0, & \text{if } p > 0. \end{cases}$$

*Proof.* Due to lemma 8 it is sufficient to show that for  $p \geq 1$  one has

$$\text{Tor}_p^{R(G)}(\mathbb{Z}, R(H) \otimes_{\mathbb{Z}} K_n(k)) = 0.$$

Replace  $K_n(k)$  with an arbitrary abelian group  $M$ . Since  $\text{Tor}$  commutes with the direct limits we can reduce the problem to the finitely generated abelian groups, and, moreover, to  $M = \mathbb{Z}$  or  $M = \mathbb{Z}/m\mathbb{Z}$ .

In the first case we at once get the claim from theorem 2,

$$\text{Tor}_p^{R(G)}(\mathbb{Z}, R(H)) = 0,$$

since  $R(H)$  is a free  $R(G)$ -module.

In the second case we can write the resolution

$$0 \longrightarrow R(H) \xrightarrow{m} R(H) \longrightarrow R(H) \otimes_{\mathbb{Z}} \mathbb{Z}/m\mathbb{Z} \longrightarrow 0,$$

which is exact since  $R(H)$  is a domain of a zero characteristic. Denoting the rank of  $R(H)$  over  $R(G)$  by  $r$ , after tensoring with  $\mathbb{Z}$  one still gets an exact sequence

$$0 \longrightarrow \mathbb{Z}^r \xrightarrow{m} \mathbb{Z}^r \longrightarrow (\mathbb{Z}/m\mathbb{Z})^r \longrightarrow 0.$$

So, we conclude

$$\text{Tor}_p^{R(G)}(\mathbb{Z}, R(H) \otimes \mathbb{Z}/m\mathbb{Z}) = 0,$$

finishing the proof.  $\square$

In order to remove the annoying restriction that  $G$  should be simply connected we need the following lemma.

**Lemma 9.** *Let  $H \subset G$  be a pair of connected split reductive groups of the same rank. Then there exists a connected split simply connected semisimple group  $\tilde{G}$  and a connected split reductive subgroup  $\tilde{H} \subset \tilde{G}$  of the same rank such that  $\tilde{G}/\tilde{H} \cong G/H$ .*

*Proof.* Let  $\tilde{G}$  be the simply connected covering of the derived group  $G'$ . There exists a covering

$$Z \twoheadrightarrow (G_m)^l \times \tilde{G} \twoheadrightarrow G$$

with finite kernel  $Z$ . Since  $H$  contains the maximal torus the preimage of  $H$  under this projection contains the factor  $(G_m)^l$ , so we have the following diagram.

$$\begin{array}{ccc} (G_m)^l \times \tilde{G} & \twoheadrightarrow & G \\ \uparrow & & \uparrow \\ (G_m)^l \times \tilde{H} & \twoheadrightarrow & H \end{array}$$

The above consideration yields

$$\tilde{G}/\tilde{H} \cong (G_m)^l \times \tilde{G} / (G_m)^l \times \tilde{H} \cong G/H,$$

so we need to show that  $\tilde{H}$  is connected. One has

$$Z \twoheadrightarrow (G_m)^l \times \tilde{H} \twoheadrightarrow H$$

with finite central  $Z$  and connected  $H$ . The identity component of  $(G_m)^l \times \tilde{H}$  contains the maximal torus, hence it contains  $Z$  and the connectedness of  $H$  yields that the identity component coincides with the whole group. The group  $\tilde{H}$  is connected as the quotient of the connected group.  $\square$

**Theorem 3.** *Let  $H \subset G$  be a pair of connected split reductive groups of the same rank. Denote  $r = [W(G) : W(H)]$ . Then there exist  $\mathcal{V}_1, \dots, \mathcal{V}_r \in \text{Vect}(G/H)$  such that*

$$K_*(G/H) = \bigoplus_{i=1}^r K_*(k)[\mathcal{V}_i].$$

*Proof.* By lemma 9 we can pass to a simply connected semisimple group  $G$  and from the proof one has that  $[W(G) : W(H)]$  remains the same. Due to proposition 2, the spectral sequence in proposition 1 degenerates, so, using lemma 8, one has

$$\begin{aligned} K_n(X) &\cong \text{Tor}_0^{R(G)}(\mathbb{Z}, R(H) \otimes_{\mathbb{Z}} K_n(k)) = \\ &= \mathbb{Z} \otimes_{R(G)} R(H) \otimes_{\mathbb{Z}} K_n(k) = \mathbb{Z}^r \otimes_{\mathbb{Z}} K_n(k). \end{aligned}$$

The above isomorphism is induced by the elements of basis  $R(H)$  over  $R(G)$  constructed in theorem 2, so we can take as  $\mathcal{V}_i$  the corresponding elements of  $\text{Vect}(G/H)$ .  $\square$

*Remark 7.* In remark 5 we noted that there is an explicit algorithm to write down the multiplication for the basis elements and now it describes the ring structure on the  $K_0(G/H)$ . Also one can drop the assumption of the connectedness of  $G$  (but not the  $H$ ), since  $G/H$  is just the disjoint union of  $[G : G_0]$  varieties isomorphic to  $G_0/H$  and so be the  $K$ -theory.

## 8. $K$ -THEORY OF TWISTED FORMS.

In this section we deal with certain twisted forms of homogeneous varieties  $X = G/H$  with connected split reductive groups  $H \subset G$  of the same rank. From now on we suppose  $G$  to be simply connected semisimple, lemma 9 shows that in fact it is not a restriction. Denote as before  $r = [W(G) : W(H)]$ .

One has an obvious left action of  $G$  on  $G/H$  and since  $H$  contains the maximal torus hence the center, this action extends to the action of

$$\overline{G} = G/Z(G).$$

Now fix a 1-cocycle  $\gamma : \text{Gal}(k^{sep}/k) \rightarrow \overline{G}(k^{sep})$ . Twisting the variety with this cocycle we obtain

$$X_\gamma = (G/H)_\gamma.$$

The following lemma provides a splitting variety for such cocycle.

**Lemma 10.** *For the above cocycle  $\gamma$  there exists a variety  $Y$  such that the following conditions hold:*

- (1)  *$Y$  is a smooth projective variety.*
- (2) *The Euler characteristic  $\chi(Y)$  equals to 1.*
- (3) *For every point (not necessary closed)  $y \in Y$  the cocycle  $\gamma_{k(y)}$  is a coboundary.*

*Proof.* Note that  $\overline{G}$  is split semisimple group and we can twist it with  $\gamma$  as well. The last condition is equivalent to the condition that for every point  $y \in Y$  the group  $(\overline{G}_\gamma)_{k(y)}$  is split. Consider

$$Y = (G/B)_\gamma.$$

We claim that for the Borel subgroup  $B \subset \overline{G}$  the variety

$$Y_F = ((G/B)_\gamma)_F$$

has rational point if and only if  $G_\gamma$  splits over  $F$ . The existence of a rational point on this variety is equivalent to the existence of a Borel subgroup defined over  $F$ , which is the stabilizer of this point, in  $(\overline{G}_\gamma)_F$ . The existence of Borel subgroup means that group is quasi-split that in our case is equivalent to be split, since we work with inner form.

$Y = (G/B)_\gamma$  is clearly a smooth projective variety. In order to compute the Euler characteristic  $\chi(G/B)$  it can be shown [Ja, Proposition 4.5] that

$$h^i(O_{G/B}) = \begin{cases} 0, & \text{if } i > 0; \\ 1, & \text{if } i = 0. \end{cases}$$

so we get the claim. □



The idea lying behind the calculation of  $K$ -theory of twisted form is quite simple: one needs to construct some candidate for the  $K$ -theory and a morphism such that they will produce the correct answer and an isomorphism in the split case. The isomorphism in the split case is written by means of some vector bundles  $\mathcal{V}_i$ , so in general we want to twist them. And here is the problem – there is no action of  $\overline{G}$  on them since the center could act non-trivially. In order to get over that we should tensor  $\mathcal{V}_i$  with some bundles to trivialize the center action, and then for cancellation of this tensoring we should look at modules over the endomorphisms of the excessive factors. Twisting these endomorphisms algebras we get the separable algebras which produce the answer.

**Definition 7.** Let  $V_G(\lambda)$  be a representation of  $G$  then we denote

$$A(\lambda) = \text{End}_k(V_G(\lambda)) = V_G(\lambda) \otimes V_G^*(\lambda)$$

the endomorphism algebra of the underlying vector space. There is an obvious  $G$  action on  $A(\lambda)$  which extends to the  $\overline{G}$  action, so we can twist this algebra and get the separable algebra  $A(\lambda)_\gamma$ . Also one can pass to the corresponding trivial sheaf of algebras  $\mathcal{A}(\lambda)_\gamma$  over  $X_\gamma$ .

Now we fix  $\lambda_i$  from theorem 2 and the corresponding  $A(\lambda_i)$ . Denote

$$W(\lambda_i) = V_H(\lambda_i) \otimes V_G(\lambda_i),$$

and the corresponding vector bundle  $\mathcal{W}(\lambda_i)$ . Note that  $W(\lambda_i)$  is a right module over  $A(\lambda_i)$  through the second factor and so  $\mathcal{W}(\lambda_i)$  is. The center acts trivially on  $W(\lambda_i)$  and  $\mathcal{W}(\lambda_i)$  so one can obtain the twisted form  $\mathcal{W}(\lambda_i)_\gamma$ .

All the considered above structures are agreed, so now we have trivial sheafs of separable algebras  $\mathcal{A}(\lambda_i)_\gamma$  and vector bundles  $\mathcal{W}(\lambda_i)_\gamma$  that are right  $\mathcal{A}(\lambda_i)_\gamma$ -modules.

**Definition 8.** For a variety  $Z$  and a separable algebra  $A$  let  $\mathcal{P}(Z, A)$  be the category of coherent  $O_Z \otimes A$ -modules which are locally free  $O_Z$ -modules. Then we denote

$$K_*(Z, A) = K_*(\mathcal{P}(Z, A)).$$

There is a corresponding notion of  $K'_*(Z, A)$  and it satisfies all the usual properties of  $K$ -theory [M].

**Proposition 3.** *Let  $Z$  be a variety such that every point  $z \in Z$  (not necessary closed) splits  $\gamma$ , i.e.  $\gamma_{k(z)}$  is a coboundary. Then in the above notation one has an isomorphism*

$$\sum_{i=1}^r \phi_i : \bigoplus_{i=1}^r K'_*(Z, A(\lambda_i)_\gamma) \longrightarrow K'_*(X_\gamma \times Z),$$

where

$$\phi_i(U) = p_X^*(\mathcal{W}(\lambda_i)_\gamma) \otimes_{A(\lambda_i)_\gamma} p_Z^*(U).$$

*Proof.* This is proved by induction on the variety dimension.

Suppose first that  $\dim Z = 0$ , i.e.  $Z$  is a point, then we are in fact in the split case. Let  $F = k(Z)$ , so we have

$$X_\gamma \times Z = X_F, \quad (A(\lambda_i)_\gamma)_F = A(\lambda_i)_F = \text{End}_F(V_G(\lambda_i) \otimes F),$$

$$(\mathcal{W}(\lambda_i)_\gamma)_F = \mathcal{W}(\lambda_i)_F = \mathcal{V}_H(\lambda_i) \otimes \mathcal{V}_G^*(\lambda_i) \otimes F.$$

Since every module over  $\text{End}_F(V_G(\lambda_i) \otimes F)$  is isomorphic to  $V_G(\lambda_i) \otimes F^n$  and

$$V_G^*(\lambda_i) \otimes F \otimes_{\text{End}_F(V_G(\lambda_i) \otimes F)} V_G(\lambda_i) \otimes F \cong F,$$

one has

$$\begin{aligned} \phi_i(V_G(\lambda_i) \otimes F^n) &= \mathcal{V}_H(\lambda_i) \otimes \mathcal{V}_G^*(\lambda_i) \otimes F \otimes_{\text{End}_F(V_G(\lambda_i) \otimes F)} \mathcal{V}_G(\lambda_i) \otimes F^n = \\ &= \mathcal{V}_H(\lambda_i) \otimes F^n. \end{aligned}$$

The above considerations show that we are in the setting of theorem 3 claiming  $\sum \phi_i$  to be an isomorphism.

For the dimension greater than 0 we can write the localization sequence for all subvarieties  $Z' \subset Z$  of codimension one, so for  $F = k(Z)$  one has

$$\begin{array}{ccccc} \varinjlim_{Z' \subset Z} \bigoplus_{i=1}^r K'_*(Z', A(\lambda_i)_\gamma) & \longrightarrow & \bigoplus_{i=1}^r K'_*(Z, A(\lambda_i)_\gamma) & \longrightarrow & \bigoplus_{i=1}^r K'_*(\text{Spec } F, A(\lambda_i)_\gamma) \\ \downarrow & & \downarrow \Sigma \phi_i & & \downarrow \\ \varinjlim_{Z' \subset Z} K'_*(X_\gamma \times Z') & \longrightarrow & K'_*(X_\gamma \times Z) & \longrightarrow & K'_*(X_\gamma \times \text{Spec } F) \end{array}$$

This sequence extends to the right and to the left with the shifts in  $K$ -theory, and both the side vertical morphisms in each triple are isomorphisms by induction, so using the five lemma one concludes that the middle one is an isomorphism.  $\square$

**Corollary 2.** *In the notation of proposition 3 for the smooth  $Z$  one has*

$$\sum_{i=1}^r \phi_i : \bigoplus_{i=1}^r K_*(Z, A(\lambda_i)_\gamma) \longrightarrow K_*(X_\gamma \times Z).$$

*Proof.* One has  $K_*(Z, A) = K'_*(Z, A)$  and  $K_*(X_\gamma \times Z) = K'_*(X_\gamma \times Z)$ .  $\square$

**Theorem 4.** *In the above notation there is an isomorphism*

$$\sum_{i=1}^r \psi_i : \bigoplus_{i=1}^r K_*(A(\lambda_i)_\gamma) \longrightarrow K_*(X_\gamma),$$

where

$$\psi_i(U) = \mathcal{W}(\lambda_i)_\gamma \otimes_{A(\lambda_i)_\gamma} U.$$

*Proof.* We can insert two copies of our morphism into the following diagram with the middle arrow from corollary 2 and  $Y$  being the splitting variety

constructed in lemma 10.

$$\begin{array}{ccc}
 \bigoplus_{i=1}^r K_*(A(\lambda_i)_\gamma) & \xrightarrow{\sum \psi_i} & K_*(X_\gamma) \\
 \downarrow \chi(Y) \cdot & \searrow p^* & \swarrow p^* \\
 & \bigoplus_{i=1}^r K_*(Y, A(\lambda_i)_\gamma) \xrightarrow{\sum \phi_i} K_*(X_\gamma \times Y) & \\
 & \swarrow p_* & \searrow p_* \\
 \bigoplus_{i=1}^r K_*(A(\lambda_i)_\gamma) & \xrightarrow{\sum \psi_i} & K_*(X_\gamma) \\
 & & \downarrow \chi(Y) \cdot
 \end{array}$$

A direct verification shows that the diagram is commutative. The vertical morphisms are just multiplications by  $\chi(Y) = 1$  since they are equal to the composition  $p_*p^*$  with  $p$  being a projection from  $Y$  to a point. The above yields that our morphism  $\sum \psi_i$  is a retraction of an isomorphism  $\sum \phi_i$  hence is an isomorphism itself.  $\square$

*Remark 8.* It can be shown [P1] that  $K_*(A(\lambda_i))$  depends only on the  $Z(G)$  action on  $V_G(\lambda_i)$ . The explicit description of the arising algebras could be found in [Ti].

## 9. EXAMPLES.

**9.1. Twisted flag variety.**  $K$ -theory of twisted flag varieties was computed in [P1] and our computation gives the same description for the inner forms. Flag variety is a homogeneous variety  $G/P$  with split semisimple  $G$  and parabolic  $P \subset G$ , and this definition includes projective spaces, flag varieties in usual sense (for  $G = SL_n$ ), split projective quadrics, etc.

There is a decomposition  $P = LU$  into the semidirect product of Levi subgroup and unipotent radical of  $P$  hence there is a morphism

$$G/L \longrightarrow G/P$$

with the fiber  $U$ . Over the extinguished point  $P$  acts on  $U$  by

$$l_p u_p \cdot u = l_p u_p u l_p^{-1},$$

so one has the representation of  $P$  on the Lie algebra of  $U$ , and  $G/L$  is the corresponding vector bundle.

Levi subgroup is a reductive subgroup of maximal rank hence theorem 4 gives an explicit answer for the  $K_*(G/L)$  and by the homotopy invariance of  $K$ -theory this is an answer for  $K_*(G/P)$ . Note that the center  $Z(G)$  acts trivially on the above bundle so one can twist it and get a new vector bundle

$$(G/L)_\gamma \longrightarrow (G/P)_\gamma,$$

hence  $K$ -theory for the inner form of a flag variety could be computed by our method as well.

**9.2. Even dimensional affine quadric.** This case corresponds to the inclusion  $SO_{2n} \subset SO_{2n+1}$  and for the root systems it is  $D_n \subset B_n$ .

First of all we pass to the simply connected group,

$$Spin_{2n+1}/Spin_{2n} = SO_{2n+1}/SO_{2n}.$$

One has  $[W(B_n) : W(D_n)] = 2$  so there are two elements in the basis. First element  $V(\lambda_1)$  as usual corresponds to the trivial representation of  $Spin_{2n}$  and as the second we can take one of the half-spin representations  $V(\omega_{n-1}), V(\omega_n)$ , since the algorithm from theorem 2 suggests one of the fundamental weights having the orbit consisting of two points.

After twisting with  $\gamma$  we get a quadric  $X(q)$  defined by a quadratic form  $q$ , then the algebra  $A(\lambda_2)_\gamma = C_0(q)$  is the even Clifford algebra for the form  $q$  [Ti]. Hence one has

$$K_*(X(q)) = K_*(k) \oplus K_*(C_0(q)).$$

This answer coincides with the one obtained in [Sw].

**9.3. Quaternionic projective space.** We consider

$$HP^n = Sp_{2n+2}/(Sp_2 \times Sp_{2n})$$

as an algebraic model for the quaternionic projective space. The motivation comes from the fact that  $HP^n(\mathbb{C})$  is homotopy equivalent to the usual quaternionic projective space  $\mathbb{H}P^n$ . An extensive treatment of the quaternionic flag varieties including the simplest case of projective spaces one can find in [PW].

The root systems in this case are  $C_1 + C_n \subset C_{n+1}$ , so basis consists of

$$[W(C_{n+1}) : (W(C_1) \times W(C_n))] = n + 1$$

elements. We have dealt with this case in section 5.3 and the basis consists of  $\Lambda^i W$  for regular representation  $W$  of  $Sp_{2n}$ . The center acts trivially on the even degrees and nontrivially on the odd ones, so one has

$$K_*(HP_\gamma^n) = K_*(k)^{\lceil \frac{n+1}{2} \rceil} \oplus K_*(A(\lambda_1)_\gamma)^{\lfloor \frac{n+1}{2} \rfloor}.$$

In the split case it reduces to  $K_*(HP^n) = K_*(k)^{n+1}$ , and it agrees with the result obtained in [PW].

**9.4. Zero characteristic.** In this case we can treat non-reductive groups. When  $char k = 0$  one has the Levi decomposition  $G = L_G U_G$  of group  $G$  into the semidirect product of some reductive subgroup and the unipotent radical [Mc], which in general fails in the positive characteristic. Also in this case the unipotent radical  $U_G$  splits, i.e. it has a filtration with vector factors [KMT] so the underlying variety is  $\mathbb{A}^n$ , which also can fail over nonperfect fields. Hence for the connected split groups of the same rank  $H \subset G$  one has the following triangle.

$$\begin{array}{ccc} G/L_H & \xrightarrow{p_1} & G/H \\ & \downarrow p_2 & \\ & L_G/L_H & \end{array}$$

The fibres of  $p_1$  and  $p_2$  are isomorphic to  $U_G$  and  $U_H$  respectively and both are affine spaces. One can show that both the projections define some vector

bundles with the trivial action of the center  $Z(L_G)$ , so one can twist with  $L_G/Z(L_G)$ -cocycle  $\gamma$  and from the homotopy invariance obtain that

$$K_*((G/H)_\gamma) = K_*((L_G/L_H)_\gamma).$$

The last one could be computed using the methods introduced in this paper.

**9.5. Octonionic projective plain.** It could be shown [Ba] that

$$\mathbb{O}\mathbb{P}^2 \cong G(F_4)/Spin(9),$$

where  $G(F_4)$  stands for the compact form of the simple algebraic group with the root system  $F_4$ . We consider as an algebraic model

$$OP^2 = G(F_4)/Spin_9$$

with split  $G(F_4)$ . It corresponds to the root systems  $B_4 \subset F_4$  treated in section 5.2. One has

$$[W(F_4) : W(B_4)] = 3,$$

and the corresponding representations for  $Spin_9$  are  $k, V(\omega_3), V(\omega_4)$ , i.e. the trivial one, 84-dimensional  $\Lambda^3 W$  and 16-dimensional spinor representation. Since the center of  $G(F_4)$  is trivial the twisting does not produce interesting algebras, though it changes the variety. Hence one has

$$K_*(OP^2_\gamma) = K_*(k) \oplus K_*(k) \oplus K_*(k).$$

#### REFERENCES

- [Ba] J. C. Baez, *The Octonions*, Bull. Amer. Math. Soc. 39 (2002), 145-205.
- [BT] A. Borel and J. Tits, *Groupes réductifs*, Inst. Hautes Études Sci. Publ. Math. 27 (1965), 55-151.
- [Bo] N. Bourbaki, *Lie groups and Lie algebras*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002.
- [H1] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Second printing, revised. Grad. Texts in Math., 9. Springer-Verlag, New York, 1978
- [H2] J. E. Humphreys, *Linear Algebraic Groups*, Grad. Texts in Math., 21, Berlin, New York, 1972
- [Ja] J. C. Jantzen, *Representations of algebraic groups*, Pure and Applied Mathematics vol. 131, Academic Press, Orlando, 1987.
- [KMT] T. Kambayashi, M. Miyanishi and M. Takeuchi, *Unipotent Algebraic Groups*, Springer Lecture Notes in Math. 414 (1974)
- [Mc] G. McNinch, *Levi decompositions of a linear algebraic group*, to appear, Transformation Groups, preprint arXiv:1007.2777.
- [M] A. Merkurjev, *Equivariant K-theory*, J. Handbook of K-theory. Vol. 1, 2 (2005), Springer, Berlin, 925-954.
- [P1] I. Panin, *On the algebraic K-theory of twisted flag varieties*, K-Theory 8 (1994), 541-585.
- [P2] I. Panin, *Splitting principle and K-theory of simply connected semisimple algebraic groups*, St. Petersburg Math. J. 10 (1999), 69-101.
- [PW] I. Panin and C. Walter, *Quaternionic Grassmannians and Pontryagin classes in algebraic geometry*, arXiv:1011.0649.
- [Q] D. Quillen, *Higher algebraic K-theory I*, Lect. Notes in Math. 341 (1972), 85-147
- [Sp] T. Springer, *Linear algebraic groups*, 2nd ed., Progr. in Math., vol. 9, Birkhuser, Boston, 1998.
- [St] R. Steinberg, *On a theorem of Pittie*, Topology 14 (1975), 173-177.
- [Sw] R. Swan, *K-theory of quadric hypersurfaces*, Ann. Math., 121 (1985) 113-153.
- [Ti] J. Tits, *Représentations linéaires irréductibles d'un groupe réductif sur un corps quelconque*, J. Reine Angew. Meth. 247 (1971), 196-220.