

VARIATIONS ON A THEME OF GROUPS  
SPLITTING BY A QUADRATIC EXTENSION  
AND GROTHENDIECK-SERRE CONJECTURE FOR  
GROUP SCHEMES  $F_4$  WITH TRIVIAL  $g_3$  INVARIANT

V. CHERNOUSOV<sup>1</sup>

Received: August 17, 2009

Revised: February 1, 2010

ABSTRACT. We study structure properties of reductive group schemes defined over a local ring and splitting over its étale quadratic extension. As an application we prove Serre–Grothendieck conjecture on rationally trivial torsors over a local regular ring containing a field of characteristic 0 for group schemes of type  $F_4$  with trivial  $g_3$  invariant.

2010 Mathematics Subject Classification: 20G07, 20G10, 20G15, 20G41

Keywords and Phrases: Linear algebraic groups, exceptional groups, torsors, non-abelian cohomology, local regular rings, Grothendieck–Serre conjecture

*To A. Suslin on his 60th birthday*

## 1 INTRODUCTION

In the present paper we prove the Grothendieck-Serre conjecture on rationally trivial torsors for group schemes of type  $F_4$  whose generic fiber has trivial  $g_3$  invariant. The Grothendieck-Serre conjecture [Gr58], [Gr68], [S58] asserts that if  $R$  is a regular local ring and if  $G$  is a reductive group scheme defined over  $R$  then a  $G$ -torsor over  $R$  is trivial if and only if its fiber at the generic point of  $\text{Spec}(R)$  is trivial. In other words the kernel of a natural map  $H_{\text{ét}}^1(R, G) \rightarrow H_{\text{ét}}^1(K, G)$  where  $K$  is a quotient field of  $R$  is trivial.

---

<sup>1</sup>Partially supported by the Canada Research Chairs Program and an NSERC research grant

Many people contributed to this conjecture by considering various particular cases. If  $R$  is a discrete valuation ring the conjecture was proved by Y. Nisnevich [N]. If  $R$  contains a field  $k$  and  $G$  is defined over  $k$  this is due to J.-L. Colliot-Thélène, M. Ojanguren [CTO] when  $k$  is infinite perfect and it is due to M. S. Raghunathan [R94], [R95] when  $k$  is infinite. The case of tori was done by J.-L. Colliot-Thélène and J.-L. Sansuc [CTS]. For certain simple simply connected group of classical type the conjecture was proved by Ojanguren, Panin, Suslin and Zainoulline [PS], [OP], [Z], [OPZ]. For a recent progress on isotropic group schemes we refer to preprints [PSV], [Pa09], [PPS].<sup>2</sup>

In the paper we deal with a still open case related to group schemes of type  $F_4$ . Recall that if  $G$  is a group of type  $F_4$  defined over a field  $k$  of characteristic  $\neq 2, 3$  one can associate (cf. [S93], [GMS03], [PetRac], [Ro]) cohomological invariants  $f_3(G)$ ,  $f_5(G)$  and  $g_3(G)$  of  $G$  in  $H^3(k, \mu_2)$ ,  $H^5(k, \mu_2)$  and  $H^3(k, \mathbb{Z}/3\mathbb{Z})$  respectively. The group  $G$  can be viewed as the automorphism group of a corresponding 27-dimensional Jordan algebra  $J$ . The invariant  $g_3(G)$  vanishes if and only if  $J$  is reduced, i.e. it has zero divisors. The main result of the paper is the following.

**THEOREM 1.** *Let  $R$  be a regular local ring containing a field of characteristic 0. Let  $G$  be a group scheme of type  $F_4$  over  $R$  such that its fiber at the generic point of  $\text{Spec}(R)$  has trivial  $g_3$  invariant. Then the canonical mapping  $H_{\text{ét}}^1(R, G) \rightarrow H_{\text{ét}}^1(K, G)$  where  $K$  is a quotient field of  $R$  has trivial kernel.*

We remark that for a group scheme  $G$  of type  $F_4$  we have  $\text{Aut}(G) \simeq G$ , so that by the twisting argument the above theorem is equivalent to the following:

**THEOREM 2.** *Let  $R$  be as above and let  $G_0$  be a split group scheme of type  $F_4$  over  $R$ . Let  $H_{\text{ét}}^1(R, G_0)_{\{g_3=0\}} \subset H_{\text{ét}}^1(R, G_0)$  be the subset consisting of isomorphism classes  $[\mathcal{T}]$  of  $G_0$ -torsors such that the corresponding twisted group  $({}^{\mathcal{T}}G_0)_K$  has trivial  $g_3$  invariant. Then a canonical mapping*

$$H_{\text{ét}}^1(R, G_0)_{\{g_3=0\}} \rightarrow H_{\text{ét}}^1(K, G_0)$$

*is injective, i.e. two  $G_0$ -torsors in  $H_{\text{ét}}^1(R, G_0)_{\{g_3=0\}}$  are isomorphic over  $R$  if and only if they are isomorphic over  $K$ .*

The characteristic restriction in the theorem is due to the fact that the purity result [ChP] is used in the proof and the latter is based on the use of the main result in [P09] on rationally isotropic quadratic spaces which was proven in characteristic zero only (the resolution of singularities is involved in that proof). We remark that if the Panin's result is true in full generality (except probably characteristic 2 case) then our arguments can be easily modify in such way that the theorem holds for all regular local rings where 2 is invertible.<sup>3</sup>

<sup>2</sup>We also remark that experts know the proof of the conjecture for group schemes of type  $G_2$  but it seems to us that a proof is not available in the literature.

<sup>3</sup>I. Panin has informed the author that his main theorem in [P09] holds for quadratic spaces defined over a regular local ring containing an infinite perfect field.

The proof of the theorem heavily depends on the fact that group schemes of type  $F_4$  with trivial  $g_3$  invariant are split by an étale quadratic extension of the ground ring  $R$ . This is why the main body of the paper consists of studying structure properties of simple group schemes of an arbitrary type over  $R$  (resp.  $K$ ) splitting by an étale quadratic extension  $S/R$  (resp.  $L/K$ ) which is of independent interest.

We show that the structure of such group schemes is completely determined by a finite family of units in  $R$  which we call structure constants of  $G$ . These constants depend on a chosen maximal torus  $T \subset G$  defined over  $R$  and splitting over  $S$ . Such a torus is not unique in  $G$ . Giving two tori  $T$  and  $T'$  we find formulas which express structure constants of  $G$  related to  $T$  in terms of that of related to  $T'$  and this leads us quickly to the proof of the main theorem.

Of course we are using a group point view. It seems plausible that our proof can be carried over in terms of Jordan algebras and their trace quadratic forms, but we do not try to do it here.

The paper is divided into four parts. We begin by introducing notation, terminology that are used throughout the paper as well as by reminding properties of algebraic groups defined over a field and splitting by a quadratic field extension. This is followed by two sections on explicit formulas for cohomological invariants  $f_3$  and  $f_5$  in terms of structure constants for groups of type  $F_4$  and their classification. In the third part of the paper we study structure properties of group schemes splitting by an étale quadratic extension of the ground ring. The proof of the main theorem is the content of the last section.

NOTATION. Let  $R$  be a (commutative) ring. We let  $G_0$  denote a split reductive group scheme over  $R$  and we let  $T_0 \subset G_0$  denote a maximal split torus over  $R$ . We denote by  $\Sigma(G_0, T_0)$  the root system of  $G_0$  with respect to  $T_0$ . We use standard terminology related to algebraic groups over rings. For the definition of reductive group schemes (and in particular split reductive group schemes), maximal tori, root systems of split group schemes and their properties we refer to [SGA3].

We number the simple roots as in [Bourb68].

*Acknowledgments.* We thank the referee for useful comments and remarks which helped to improve the exposition.

## 2 LEMMA ON REPRESENTABILITY OF UNITS BY QUADRATIC FORMS

Throughout the paper  $R$  denotes a (commutative) ring where 2 is invertible and  $R^\times$  denotes the group of invertible elements of  $R$ . Also, all fields considered in the paper have characteristic  $\neq 2$ .

If  $R$  is a local ring with the maximal ideal  $M$  we let  $k = \overline{R} = R/M$ . Similarly, if  $V$  is a free module on rank  $n$  over  $R$  we let  $\overline{V} = V \otimes_R \overline{R} = V \otimes_R k$  and for a vector  $v \in V$  we set  $\overline{v} = v \otimes 1$ . If  $R$  is a regular local ring it is a unique factorization domain ([Ma, Theorem 48, page 142]). Throughout the paper a quotient field of  $R$  will be denoted by  $K$ .

Let  $f = \sum_{i=1}^n a_i x_i^2$  be a quadratic form over  $R$  where  $a_1, \dots, a_n \in R^\times$  given on a free  $R$ -module  $V$ . If  $I \subset \{1, \dots, n\}$  is a non-empty subset we denote by  $f_I = \sum_{i \in I} a_i x_i^2$  the corresponding subform of  $f$ . If  $v = (v_1, \dots, v_n) \in V$  we set  $f_I(v) = \sum_{i \in I} a_i v_i^2$ . Finally, let  $g = \prod_I f_i$  where the product is taken over all non-empty subsets of  $\{1, \dots, n\}$ . For a vector  $v$  we set  $g(v) = \prod_I f_I(v)$ .

LEMMA 3. *Let  $f$  and  $g$  be as above. Assume that (the residue field)  $k$  is infinite. Let  $a \in R^\times$  be a unit such that  $f(v) = a$  for some vector  $v \in V$ . Then there exists a vector  $u \in V$  such that  $f(u) = a$  and  $g(u)$  is a unit.*

*Proof.* If  $n = 1$ ,  $v$  has the required properties. Hence we may assume  $n \geq 2$ . If  $w \in V$  is a vector whose length  $f(w)$  with respect to  $f$  is a unit we denote by  $\tau_w$  an orthogonal reflection with respect to  $w$  given by

$$\tau_w(x) = x - 2f(x, w)f(w)^{-1}w$$

for all vectors  $x$  in  $V$ . Since orthogonal reflections preserve length of vectors it suffices to find vectors  $w_1, \dots, w_s \in V$  such that  $g(\tau_{w_1} \cdots \tau_{w_s}(v))$  is a unit. For that, in turn, it suffices to find  $\bar{w}_1, \dots, \bar{w}_s \in \bar{V}$  such that  $\bar{g}(\tau_{\bar{w}_1} \cdots \tau_{\bar{w}_s}(\bar{v})) \neq 0$ . It follows that we can pass to a vector space  $\bar{V}$  over  $k$ . Consider a quadric

$$Q_{\bar{a}} = \{x \in \bar{V} \mid \bar{f}(x) = \bar{a}\}$$

defined over  $k$ . We have  $\bar{v} \in Q_{\bar{a}}(k)$ , hence  $Q_{\bar{a}}(k) \neq \emptyset$  implying  $Q_{\bar{a}}$  is a rational variety over  $k$ .

Let  $U \subset \bar{V}$  be an open subset given by  $\bar{g}(x) \neq 0$ . It is easy to see that  $Q_{\bar{a}} \cap U \neq \emptyset$  (indeed, if we pass to an algebraic closure  $\bar{k}$  of  $k$  then obviously we have  $U(\bar{k}) \cap Q_{\bar{a}}(\bar{k}) \neq \emptyset$ ). Since  $k$  is infinite,  $k$ -points of  $Q_{\bar{a}}$  are dense in  $Q_{\bar{a}}$ . Hence  $Q_{\bar{a}}(k) \cap U$  is nonempty. Take a vector  $\bar{w} \in Q_{\bar{a}}(k) \cap U$ . Since the orthogonal group  $O(\bar{f})$  acts transitively on vectors of  $Q_{\bar{a}}$  there exists  $\bar{\tau} \in O(\bar{f})$  such that  $\bar{w} = \bar{\tau}(\bar{v})$ . It remains to note that orthogonal reflections generate  $O(\bar{f})$ .  $\square$

### 3 ALGEBRAIC GROUPS SPLITTING BY QUADRATIC FIELD EXTENSIONS

The aim of this section is to remind structure properties of a simple simply connected algebraic group  $G$  defined over a field  $K$  and splitting over its quadratic extension  $L/K$ . There is nothing special in type  $F_4$  and we will assume in this section that  $G$  is of an arbitrary type of rank  $n$ . The only technical restriction which we need later on to simplify the exposition of the material on the structure of such groups relates to the Weyl group  $W$  of  $G$ . Namely, we will assume that  $W$  contains  $-1$ , i.e. an element which takes an arbitrary root  $\alpha$  into  $-\alpha$ .<sup>4</sup> Let  $\tau$  be the nontrivial automorphism of  $L/K$ . If  $B_L \subset G_L$  is a Borel subgroup over  $L$  in  $G_L$  in generic position then  $B_L \cap \tau(B_L) = T$  is a maximal torus in

<sup>4</sup>For groups  $G$  splitting over a quadratic extension of the ground field and whose whose Weyl group doesn't contain  $-1$  the Galois descent data looks more complicated; for instance, Lemma 4 doesn't hold for them.

$G_L$ . Clearly, it is defined over  $K$  and splitting over  $L$  (because it is contained in  $B_L$  and all tori in  $B_L$  are  $L$ -split).

LEMMA 4.  *$T$  is anisotropic over  $K$ .*

*Proof.* The Galois group of  $L/K$  acts in a natural way on characters of  $T$  and hence on the root system  $\Sigma = \Sigma(G_K, T)$  of  $G_K$  with respect to  $T_K$ . Thus we have a natural embedding  $\text{Gal}(L/F) \hookrightarrow W$  which allows us to view  $\tau$  as an element of  $W$ . Since the intersection of two Borel subgroups  $B_L$  and  $\tau(B_L)$  is a maximal torus in  $G_L$ , one of them, say  $\tau(B_L)$ , is the opposite Borel subgroup to the second one  $B_L$  with respect to the ordering on  $\Sigma$  determined by the pair  $(T_L, B_L)$ . One knows that  $W$  contains a unique element which takes  $B_L$  to  $\tau(B_L) = B_L^-$ . Since  $-1 \in W$  such an element is necessary  $-1$ . Of course this implies  $\tau = -1$ , hence  $\tau$  acts on characters of  $T$  as  $-1$ . In particular  $T$  is  $K$ -anisotropic.  $\square$

Our Borel subgroup  $B_L$  determines an ordering of the root system  $\Sigma$  of  $G_L$ , hence the system of simple roots  $\Pi = \{\alpha_1, \dots, \alpha_n\}$ . Let  $\Sigma^+$  (resp.  $\Sigma^-$ ) be the set of positive (resp. negative) roots. Let us choose a Chevalley basis [St]

$$\{H_{\alpha_1}, \dots, H_{\alpha_n}, X_\alpha, \alpha \in \Sigma\} \quad (5)$$

in the Lie algebra  $\mathfrak{g}_L = \mathcal{L}(G_L)$  of  $G_L$  corresponding to the pair  $(T_L, B_L)$ . Recall that elements from (5) are eigenvectors of  $T_L$  with respect to the adjoint representation  $ad : G \rightarrow \text{End}(\mathfrak{g}_L)$  satisfying some additional relations; in particular for each  $t \in T_L$  we have

$$tX_\alpha t^{-1} = \alpha(t)X_\alpha \quad (6)$$

where  $\alpha \in \Sigma$  and  $tH_{\alpha_i} t^{-1} = H_{\alpha_i}$ . A Chevalley basis is unique up to signs and automorphisms of  $\mathfrak{g}_L$  which preserve  $B_L$  and  $T_L$  (see [St], §1, Remark 1).

Since  $G_L$  is a Chevalley group over  $L$ , the structure of  $G(L)$  as an abstract group, i.e. its generators and relations, is well known. For more details and proofs of all standard facts about  $G(L)$  used in this paper we refer to [St]. Recall that  $G(L)$  is generated by the so-called root subgroups  $U_\alpha = \langle x_\alpha(u) \mid u \in L \rangle$ , where  $\alpha \in \Sigma$  and  $T$  is generated by the one-parameter subgroups

$$T_\alpha = T \cap G_\alpha = \text{Im } h_\alpha$$

Here  $G_\alpha$  is the subgroup generated by  $U_{\pm\alpha}$  and  $h_\alpha : G_{m,L} \rightarrow T_L$  is the corresponding cocharacter (coroot) of  $T$ . Furthermore, since  $G_L$  is a simply connected group, the following relations hold in  $G_L$  (cf. [St], Lemma 28 b), Lemma 20 c):

- (i)  $T \simeq T_{\alpha_1} \times \dots \times T_{\alpha_n}$ ;
- (ii) for any two roots  $\alpha, \beta \in \Sigma$  and  $t, u \in L$  we have

$$h_\alpha(t) x_\beta(u) h_\alpha(t)^{-1} = x_\beta(t^{\langle \beta, \alpha \rangle} u)$$

where  $\langle \beta, \alpha \rangle = 2(\beta, \alpha)/(\alpha, \alpha)$  and

$$h_\alpha(t)X_\beta h_\alpha(t)^{-1} = t^{\langle \beta, \alpha \rangle} X_\beta \quad (7)$$

If  $\Delta \subset \Sigma^+$  is a subset, we let  $G_\Delta$  denote the subgroup generated by  $U_{\pm\alpha}$ ,  $\alpha \in \Delta$ . We shall now describe explicitly the  $K$ -structure of  $G$ , i.e. the action of  $\tau$  on the generators  $\{x_\alpha(u), \alpha \in \Sigma\}$  of  $G_L$ . As we already know  $\tau(\alpha) = -\alpha$  for any  $\alpha \in \Sigma$  and this implies  $T_\alpha \simeq R_{L/K}^{(1)}(G_{m,L})$  (see [V, 4.9, Example 6]).

Let  $\alpha \in \Sigma$ . Since  $\tau(\alpha) = -\alpha$  there exists a constant  $c_\alpha \in L^\times$  such that  $\tau(X_\alpha) = c_\alpha X_{-\alpha}$ . It follows that the action of  $\tau$  on  $G(L)$  is determined completely by the family  $\{c_\alpha, \alpha \in \Sigma\}$ . We call these constants by *structure constants* of  $G$  with respect to  $T$  and Chevalley basis (5). Of course, they depend on the choice of  $T$  and a Chevalley basis. We summarize their properties in the following two lemmas (for their proofs we refer to [Ch, Lemmas 4.4, 4.5, 4.11]).

LEMMA 8. *Let  $\alpha \in \Sigma$ . Then we have*

- (i)  $c_{-\alpha} = c_\alpha^{-1}$ ;
- (ii)  $c_\alpha \in K^\times$ ;
- (iii) *if  $\beta \in \Sigma$  is a root such that  $\alpha + \beta \in \Sigma$ , then  $c_{\alpha+\beta} = -c_\alpha c_\beta$ ; in particular, the family  $\{c_\alpha, \alpha \in \Sigma\}$  is determined completely by its subfamily  $\{c_{\alpha_1}, \dots, c_{\alpha_n}\}$ .*

LEMMA 9. (i)  $\tau[x_\alpha(u)] = x_{-\alpha}(c_\alpha \tau(u))$  for every  $u \in L$  and every  $\alpha \in \Sigma$ .

(ii) *Let  $L = K(\sqrt{d})$ . Then the subgroup  $G_\alpha$  of  $G$  is isomorphic to  $\mathrm{SL}(1, D)$  where  $D$  is a quaternion algebra over  $K$  of the form  $D = (d, c_\alpha)$ .*

#### 4 MOVING TORI

We follow the notation of the previous section. The family  $\{c_\alpha, \alpha \in \Sigma\}$  determining the action of  $\tau$  on  $G(L)$  depends on a chosen Borel subgroup  $B_L$  and the corresponding Chevalley basis. Given another Borel subgroup and Chevalley basis we get another family of constants and we now are going to describe the relation between the old ones and the new ones.

Let  $B'_L \subset G_L$  be a Borel subgroup over  $L$  such that the intersection  $T' = B'_L \cap \tau(B'_L)$  is a maximal  $K$ -anisotropic torus. Clearly both tori  $T$  and  $T'$  are isomorphic over  $K$  (because both of them are isomorphic to the direct product of  $n$  copies of  $R_{L/K}^{(1)}(G_{m,L})$ ). Furthermore, there exists a  $K$ -isomorphism  $\lambda : T \rightarrow T'$  preserving positive roots, i.e. which takes  $(\Sigma')^+ = \Sigma(G, T')^+$  into  $\Sigma^+ = \Sigma(G, T)^+$ . Any such isomorphism can be extended to an inner automorphism

$$i_g : G \longrightarrow G, \quad x \rightarrow g x g^{-1}$$

for some  $g \in G(K_s)$ , where  $K_s$  is a separable closure of  $K$ , which takes  $B_L$  into  $B'_L$  (see [Hum], Theorem 32.1). Note that  $g$  is not unique since for any  $t \in T(K_s)$  the inner conjugation by  $gt$  also extends  $\lambda$  and it takes  $B_L$  into  $B'_L$ .

LEMMA 10. *The element  $g$  can be chosen in  $G(L)$ .*

*Proof.* Take an arbitrary  $g' \in G(K_s)$  such that  $i_{g'}$  extends  $\lambda$  and  $i_{g'}(B_L) = B'_L$ . Since the restriction  $i_{g'}|_T$  is a  $K$ -defined isomorphism, we have

$$t_\sigma = (g')^{-1+\sigma} \in T(K_s)$$

for any  $\sigma \in \text{Gal}(K_s/K)$ . The family  $\{t_\sigma, \sigma \in \text{Gal}(K_s/F)\}$  determines a cocycle  $\xi = (t_\sigma) \in Z^1(K, T)$ . Since  $T$  splits over  $L$ ,  $\text{res}_L(\xi)$  viewed as a cocycle in  $T$  is trivial, by Hilbert’s Theorem 90. It follows there is  $z \in T(K_s)$  such that  $t_\sigma = z^{1-\sigma}$ ,  $\sigma \in \text{Gal}(K_s/L)$ . Then  $g = g'z$  is stable under  $\text{Gal}(K_s/L)$ . This implies  $g \in G(L)$  and clearly we have  $gB_Lg^{-1} = B'_L$ .  $\square$

Let  $g$  be an element from Lemma 10 and let  $t = g^{-1+\tau}$ . Since  $t \in T(L)$ , it can be written uniquely as a product  $t = h_{\alpha_1}(t_1) \cdots h_{\alpha_n}(t_n)$ , where  $t_1, \dots, t_n \in L^\times$  are some parameters.

LEMMA 11. *We have  $t_1, \dots, t_n \in K^\times$ .*

*Proof.* We first note that, by the construction of  $t$ , we have  $t\tau(t) = 1$ . Since  $\tau$  acts on characters of  $T$  as multiplication by  $-1$  we have  $\tau(h_{\alpha_i}(t_i)) = h_{\alpha_i}(1/\tau(t_i))$  for every  $i = 1, \dots, n$ . Also, the equality  $t\tau(t) = 1$  implies  $h_{\alpha_i}(t_i)h_{\alpha_i}(1/\tau(t_i)) = 1$ , hence  $t_i = \tau(t_i)$ .  $\square$

The set

$$\{H'_{\alpha_1} = gH_{\alpha_1}g^{-1}, \dots, H'_{\alpha_n} = gH_{\alpha_n}g^{-1}, X'_\alpha = gX_\alpha g^{-1}, \alpha \in \Sigma\} \tag{12}$$

is a Chevalley basis related to the pair  $(T', B'_L)$ . Let  $\{c'_\alpha, \alpha \in \Sigma\}$  be the corresponding structure constants of  $G$  with respect to  $T'$  and Chevalley basis (12).

LEMMA 13. *For every root  $\alpha \in \Sigma'$  one has  $c'_\alpha = t_1^{-\langle \alpha, \alpha_1 \rangle} \cdots t_n^{-\langle \alpha, \alpha_n \rangle} \cdot c_\alpha$ .*

*Proof.* Apply  $\tau$  to the equality  $X'_\alpha = gX_\alpha g^{-1}$  and use relation (7).  $\square$

Our element  $g$  constructed in Lemma 10 has the property  $g^{-1+\tau} \in T(L)$ . Conversely, it is easy to see that an arbitrary  $g \in G(L)$  with this property gives rise to a new pair  $(B'_L, T')$  and hence to the new structure constants  $\{c'_\alpha\}$  which are given by the formulas in Lemma 13. Thus we have

LEMMA 14. *Let  $g \in G(L)$  be an element such that  $t = g^{-1+\tau} \in T(L)$ . Then  $T' = gTg^{-1}$  is a  $K$ -defined maximal torus splitting over  $L$  and the restriction of the inner automorphism  $i_g$  to  $T$  is a  $K$ -defined isomorphism. The structure constants  $\{c'_\alpha\}$  related to  $T'$  are given by the formulas in Lemma 13.*

EXAMPLE 15. Let  $G, T$  be as above and let  $\Sigma = \Sigma(G, T)$ . Take an element

$$g = x_{-\alpha}(-c_\alpha v)x_\alpha \left( \frac{-\tau(v)}{1 - c_\alpha v\tau(v)} \right) \tag{16}$$

where  $\alpha \in \Sigma$  is an arbitrary root and  $v \in L^\times$  is such that  $1 - c_\alpha v \tau(v) \neq 0$ . One easily checks that

$$g^{-1+\tau} = h_\alpha \left( \frac{1}{1 - c_\alpha v \tau(v)} \right)$$

and hence  $g$  gives rise to a new torus  $T' = gTg^{-1}$  and to a new structure constants.

DEFINITION 17. We say that we apply an elementary transformation of  $T$  with respect to a root  $\alpha$  and a parameter  $v \in L^\times$  when we move from  $T$  to  $T' = gTg^{-1}$  where  $g$  is given by (16) and  $1 - c_\alpha v \tau(v) \neq 0$ .

REMARK 18. The main property of an elementary transformation with respect to a root  $\alpha$  is that the new structure constant  $c'_\beta$  with respect to  $T'$  doesn't change (up to squares) if  $\beta$  is orthogonal to  $\alpha$  or  $\langle \beta, \alpha \rangle = \pm 2$  and it is equal to  $(1 - c_\alpha v \tau(v))c_\beta$  (up to squares) if  $\langle \beta, \alpha \rangle = \pm 1$ . Thus in the context of algebraic groups this is an analogue of an elementary chain equivalence of quadratic forms.

REMARK 19. An arbitrary reduced norm in the quaternion algebra  $D = (d, c_\alpha)$  can be written as a product of two elements of the form  $1 - c_\alpha v \tau(v)$ , hence in the case  $\langle \beta, \alpha \rangle = \pm 1$  we can change  $c_\beta$  by any reduced norm in  $D$ .

## 5 COHOMOLOGICAL INTERPRETATION

While considering cohomological invariants of  $G$  of type  $F_4$  sometimes it is convenient to consider  $G$  as a twisting group. Let  $G^{ad}$  be the corresponding adjoint group. Note that groups of type  $F_4$  are simply connected and adjoint so that for them we have  $G = G^{ad}$ . Let  $G_0$  (resp.  $G_0^{ad}$ ) be a  $K$ -split simple simply connected (resp. adjoint) group of the same type as  $G^{ad}$  and let  $T_0 \subset G_0$  (resp.  $T_0^{ad} \subset G_0^{ad}$ ) be a maximal  $K$ -split torus. We denote by  $c \in \text{Aut}(G_0)$  an element such that  $c^2 = 1$  and  $c(t) = t^{-1}$  for every  $t \in T_0$  (it is known that such an automorphism exists, see e.g. [DG], Exp. XXIV, Prop. 3.16.2, p. 355). We assume additionally that  $c \in N_{G_0^{ad}}(T_0^{ad})$ .

REMARK 20. In general case  $c$  can not be lifted to  $N_{G_0}(T_0)$ . However it is known that if  $G_0$  has type  $D_4$  or  $F_4$  such an element can be chosen inside the normalizer  $N_{G_0}(T_0)$  of  $T_0$ . So when we deal with such groups we will assume that  $c \in N_{G_0}(T_0)$ .

LEMMA 21. Let  $t \in T_0^{ad}(K)$  and let  $a_\tau = ct$ . Then  $\xi = (a_\tau)$  is a cocycle in  $Z^1(L/K, G_0^{ad}(L))$ .

*Proof.* We need to check that  $a_\tau \tau(a_\tau) = 1$ . Indeed,

$$a_\tau \tau(a_\tau) = ct \tau(ct) = ctct = t^{-1}t = 1$$

as required. □



For further reference we note that every cocycle  $\eta \in Z^1(K, G_0^{ad})$  acts by inner conjugation on both  $G_0$  and  $G_0^{ad}$  and hence we can twist  ${}^\eta G_0, {}^\eta G_0^{ad}$  both groups. Since  $G_0^{ad}$  is adjoint the character group of  $T_0^{ad}$  is generated by simple roots  $\{\alpha_1, \dots, \alpha_n\}$  of the root system  $\Sigma = \Sigma(G_0^{ad}, T_0^{ad})$  of  $G_0^{ad}$  with respect to  $T_0^{ad}$ . Choose a decomposition  $T_0^{ad} = G_m \times \dots \times G_m$  such that the canonical embeddings  $\pi_i : G_m \rightarrow T_0^{ad}$  onto the  $i$ th factor,  $i = 1, \dots, n$ , are the cocharacters dual to  $\alpha_1, \dots, \alpha_n$ .

PROPOSITION 22. *Let  $G$  be as above with structure constants  $c_{\alpha_1}, \dots, c_{\alpha_n}$ . Let  $\xi = (a_\tau)$  where  $a_\tau = ct$  and  $t = \prod_i \pi_i(c_{\alpha_i})$ . Then the twisted group  ${}^\xi G_0$  is isomorphic to  $G$  over  $K$ .*

*Proof.* It is known that  $cX_\alpha c^{-1} = X_{-\alpha}$  and according to (6) we have  $tX_\alpha t^{-1} = \alpha(t)X_\alpha$  for every root  $\alpha \in \Sigma$ . Since the cocharacters  $\pi_1, \dots, \pi_n$  are dual to the roots  $\alpha_1, \dots, \alpha_n$ , we have  $\langle \pi_i, \alpha_j \rangle = \delta_{ij}$ , hence

$$\pi_i(c_{\alpha_i})X_{\alpha_i}\pi_i(c_{\alpha_i})^{-1} = c_{\alpha_i}X_{\alpha_i}$$

and

$$\pi_i(c_{\alpha_i})X_{\alpha_j}\pi_i(c_{\alpha_i})^{-1} = X_{\alpha_j}$$

if  $i \neq j$ . Thus for the twisted group  ${}^\xi G_0$  the structure constant for the simple root  $\alpha_i, i = 1, \dots, n$ , is  $c_{\alpha_i}$  because

$$X_{\alpha_i} \rightarrow a_\tau X_{\alpha_i} a_\tau^{-1} = (c \prod_i \pi_i(c_{\alpha_i}))X_{\alpha_i}(c \prod_i \pi_i(c_{\alpha_i}))^{-1} = c_{\alpha_i}X_{-\alpha_i}.$$

If  $\alpha \in \Sigma$  is an arbitrary root, then by Lemma 8 the structure constant  $c_\alpha$  of  ${}^\xi G_0$  can be expressed uniquely in terms of the constants  $c_{\alpha_1}, \dots, c_{\alpha_n}$ , so that the twisted group  ${}^\xi G_0$  has the same structure constants as  $G$ . It follows that the Lie algebras  $\mathcal{L}(G)$  and  $\mathcal{L}({}^\xi G_0)$  of  $G$  and  ${}^\xi G_0$  have the same Galois descent data. This yields  $\mathcal{L}(G) \simeq \mathcal{L}({}^\xi G_0)$  and as a consequence we obtain that their automorphism groups (and in particular their connected components) are isomorphic over  $K$  as well.  $\square$

REMARK 23. Assume that  $R$  is a domain where 2 is invertible with a field of fractions  $K$  and  $G_0$  is a split group scheme over  $R$ . Let  $S = R(\sqrt{d})$  be an étale quadratic extension of  $R$  where  $d$  is a unit in  $R$ . Let  $\tau$  be the generator of  $\text{Gal}(S/R)$ . Assume that  $c_{\alpha_1}, \dots, c_{\alpha_n} \in R^\times$ . Then we may view  $\xi = (a_\tau)$  where  $a_\tau = c \prod_i \pi_i(c_{\alpha_i})$  as a cocycle in  $Z^1(S/R, G_0^{ad}(S))$  and hence the twisted group  ${}^\xi G_0$  is a group scheme over  $R$  whose fiber at the generic point of  $\text{Spec}(R)$  is isomorphic to  $G_K$ .

As an application of the above proposition we get

LEMMA 24. *Let  $G$  and  $G'$  be groups over  $K$  and splitting over  $L$  with structure constants  $\{c_{\alpha_1}, \dots, c_{\alpha_n}\}$  and  $\{c_{\alpha_1}u_1, \dots, c_{\alpha_n}u_n\}$  where  $u_1, \dots, u_n$  are in the image of the norm map  $N_{L/K} : L^\times \rightarrow K^\times$ . Then  $G$  and  $G'$  are isomorphic over  $K$ .*

*Proof.* Let  $u_i = N_{L/K}(v_i)$ . By Proposition 22, we have  $G$  and  $G'$  are twisted forms of  $G_0$  by means of cocycles  $\xi = (a_\tau)$  and  $\xi' = (a'_\tau)$  with coefficients in  $G_0^{ad}(S)$  where  $a_\tau = c \prod_i \pi_i(c_{\alpha_i})$  and  $a'_\tau = c \prod_i \pi_i(c_{\alpha_i} u_i)$ . Since  $T_0^{ad}$  is a  $K$ -split torus and since  $\pi_i$  is a  $K$ -defined morphism we have  $\tau(\pi_i(v_i)) = \pi_i(\tau(v_i))$ . Also, we have  $c^2 = 1$  and  $c\pi_i(v_i)c^{-1} = \pi_i(v_i^{-1})$ . Then it easily follows

$$a_\tau = \left( \prod_i \pi_i(v_i) \right) a'_\tau \left( \prod_i \pi_i(v_i) \right)^{-\tau}$$

and this implies  $\xi$  is equivalent to  $\xi'$ . □

The statement of the lemma can be equivalently reformulated as follows.

**COROLLARY 25.** *Let  $T \subset G$  be a maximal torus with the structure constants  $\{c_{\alpha_1}, \dots, c_{\alpha_n}\}$  and let  $u_1, \dots, u_n \in N_{L/K}(L^\times)$ . Then  $G$  contains a maximal torus  $T'$  whose structure constants are  $\{c_{\alpha_1} u_1, \dots, c_{\alpha_n} u_n\}$ .*

6 STRONGLY INNER FORMS OF TYPE  $D_4$

For later use we need some classification results on strongly inner forms of type  ${}^1D_4$ ; in other words we need an explicit description of the image of  $H^1(K, G_0) \rightarrow H^1(K, \text{Aut}(G_0))$  where  $G_0$  is a simple simply connected group over a field  $K$  of type  $D_4$ .

For an arbitrary cocycle  $\xi \in Z^1(K, G_0)$  the twisted group  $G = {}^\xi G_0$  is isomorphic to  $\text{Spin}(f)$  where  $f$  is an 8-dimensional quadratic form having trivial discriminant and trivial Hasse-Witt invariant. By Merkurjev's theorem [M],  $f$  belongs to  $I^3$  where  $I$  is the fundamental ideal of even dimensional quadratic forms in the Witt group  $W(K)$ . We may assume that  $f$  represents 1 (because  $\text{Spin}(f) \simeq \text{Spin}(af)$  for  $a \in K^\times$ ). Since  $\dim f = 8$ , by the Arason-Pfister Hauptsatz,  $f$  is a 3-fold Pfister form over  $K$  and as a consequence we obtain  $G$  is splitting over a quadratic extension  $L/K$  of  $K$ , say  $L = K(\sqrt{d})$ .

**LEMMA 26.** *There exist parameters  $u_1, \dots, u_4 \in K^\times$  such that  $G \simeq {}^\eta G_0$  where  $\eta$  is of the form  $\eta = (a_\tau)$  and  $a_\tau = c \prod_i h_{\alpha_i}(u_i)$ .*

*Proof.* By Remark 20 we may assume that  $c \in N_{G_0}(T_0)$ . Let  $\xi'$  be the image of  $\xi$  in  $H^1(K, G_0^{ad})$  and let  $c'$  be the image of  $c$  in  $G_0^{ad}$ . By Proposition 22, we may assume that  $\xi'$  is of the form  $\xi' = (a'_\tau)$  where  $a'_\tau = c' \prod_i \pi_i(c_{\alpha_i})$  and  $c_{\alpha_i}$  are structure constants of  $G^{ad} = {}^{\xi'} G_0^{ad}$  with respect to some maximal torus in  $G^{ad}$  defined over  $K$  and splitting over  $L$ .

The element  $c$  gives rise to a cocycle  $\lambda = (b_\tau) \in Z^1(L/K, G_0(L))$  where  $b_\tau = c$ . Twisting  $G_0$  by  $\lambda$  yields a commutative diagram

$$\begin{array}{ccc} H^1(K, G_0) & \xrightarrow{f_1} & H^1(K, {}^\lambda G_0) \\ \downarrow & & \downarrow \\ H^1(K, G_0^{ad}) & \xrightarrow{f_2} & H^1(K, {}^\lambda G_0) \end{array}$$

where  $f_1$  and  $f_2$  are the canonical bijections. Let  $f_2(\xi') = \xi''$ . It is of the form  $\xi'' = (a''_\tau)$  where  $a''_\tau = \prod_i \pi_i(c_{\alpha_i})$ ; hence  $f_2(\xi')$  takes values in a maximal torus  $T^{ad} = {}^\lambda T_0^{ad}$  of  ${}^\lambda(G_0^{ad})$  defined over  $K$  and splitting over  $L$ .

Let  $Z$  be the center of  $G_0$ . We have an exact sequence

$$0 \rightarrow Z \rightarrow {}^\lambda T_0 \rightarrow T^{ad} \rightarrow 1$$

It induces a morphism  $f_3 : H^1(K, T^{ad}) \rightarrow H^2(K, Z)$ . Since  $c'$  and  $\xi'$  can be lifted to  $G_0$ , we have  $f_3(\xi'') = 0$ . Hence  $\xi''$  has a lifting into the torus  ${}^\lambda T_0$ , say  $\tilde{\eta} \in H^1(L/K, {}^\lambda T_0)$ . Going back to  $H^1(K, T_0)$  we see that  $\eta = f_1^{-1}(\tilde{\eta})$  has the required property.  $\square$

Since we are interesting in the description of  $G = {}^\xi G_0$  we may assume without loss of generality that  $\xi = \eta$ . It is known that  $Z \simeq \mu_2 \times \mu_2$  (see [PR94, §6.5]), hence  $Z$  contains three elements of order 2. They give rise to three homomorphisms  $\phi_i : G_0 \rightarrow \text{SO}(f_0)$  where  $i = 1, 2, 3$  and  $f_0$  is a split 8-dimensional quadratic form. The images  $\phi_i(\xi)$ ,  $i = 1, 2, 3$ , of  $\xi$  in  $Z^1(K, \text{SO}(f_0))$  correspond to three quadratic form  $f_1, f_2, f_3$  and we are going to give an explicit description of  $f_i$  in terms of the parameters  $u_1, u_2, u_3, u_4$  and  $d$ .

LEMMA 27. *Up to numbering we have  $f_1 = u_3 f$ ,  $f_2 = u_4 f$  and  $f_3 = u_3 u_4 f$  where  $f = \langle\langle d, v_1, v_2 \rangle\rangle$  and  $v_1 = u_1 u_3^{-1} u_4^{-1}$ ,  $v_2 = u_2$ . In particular  $G$  is split over a field extension  $E/K$  if and only if so is  $f_E$ .*

*Proof.* One easily checks that  $Z$  is generated by

$$h_{\alpha_1}(-1)h_{\alpha_3}(-1) \quad \text{and} \quad h_{\alpha_1}(-1)h_{\alpha_4}(-1).$$

We now rewrite the cocycle  $\xi = (a_\tau)$  in the form

$$a_\tau = ch_{\alpha_1}(v_1)h_{\alpha_2}(v_2)z_1z_2$$

where  $v_1 = u_1 u_3^{-1} u_4^{-1}$ ,  $v_2 = u_2$  and

$$z_1 = h_{\alpha_1}(u_3)h_{\alpha_3}(u_3), \quad z_2 = h_{\alpha_1}(u_4)h_{\alpha_4}(u_4).$$

Using relation (7) we find that the structure constants of  $G$  with respect to the twisted torus  $T = {}^\xi T_0$  up to squares are  $c_{\alpha_2} = v_1$  and  $c_{\alpha_1} = c_{\alpha_3} = c_{\alpha_4} = v_2$ . Also, applying the same twisting argument as in [ChS, 4.1] we find that up to numbering we have  $f_1 = u_3 f$ ,  $f_2 = u_4 f$  and  $f_3 = u_3 u_4 f$  where

$$f = \langle\langle d, v_1, v_2 \rangle\rangle = \langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle.$$

$\square$

We are now going to show that we don't change the equivalence class  $[\xi]$  if we multiply the parameters  $u_3, u_4$  in the expression for  $\xi$  by elements in  $K^\times$  represented by  $f$ . Let  $V, V_1, V_2, V_3$  be 8-dimensional vector space over  $K$  equipped with the quadratic forms  $f, f_1, f_2, f_3$ .

PROPOSITION 28. *Let  $w_1, w_2 \in V$  be two anisotropic vectors and let  $a = f(w_1), b = f(w_2)$ . Let  $\xi' = (a'_\tau)$  where  $a'_\tau = ch_{\alpha_1}(v_1)h_{\alpha_2}(v_2)z'_1z'_2$  and*

$$z'_1 = h_{\alpha_1}(au_3)h_{\alpha_3}(au_3), \quad z'_2 = h_{\alpha_1}(bu_4)h_{\alpha_4}(bu_4).$$

*Then  $\xi'$  is equivalent to  $\xi$ .*

*Proof.* Consider two embeddings  $\psi_1 \psi_2 : \mu_2 \rightarrow G_0$  given by

$$-1 \rightarrow h_{\alpha_1}(-1)h_{\alpha_3}(-1)$$

and

$$-1 \rightarrow h_{\alpha_1}(-1)h_{\alpha_4}(-1).$$

Up to numbering we may assume that

$${}^\xi G_0/\psi_1(\mu_2) \simeq \text{SO}(f_1) \quad \text{and} \quad {}^\xi G_0/\psi_2(\mu_2) \simeq \text{SO}(f_2).$$

We also have a canonical bijection  $H^1(K, G_0) \rightarrow H^1(K, {}^\xi G_0)$  (translation by  $\xi$ ) under which  $\xi'$  goes to  $\eta = (h_{\alpha_1}(a)h_{\alpha_3}(a)h_{\alpha_1}(b)h_{\alpha_4}(b))$  and we need to show that  $\eta$  is trivial in  $H^1(K, {}^\xi G_0)$ .

We now note that  $\eta$  is the product of two cocycles  $\eta_1 = (h_{\alpha_1}(a)h_{\alpha_3}(a))$  and  $\eta_2 = (h_{\alpha_1}(b)h_{\alpha_4}(b))$  first of which being in the image of  $\psi_1^* : H^1(K, \mu_2) \rightarrow H^1(K, {}^\xi G_0)$  induced by  $\psi_1$  and the second one being in the image of  $\psi_2^* : H^1(K, \mu_2) \rightarrow H^1(K, {}^\xi G_0)$  induced by  $\psi_2$ . We may identify  $H^1(K, \mu_2) = K^\times/(K^\times)^2$ . It is known that  $\text{Ker } \psi_1^*$  (resp.  $\text{Ker } \psi_2^*$ ) consists of spinor norms of  $f_1$  (resp.  $f_2$ ). Thus the statement of the proposition is amount to saying that  $a, b$  are spinor norms for the twisted group  $G = {}^\xi G_0$  with respect to the quadratic forms  $f_1$  and  $f_2$  respectively. Since spinor norms of  $f_i$  are generated by  $f_i(s_1)f_i(s_2)$  where  $s_1, s_2 \in V_i$  are anisotropic vectors and since  $f_i$  is proportional to  $f$  we are done.  $\square$

REMARK 29. Assume that  $R$  and  $S$  are as in Remark 23. Take a cocycle  $\xi = (a_\tau)$  in  $Z^1(S/R, G_0(S))$  given by  $a_\tau = ch_{\alpha_1}(u_1) \cdots h_{\alpha_4}(u_4)$  where  $u_1, \dots, u_4 \in R^\times$ . Then arguing literally verbatim we find that the twisted group  $G = {}^\xi G_0$  is isomorphic to  $\text{Spin}(f)$  where  $f$  is a 3-fold Pfister form given by  $f = \langle\langle d, u_2, u_1u_3u_4 \rangle\rangle$  and that for all units  $a, b \in R^\times$  represented by  $f$  the cocycle  $\xi'$  from Proposition 28 is equivalent to  $\xi$ .

PROPOSITION 30. *Let  $G$  be as above and let  $f = \langle\langle d, v_1, v_2 \rangle\rangle$  be the corresponding 3-fold Pfister form. Assume that  $f$  has another presentation  $f = \langle\langle d, a, b \rangle\rangle$  over  $K$ . Then there exists a maximal torus  $T' \subset G$  defined over  $K$  and splitting over  $L$  such that structure constants of  $G$  with respect to  $T'$  (up to squares) are  $c'_{\alpha_1} = a$  and  $c'_{\alpha_2} = b$ .*

*Proof.* We proved in Lemma 27 that the structure constants of  $G$  with respect to the torus  $T = {}^\xi T_0$  are  $c_{\alpha_1} = v_2$  and  $c_{\alpha_1} = v_1$ . We now construct a sequence of elementary transformations of  $T$  with respect to the roots  $\alpha_1$  and  $\alpha_2$  such that

at the end we arrive to a torus with the required structure constants. Recall that, by Remarks 18 and 19, an application of an elementary transformation of  $T$  with respect to  $\alpha_1$  (resp.  $\alpha_2$ ) does not change  $c_{\alpha_1}$  (resp.  $c_{\alpha_2}$ ) modulo squares and multiplies  $c_{\alpha_2}$  (resp.  $c_{\alpha_1}$ ) by a reduced norm from the quaternion algebra  $(d, c_{\alpha_1})$  (resp.  $(d, c_{\alpha_2})$ ).

By Witt cancellation we may write  $a$  in the form  $a = w_1 c_{\alpha_1} + w_2 c_{\alpha_2} - w_3 c_{\alpha_1} c_{\alpha_2}$  where  $w_1, w_2, w_3 \in N_{L/K}(L^\times)$ . By Corollary 25, passing to another maximal torus and Chevalley basis (if necessary) we may assume without loss of generality that  $w_1 = w_2 = 1$  and hence we may assume that  $a$  is of the form  $a = c_{\alpha_1}(1 - w_3 c_{\alpha_2}) + c_{\alpha_2}$  where  $w_3$  is still in  $N_{L/K}(L^\times)$ .

If  $1 - w_3 c_{\alpha_2} = 0$  then  $a = c_{\alpha_2}$  and we pass to the last paragraph of the proof. Otherwise applying a proper elementary transformation with respect to  $\alpha_2$  we pass to a new torus with structure constants  $c'_{\alpha_1} = c_{\alpha_1}(1 - w_3 c_{\alpha_2})$  and  $c'_{\alpha_2} = c_{\alpha_2}$ . Thus abusing notation without loss of generality we may assume

$$a = c_{\alpha_1} + c_{\alpha_2} = c_{\alpha_1}(1 - (-c_{\alpha_1})^{-1}c_{\alpha_2}).$$

Applying again a proper elementary transformation with respect to  $\alpha_1$  we can pass to a torus whose second structure constant is  $(-c_{\alpha_1})^{-1}c_{\alpha_2}$ , so that we may assume  $a = c_{\alpha_1}(1 - c_{\alpha_2})$ . Lastly, applying an elementary transformation with respect to  $\alpha_2$  we pass to a torus such that  $a = c_{\alpha_1}$ .

We finally observe that from

$$\langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle = \langle\langle d, a, b \rangle\rangle = \langle\langle d, c_{\alpha_1}, b \rangle\rangle$$

it follows that  $b$  is of the form  $b = w c_{\alpha_2}$  where  $w \in \text{Nrd}(d, c_{\alpha_1})$ . So a proper elementary transformation with respect to  $\alpha_1$  completes the proof.  $\square$

## 7 ALTERNATIVE FORMULAS FOR $f_3$ AND $f_5$ INVARIANTS

We are going to apply the previous technique to produce explicit formulas for the  $f_3$  and  $f_5$  invariants of a group  $G$  of type  $F_4$  over a field  $K$  of characteristic  $\neq 2$  with trivial  $g_3$  invariant. Recall (cf. [S93], [GMS03], [PetRac]) that given such  $G$  one can associate the cohomological invariants  $f_3(G) \in H^3(K, \mu_2)$  and  $f_5(G) \in H^5(K, \mu_2)$  with the following properties (cf. [Sp], [Ra]):

- (a) *The group  $G$  is split over a field extension  $E/K$  if and only if  $f_3(G)$  is trivial over  $E$ ;*
- (b) *The group  $G$  is isotropic over a field extension  $E/K$  if and only if  $f_5(G)$  is trivial over  $E$ .*

These two invariants  $f_3, f_5$  are symbols given in terms of the trace quadratic form of the Jordan algebra  $J$  corresponding to  $G$  and hence we may associate to them 3-fold and 5-fold Pfister forms. Abusing notation we denote them by the same symbols  $f_3(G)$  and  $f_5(G)$ . It is well known that  $f_3(G)$  and  $f_5(G)$  completely classify groups of type  $F_4$  with trivial  $g_3$  invariant (see [Sp], [S93])

and we would like to produce explicit formulas of  $f_3(G)$  and  $f_5(G)$  in group terms only in order to generalize them later on to the case of local rings.

It follows from (a) that our group  $G$  is splitting by a quadratic extension. Indeed, if  $f_3(G) = (d) \cup (a) \cup (b)$  then passing to  $L = K(\sqrt{d})$  we get  $G_L$  has trivial  $f_3$  invariant and as a consequence  $G$  is  $L$ -split by property (a).

We next construct a subgroup  $H$  in  $G$  of type  $D_4$  and compute structure constants of  $G$  and  $H$ . By Proposition 22 we may view  $G$  as a twisted group  ${}^\xi G_0$  where  $\xi = (a_\tau)$ ,  $a_\tau = c \prod_{i=1}^4 h_{\alpha_i}(u_i)$  and  $u_1, \dots, u_4 \in K^\times$  where  $G_0$  is a split group of type  $F_4$ . Looking at the tables in [Bourb68] we find that the subroot system  $\Sigma'$  in  $\Sigma(G_0, T_0)$  generated by the long roots has type  $D_4$ . One checks that

$$\beta_1 = -\epsilon_1 - \epsilon_2, \quad \beta_2 = \alpha_1, \quad \beta_3 = \alpha_2, \quad \beta_4 = \epsilon_3 + \epsilon_4$$

is its basis. Since  $\epsilon_3 + \epsilon_4 = \alpha_2 + 2\alpha_3$  and  $\epsilon_1 + \epsilon_2 = 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 2\alpha_4$ , it follows that the cocharacters  $h_{\epsilon_3+\epsilon_4}$  and  $h_{\epsilon_1+\epsilon_2}$  are equal to

$$h_{\epsilon_3+\epsilon_4} = h_{\alpha_2} + h_{\alpha_3} \quad \text{and} \quad h_{\epsilon_1+\epsilon_2} = 2h_{\alpha_1} + 3h_{\alpha_2} + 2h_{\alpha_3} + h_{\alpha_4}$$

so that

$$h_{\epsilon_3+\epsilon_4}(u) = h_{\alpha_2}(u)h_{\alpha_3}(u) \tag{31}$$

and

$$h_{\epsilon_1+\epsilon_2}(u) = h_{\alpha_1}(u^2)h_{\alpha_2}(u^3)h_{\alpha_3}(u^2)h_{\alpha_4}(u) \tag{32}$$

for all parameters  $u \in L^\times$ .

These relations shows that  $a_\tau$  can be rewritten in the form

$$a_\tau = ch_{\alpha_1}(v_1)h_{\alpha_2}(v_2)[h_{\epsilon_1+\epsilon_2}(v_3)h_{\alpha_2}(v_3)][h_{\epsilon_3+\epsilon_4}(v_4)h_{\alpha_2}(v_4)] \tag{33}$$

where  $v_1, v_2, v_3, v_4 \in K^\times$ .

Let  $H_0$  be the subgroup in  $G_0$  generated by  $\Sigma'$ . It is stable with respect to the conjugation by  $a_\tau$ , hence  $G$  contains the subgroup  $H = {}^\xi H_0$  of type  $D_4$ . Using (7) we easily find that modulo squares in  $K^\times$  one has  $c_{\alpha_3} = v_2v_3$  and  $c_{\alpha_4} = v_4$  and  $c_{\alpha_1} = v_2, c_{\alpha_2} = v_1$ ; in particular  $c_{\alpha_1}, c_{\alpha_2}$  don't depend on  $v_3, v_4$  modulo squares.

Recall that two  $n$ -fold Pfister forms, say  $g_1$  and  $g_2$ , are isomorphic over the ground field  $K$  if and only if  $g_1$  is hyperbolic over the function field of  $K(g_2)$  of  $g_2$ .

**THEOREM 34.** *One has  $f_3(G) = (d) \cup (c_{\alpha_1}) \cup (c_{\alpha_2})$ .*

*Proof.* Let  $f = \langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle$  and let  $E$  be the function field of  $f$ . According to property (a), it suffices to show that  $G$  is split over  $E$  or  $H$  is split over  $E$ . But in Lemma 27 we showed that  $H \simeq \text{Spin}(f)$  and so we are done.  $\square$

The following proposition shows that the structure constants  $c_{\alpha_3}$  are  $c_{\alpha_4}$  of  $G$  are well defined modulo values of  $f = f_3(G) = \langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle$ .

PROPOSITION 35. *Let  $a, b \in K^\times$  be represented by  $f$  over  $K$ . Then there exists a maximal torus  $T' \subset G$  defined over  $K$  and splitting over  $L$  such that modulo squares  $G$  has structure constants  $c_{\alpha_1}, c_{\alpha_2}, ac_{\alpha_3}, bc_{\alpha_4}$  with respect to  $T'$ .*

*Proof.* According to Proposition 28, if multiply the parameters  $v_3, v_4$  in the expression (33) by  $a, b$  respectively we obtain a cocycle equivalent to  $\xi$ , so the result follows.  $\square$

THEOREM 36. *One has  $f_5(G) = (d) \cup (c_{\alpha_1}) \cup (c_{\alpha_2}) \cup (c_{\alpha_3}) \cup (c_{\alpha_4})$ .*

*Proof.* Let  $g = \langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c_{\alpha_3}, c_{\alpha_4} \rangle\rangle$ . Arguing as in Theorem 34 and using property (b) we may assume that  $g$  is split and we have to prove that  $G$  is isotropic. Since  $g$  is split we may write  $c_{\alpha_4}$  in the form

$$c_{\alpha_4} = a^{-1}(1 - bc_{\alpha_3}) \quad (37)$$

where  $a, b$  are represented by  $f = \langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle$ . Our aim is to pass (with the use of elementary transformations) to a new torus  $T' \subset G$  defined over  $K$  and splitting over  $L$  such that the new structure constant  $c'_{\alpha_4}$  related to  $T'$  is equal to 1 modulo squares. The last would imply that the corresponding subgroup  $G_{\alpha_4}$  of  $G$  is isomorphic to  $\mathrm{SL}_2$  by Lemma 9 (ii) and this would show that  $G$  is isotropic as required.

By Proposition 35 there exists a maximal torus  $T'$  in  $G$  such that two last structure constants related to  $T'$  are  $c'_{\alpha_3} = bc_{\alpha_3}$  and  $c'_{\alpha_4} = ac_{\alpha_4}$ . Then by (37) we have  $c'_{\alpha_4} = 1 - c'_{\alpha_3}$ . Applying a proper elementary transformation with respect to  $\alpha_3$  we pass to the third torus  $T''$  for which  $c''_{\alpha_4} = 1$  modulo squares and we are done.  $\square$

## 8 CLASSIFICATION OF GROUPS OF TYPE $F_4$ WITH TRIVIAL $g_3$ INVARIANT

The theorem below is due to T. Springer [Sp]. In this section we produce an alternative short proof which can be easily adjusted to the case of local rings.

THEOREM 38. *Let  $G_0$  be a split group of type  $F_4$  over a field  $K$ . A mapping*

$$H_{\acute{e}t}^1(K, G_0)_{\{g_3=0\}} \rightarrow H^3(K, \mu_2) \times H^5(K, \mu_2)$$

*given by  $G \rightarrow (f_3(G), f_5(G))$  is injective.*

We need the following preliminary result.

PROPOSITION 39. *Let  $G$  be a group of type  $F_4$  defined over  $K$  and splitting over  $L$  with structure constants  $c_{\alpha_1}, \dots, c_{\alpha_4}$  with respect to a torus  $T$ . Let  $a \in K^\times$  be represented by  $g = \langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c_{\alpha_3} \rangle\rangle$  over  $K$ . Then there is a maximal torus  $T' \subset G$  such that the corresponding structure constants are  $c_{\alpha_1}, c_{\alpha_2}, c_{\alpha_3}, ac_{\alpha_4}$  modulo squares.*

*Proof.* Write  $a$  in the form  $a = a_1(1 - a_2c_{\alpha_3})$  where  $a_1, a_2$  are represented by  $f = \langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle$ . By Proposition 35 the structure constants  $c_{\alpha_3}$  and  $c_{\alpha_4}$  are well defined modulo values of  $f$ . Hence passing to another maximal torus in  $G$  we may assume without loss of generality that  $a_1 = a_2 = 1$  so that  $a = 1 - c_{\alpha_3}$ . Since  $1 - c_{\alpha_3}$  is a reduced norm in the quaternion algebra  $(d, c_{\alpha_3})$  a proper elementary transformation with respect to  $\alpha_3$  lead us to a torus whose first three structure constants are the same modulo squares and the last one is  $(1 - c_{\alpha_3})c_{\alpha_4}$ .  $\square$

*Proof of Theorem 38.* Let  $G, G'$  be two groups of type  $F_4$  over  $K$  such that  $f_3(G) = f_3(G')$  and  $f_5(G) = f_5(G')$ . Choose a quadratic extension  $L/K$  splitting  $f_3(G)$ . It splits both  $G$  and  $G'$ . Our strategy is to show that  $G, G'$  contain maximal tori defined over  $K$  and splitting over  $L$  with the same structure constants.

Choose arbitrary maximal tori  $T \subset G, T' \subset G'$  defined over  $K$  and splitting over  $L$ . Let  $c_{\alpha_1}, \dots, c_{\alpha_4}$  and  $c'_{\alpha_1}, \dots, c'_{\alpha_4}$  be the corresponding structure constants. As we know,  $G, G'$  contain subgroups  $H, H'$  of type  $D_4$  over  $K$  generated by the long roots. By Theorem 34 we have  $f_3(G) = (d) \cup (c_{\alpha_1}) \cup (c_{\alpha_2})$  and  $f_3(G') = (d) \cup (c'_{\alpha_1}) \cup (c'_{\alpha_2})$ , hence

$$\langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle = \langle\langle d, c'_{\alpha_1}, c'_{\alpha_2} \rangle\rangle.$$

Then according to Proposition 30 applied to  $H'$  and  $f = \langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle$  we may assume without loss of generality that  $c_{\alpha_1} = c'_{\alpha_1}$  and  $c_{\alpha_2} = c'_{\alpha_2}$ .

We next show that up to choice of maximal tori in  $G$  and  $G'$  we also may assume that  $c_{\alpha_3} = c'_{\alpha_3}$ . Since  $f_5(G) = f_5(G')$  we get

$$\langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c_{\alpha_3}, c_{\alpha_4} \rangle\rangle = \langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c'_{\alpha_3}, c'_{\alpha_4} \rangle\rangle. \quad (40)$$

By Witt cancellation we can write  $c'_{\alpha_3}$  in the form  $c'_{\alpha_3} = a_1c_{\alpha_3} + a_2c_{\alpha_4} - a_3c_{\alpha_3}c_{\alpha_4}$  where  $a_1, a_2, a_3$  are values of  $f$ . By Proposition 35 we may assume without loss of generality that  $a_1 = a_2 = 1$ . Arguing as in Proposition 30 we may pass to another maximal torus in  $G'$  such that the corresponding structure constants are

$$c'_{\alpha_1} = c_{\alpha_1}, \quad c'_{\alpha_2} = c_{\alpha_2}, \quad c'_{\alpha_3} = c_{\alpha_3}.$$

Finally, from (40) it follows that  $c'_{\alpha_4} = ac_{\alpha_4}$  for some  $a \in K^\times$  represented by  $g = \langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c_{\alpha_3} \rangle\rangle$ . Application of Proposition 39 completes the proof.  $\square$

## 9 GROUP SCHEMES SPLITTING BY ÉTALE QUADRATIC EXTENSIONS

We now pass to a simple simply connected group scheme  $G$  of an arbitrary type of rank  $n$  defined over a local ring  $R$  where 2 is invertible and splitting by an étale quadratic extension  $S = R(\sqrt{u}) \simeq R[t]/(t^2 - u)$  of  $R$  where  $u \in R^\times$ . We assume that  $R$  is a domain with a quotient field  $K$  and with a residue field  $k$  and we assume  $u$  is not square in  $K^\times$ . We also denote  $L = S \otimes_R K$  and



$l = S \otimes_R k$ . Abusing notation we denote the nontrivial automorphisms of  $S/R$ ,  $L/K$  and  $l/k$  by the same letter  $\tau$ .

Let  $\mathfrak{g}$  be the Lie algebra of  $G$ . As usual we set

$$\mathfrak{g}_S = \mathfrak{g} \otimes_R S, \quad \mathfrak{g}_K = \mathfrak{g} \otimes_R K, \quad \mathfrak{g}_L = \mathfrak{g} \otimes_R L$$

and

$$\bar{\mathfrak{g}} = \mathfrak{g}_k = \mathfrak{g} \otimes_R k, \quad \bar{\mathfrak{g}}_S = \mathfrak{g}_l = \mathfrak{g}_S \otimes_S l.$$

Let  $\mathfrak{b}_S$  be a Borel subalgebra in  $\mathfrak{g}_S$ . We say that it is in a *generic position* if  $\bar{\mathfrak{b}}_S \cap \tau(\bar{\mathfrak{b}}_S)$  is a Cartan subalgebra in  $\bar{\mathfrak{g}}_l$ . This amounts to saying that  $\bar{\mathfrak{b}}_S \cap \tau(\bar{\mathfrak{b}}_S)$  has dimension  $n$  over  $l$ .

We will systematically use below the fact that in a split simple Lie algebra defined over a field the intersection of two Borel subalgebras contains a split Cartan subalgebra; in particular this intersection has dimension at least  $n$ .

LEMMA 41. *The Lie algebra  $\mathfrak{g}_S$  contains Borel subalgebras in generic position.*

*Proof.* Let  $\mathcal{B}$  and  $\bar{\mathcal{B}}$  be the varieties of Borel subalgebras in the split Lie algebras  $\mathfrak{g}_S$  and  $\bar{\mathfrak{g}}_l$  respectively. Passing to residues we have a canonical mapping  $\mathcal{B} \rightarrow \bar{\mathcal{B}}$  whose image is dense (because  $\mathfrak{g}_S$  is split). Let  $U \subset \bar{\mathcal{B}}$  be an open subset in Zariski topology consisting of Borel subalgebras  $\mathfrak{b}_l$  such that  $\mathfrak{b}_l \cap \tau(\mathfrak{b}_l)$  has dimension  $n$ . Since  $\mathcal{B}(S)$  is dense in  $\mathcal{B}$  there exists a Borel subalgebra  $\mathfrak{b}_S$  in  $\mathfrak{g}_S$  over  $S$  whose image in  $\bar{\mathcal{B}}$  is contained in  $U$ .  $\square$

LEMMA 42. *Let  $\mathfrak{b}_S \subset \mathfrak{g}_S$  be a Borel subalgebra in generic position. Then a submodule  $\mathfrak{t}_S = \mathfrak{b}_S \cap \tau(\mathfrak{b}_S)$  of  $\mathfrak{b}_S$  has rank  $n$ .*

*Proof.* Let  $M_S \subset S$  be a maximal ideal. Our subalgebra  $\mathfrak{t}_S$  is given as an intersection of two free submodules in  $\mathfrak{g}_S$  of codimensions  $m$ , where  $m$  is the number of positive roots in  $\mathfrak{g}_S$ , each of them being a direct summand in  $\mathfrak{g}_S$ . So  $\mathfrak{t}_S$  consists of all solutions of a linear system of  $m$  equations in  $m+n$  variables. The space of solutions of this system modulo  $M$  coincides with the intersection  $\bar{\mathfrak{b}}_S \cap \tau(\bar{\mathfrak{b}}_S)$  and hence it has dimension  $n$ . This implies that the linear system has a minor of size  $m \times m$  whose determinant is a unit in  $S$  and we are done.  $\square$

Our next aim is to show that the Galois descent data for the generic fiber  $G_K$  of  $G$  described in previous sections can be pushed down at the level of  $R$ . As usual we will assume that the Weyl group of  $G$  contains  $-1$ .

PROPOSITION 43. *Let  $\mathfrak{b}_S \subset \mathfrak{g}_S$  be a Borel subalgebra in generic position and let  $\mathfrak{t}_S = \mathfrak{b}_S \cap \tau(\mathfrak{b}_S)$ . Then  $\mathfrak{t}_S$  is a split Cartan subalgebra of  $\mathfrak{g}_S$  contained in  $\mathfrak{b}_S$ .*

*Proof.* Let  $\mathfrak{u}_S$  be the ideal in  $\mathfrak{b}_S$  consisting of nilpotent elements. It is complemented in  $\mathfrak{b}_S$  by a split Cartan algebra and hence  $\mathfrak{b}_S/\mathfrak{u}_S$  is isomorphic to a split Cartan subalgebra in  $\mathfrak{b}_S$ . We want to show that a canonical projection  $p: \mathfrak{b}_S \rightarrow \mathfrak{b}_S/\mathfrak{u}_S$  restricted at  $\mathfrak{t}_S$  is an isomorphism.

Let  $\mathfrak{b}_L = \mathfrak{b}_S \otimes_S L$  be a generic fiber of  $\mathfrak{b}_S$ . We already know that  $\mathfrak{t}_L = \mathfrak{b}_L \cap \tau(\mathfrak{b}_L)$  has dimension  $n$  over  $L$ , so it is a split Cartan algebra in  $\mathfrak{g}_L$ . Since  $\mathfrak{t}_S$  embeds into  $\mathfrak{t}_L$ , it is a commutative Lie subalgebra contained in  $\mathfrak{b}_S$  and consisting of diagonalizable semisimple elements. So injectivity of  $p$  follows.

As for surjectivity, it suffices to prove it modulo maximal ideal  $M_R \subset R$ . In the course of proving of Lemma 42 we saw that  $\mathfrak{t}_S$  is the space of solutions of the linear system of  $m$  equations in  $m+n$  variables whose matrix modulo  $M$  has rank  $m$ . It follows  $\mathfrak{t}_S$  modulo  $M$  has dimension  $n$  and we are done.  $\square$

Let now  $\mathfrak{t}_S$  be as in Proposition 43 and let  $\mathfrak{t} = \mathfrak{t}_S^{(\tau)}$  be the invariant subspace. By descent we have  $\mathfrak{t} \otimes_R S = \mathfrak{t}_S$ , hence  $\mathfrak{t}$  is an  $R$ -defined Cartan subalgebra splitting over  $S$ . Let  $B_S$  be a Borel subgroup in  $G_S$  corresponding to  $\mathfrak{b}_S$ . The connected component of the automorphism group of a pair  $(\mathfrak{b}_S, \mathfrak{t}_S)$  gives rise to a maximal torus  $T_S$  in  $B_S$ . It is  $R$ -defined and  $S$ -split because so is  $\mathfrak{t}$ . Let us choose a Chevalley basis

$$\{H_{\alpha_1}, \dots, H_{\alpha_n}, X_{\alpha}, \alpha \in \Sigma\}$$

in  $\mathfrak{g}_S$  corresponding to  $(T_S, B_S)$ . Since  $W$  contains  $-1$ , we know that  $\tau$  acts on the root system  $\Sigma = \Sigma(G_S, T_S)$  as  $-1$ . Now repeating verbatim the arguments in [Ch] we easily find that for every root  $\alpha \in \Sigma$  there exists a constants  $c_{\alpha} \in R$  such that  $\tau(X_{\alpha}) = c_{\alpha} X_{-\alpha}$  and hence the action of  $\tau$  on  $G(S)$  is determined completely by the family  $\{c_{\alpha}, \alpha \in \Sigma\}$ . We call these constants by *structure constants* of  $G$  with respect to  $T$ .

As in [Ch] one checks that the structure constants satisfy the relations given in Lemmas 8, 9. Also, as in Example 15 we may obviously define the notion of an elementary transformation with respect to a root  $\alpha \in \Sigma$  (because root subgroups  $U_{\alpha}$  are defined over  $S$ ).

REMARK 44. We note that the structure constants  $\{c_{\alpha} \mid \alpha \in \Sigma\}$  are units in  $R$ . Indeed, by our construction we have surjections  $\mathfrak{b}_S \rightarrow \overline{\mathfrak{b}}_S$  and  $\mathfrak{b}_S \cap \tau(\mathfrak{b}_S) \rightarrow \overline{\mathfrak{b}}_S \cap \tau(\overline{\mathfrak{b}}_S)$ . Hence the residues of  $c_{\alpha}$  are structure constants of  $\overline{G} = G \otimes_R k$  in the corresponding basis.

## 10 PROOF OF THEOREM 2

Let  $R$  be a ring satisfying all hypothesis in Theorem 2. As usual we denote its quotient field by  $K$ . Let  $G_0$  be a split group of type  $F_4$  over  $R$  and let  $[\xi] \in H^1(R, G_0)_{\{g_3=0\}}$ . We first claim that the twisted group  $G = {}^{\xi}G_0$  is split by an étale quadratic extension of  $R$ . The proof is based on the following.

LEMMA 45. *There exist  $u, v, w \in R^{\times}$  such that  $f_3(G_K) = (u) \cup (v) \cup (w)$ .*

*Proof.* Let  $f_3(G_K) = (a) \cup (b) \cup (c)$  where  $a, b, c \in K^{\times}$ . By [ChP] the functor of 3-fold Pfister forms satisfies purity, hence it suffices to show that  $f_3(G)$  is unramified at prime ideals of  $R$  of height 1.

Let  $\mathfrak{p} \subset R$  be a prime ideal of height 1 and let  $v = v_{\mathfrak{p}}$  be the corresponding discrete valuation on  $K$  with the residue field  $k(v) = R/\mathfrak{p}$ . We need to show that the image of  $f_3(G_K)$  under the boundary map  $\partial_{v,K} : H^3(K, \mathbb{Z}/2) \rightarrow H^2(k(v), \mathbb{Z}/2)$  is trivial. Consider the following commutative diagram:

$$\begin{array}{ccccccc}
 H^1(R, G_0) & \xrightarrow{\phi_1} & H^1(K, G_0) & \xrightarrow{\mathcal{R}_{G_0, K}} & H^3(K, \mu_6^{\otimes 2}) & \xrightarrow{\partial_{v, K}} & H^2(k(v), \mu_6) \\
 \downarrow & & \downarrow & & \downarrow & & = \downarrow \\
 H^1(R_v, G_0) & \xrightarrow{\phi_2} & H^1(K_v, G_0) & \xrightarrow{\mathcal{R}_{G_0, K_v}} & H^3(K_v, \mu_6^{\otimes 2}) & \xrightarrow{\partial_{v, K_v}} & H^2(k(v), \mu_6)
 \end{array}$$

Here  $\mathcal{R}_{G_0}$  is the Rost invariant for  $G_0$  (see [GMS03]). Since  $g_3(G_K) = 0$  and since  $G_K = (\xi G_0)_K$  we have  $f_3(G_K) = \mathcal{R}_{G_0, K}(\phi_1(\xi))$ . By [G00, Theorem 2], we also have  $\partial_{v, K_v} \circ \mathcal{R}_{G_0, K_v} \circ \phi_2 = 0$ . This yields immediately  $(\partial_{v, K} \circ \mathcal{R}_{G_0, K} \circ \phi_1)(\xi) = 0$  as required.  $\square$

PROPOSITION 46.  $G$  is split by an étale quadratic extension of  $R$ .

*Proof.* By Lemma 45 we have  $f_3(G_K) = (u) \cup (v) \cup (w)$  where  $u, v, w \in R^\times$ . Take  $S = R(\sqrt{u})$  and we claim  $G_S$  is split. One of the following two cases occurs.

If  $u \in (K^\times)^2$  then we have  $f_3(G_K) = 0$ . It follows  $\mathcal{R}_{G_0}([\xi_K]) = f_3(G_K) = 0$ . Since the kernel of the Rost invariant for split groups of type  $F_4$  defined over  $K$  is trivial we have  $[\xi_K] = 0$ . Since by [CTO], [R94], [R95] Grothendieck–Serre conjecture holds for  $G_0$  we conclude  $\xi = 0$ , i.e.  $G$  is already split over  $R$ .

Assume now that  $u \notin (K^\times)^2$ . Let  $L$  be a quotient field of  $S$ . Arguing along the same lines we first get  $\mathcal{R}_{G_0}([\xi_L]) = 0$  and then  $G_S$  is split.  $\square$

The following lemma is an  $R$ -analogue of Corollary 25.

LEMMA 47. Let  $T \subset G$  be a maximal torus with the structure constants  $\{c_{\alpha_1}, \dots, c_{\alpha_4}\}$  and let  $u_1, \dots, u_4 \in N_{S/R}(S^\times)$ . Then  $G$  contains a maximal torus  $T'$  whose structure constants are  $\{c_{\alpha_1}u_1, \dots, c_{\alpha_n}u_4\}$ .

*Proof.* Apply the same argument as in Lemma 24 with the use of Remark 23.  $\square$

*Proof of Theorem 2.* Let  $[\xi], [\xi'] \in H^1(R, G_0)_{\{g_3=0\}}$  be two classes and let  $G, G'$  be the corresponding twisted group schemes over  $R$ . Assume that the generic fibers  $G_K, G'_K$  of  $G$  and  $G'$  are isomorphic over  $K$ . If  $G_K$  is  $K$ -split, there is nothing to prove, because Grothendieck-Serre conjecture is already proven for  $G_0$ , and so we may assume that  $G_K, G'_K$  are not split over  $K$  (and hence  $G, G'$  are not split over  $R$ ) which amounts to saying that  $f_3(G_K) \neq 0$  and  $f_3(G'_K) \neq 0$ .

By Proposition 46 there exists an étale quadratic extension  $S = R(\sqrt{d})$ , where  $d \in R^\times$ , splitting  $G$ . Of course, it is split  $G'$  as well. It now suffices to show

that  $G, G'$  contain maximal tori  $T, T'$  defined over  $R$  and splitting over  $S$  and such that the corresponding structure constants for  $G$  and  $G'$  are the same. Let  $T, T'$  be arbitrary  $R$ -defined and  $S$ -splitting maximal tori in  $G, G'$ . Let  $c_{\alpha_1}, \dots, c_{\alpha_4}$  and  $c'_{\alpha_1}, \dots, c'_{\alpha_4}$  be structure constants of  $G, G'$  with respect to  $T$  and  $T'$ . By Theorem 34 we have  $f_3(G_K) = (d) \cup (c_{\alpha_1}) \cup (c_{\alpha_2})$  and  $f_3(G'_K) = (d) \cup (c'_{\alpha_1}) \cup (c'_{\alpha_2})$ . Since  $f_3(G_K) = f_3(G'_K)$  we get

$$\langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle_K \overset{K}{\simeq} \langle\langle d, c'_{\alpha_1}, c'_{\alpha_2} \rangle\rangle_K$$

and hence

$$\langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle \overset{R}{\simeq} \langle\langle d, c'_{\alpha_1}, c'_{\alpha_2} \rangle\rangle.$$

We first claim that up to choice of  $T$  and  $T'$  we may assume that  $c_{\alpha_1} = c'_{\alpha_1}$  and  $c_{\alpha_2} = c'_{\alpha_2}$ . The proof of the claim is completely similar to that of Proposition 30. Namely, by Witt cancellation and by Lemma 3 we may write  $c'_{\alpha_1}$  in the form  $c'_{\alpha_1} = w_1 c_{\alpha_1} + w_2 c_{\alpha_2} - w_3 c_{\alpha_1} c_{\alpha_2}$  where  $w_1, w_2, w_3 \in N_{S/R}(S^\times)$  and  $w_1 c_{\alpha_1} - w_3 c_{\alpha_1} c_{\alpha_2}$  is a unit in  $R$ . By Lemma 47, passing to another maximal torus in  $G$  (if necessary) we may assume that  $w_1 = w_2 = 1$  and then  $c'_{\alpha_1} = c_{\alpha_1}(1 - w_3 c_{\alpha_2}) + c_{\alpha_2}$  where  $w_3$  is still in  $N_{S/R}(S^\times)$  and  $1 - w_3 c_{\alpha_2}$  is a unit in  $R$ . The rest of the proof is the same as in Proposition 30.

We next claim that up to choice of  $T$  and  $T'$  we may additionally assume that  $c_{\alpha_3} = c'_{\alpha_3}$ . To prove it we are just copying the related part of the proof of Theorem 38. Arguing as in Proposition 22 we conclude that up to equivalence  $\xi$  and  $\xi'$  are of the form  $\xi = (a_\tau)$  and  $\xi' = (a'_\tau)$  where  $a_\tau = c \prod_{i=1}^n h_{\alpha_i}(u_i)$  and  $a'_\tau = c \prod_{i=1}^n h_{\alpha_i}(u'_i)$ , so that, by Remark 29,  $G$  and  $G'$  contain simple simply connected subgroups  $H$  and  $H'$  generated by long roots such that  $H \simeq H' \simeq \text{Spin}(f)$  where  $f = \langle\langle d, c_{\alpha_1}, c_{\alpha_2} \rangle\rangle$ . Furthermore arguing as in Proposition 35 with the use of the second part of Remark 29 we see that the structure constants  $c_{\alpha_3}, c_{\alpha_4}, c'_{\alpha_3}, c'_{\alpha_4}$  are well defined modulo units in  $R$  represented by  $f$ . Since  $f_5(G_K) = f_5(G'_K)$  we get

$$\langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c_{\alpha_3}, c_{\alpha_4} \rangle\rangle \overset{K}{\simeq} \langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c'_{\alpha_3}, c'_{\alpha_4} \rangle\rangle$$

and hence

$$\langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c_{\alpha_3}, c_{\alpha_4} \rangle\rangle \overset{R}{\simeq} \langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c'_{\alpha_3}, c'_{\alpha_4} \rangle\rangle. \tag{48}$$

By Witt cancellation we can write  $c'_{\alpha_3}$  in the form  $c'_{\alpha_3} = a_1 c_{\alpha_3} + a_2 c_{\alpha_4} - a_3 c_{\alpha_3} c_{\alpha_4}$  where  $a_1, a_2, a_3$  are units in  $R$  represented by  $f$  and  $a_1 c_{\alpha_3} - a_3 c_{\alpha_3} c_{\alpha_4}$  is also a unit in  $R$ . Since  $c_{\alpha_3}, c_{\alpha_4}$  are defined modulo values of  $f$  passing to another maximal torus in  $G$  we may assume without loss of generality that  $a_1 = a_2 = 1$ . The rest of the proof is the same as in Proposition 30.

Finally we claim that we may assume that  $c_{\alpha_4} = c'_{\alpha_4}$ . Indeed, from (48) and Witt cancellation we conclude that  $c'_{\alpha_4}$  is of the form  $c'_{\alpha_4} = a c_{\alpha_4}$  where  $a$  is a unit in  $R$  represented by  $\langle\langle d, c_{\alpha_1}, c_{\alpha_2}, c_{\alpha_3} \rangle\rangle$ . Copying the proof of Proposition 39 we easily complete the proof of the claim. Thus Theorem 2 is proven.  $\square$

## REFERENCES

- [Bourb68] N. Bourbaki, Groupes et Algèbres de Lie, Chapitres 4,5 et 6, Hermann, Paris 1968.
- [Ch] V. Chernousov, The kernel of the Rost invariant, Serre’s Conjecture II and the Hasse principle for quasi-split groups  ${}^{3,6}D_4$ ,  $E_6$ ,  $E_7$ , Math. Annalen 326 (2003), 297–330.
- [ChS] V. Chernousov, J.-P. Serre, Lower bounds for essential dimensions via orthogonal representations, Journal of Algebra, 305 (2006), 1055 – 1070.
- [ChP] V. Chernousov, I. Panin, Purity of  $G_2$ -torsors, C. R. Math. Acad. Sci. Paris, 345 (2007), 307–312.
- [CTO] J.-L. Colliot-Thélène, M. Ojanguren, Espaces principaux homogènes génériquement triviaux, Publ. Math. IHES 75 (1992), 97–122.
- [CTS] J.-L. Colliot-Thélène, J.-J. Sansuc, Principal homogeneous spaces under flasque tori: Applications, Journal of Algebra 106 (1987), 148–205.
- [DG] M. Demazure, A. Grothendieck, Structure des Schémas en Groupes Réductifs, SGA 3 III, LN 153, Springer-Verlag, 1970.
- [GMS03] R. S. Garibaldi, A. Merkurjev, J.-P. Serre, Cohomological invariants in Galois cohomology, University Lecture Series, vol. 28, Amer. Math. Soc., 2003.
- [G00] P. Gille, Invariants cohomologiques de Rost en caractéristique positive, K-Theory 21 (2000), 57–100.
- [Gr58] A. Grothendieck, Torsion homologique et sections rationnelles, *Annales de Chow et applications*, Séminaire Chevalley, 2<sup>e</sup> année, Secrétariat mathématique, Paris, 1958.
- [Gr68] A. Grothendieck, Le groupe de Brauer II, in *Dix exposés sur la cohomologie des schémas*, Amsterdam, North-Holland, 1968.
- [Hum] J. Humphreys, Linear Algebraic Groups, Springer-Verlag 1975.
- [Ma] H. Matsumura, Commutative algebra, W.A. Benjamin Co., New York, 1970.
- [M] A. Merkurjev, On the norm residue symbol of degree 2, Dokl. Acad. Nauk SSSR, 261 (1981), 542–547.
- [N] Y. Nisnevich, Rationally trivial principal homogeneous spaces and arithmetic of reductive group schemes over Dedekind rings, C. R. Math. Acad. Sci. Paris, Série I, 299 (1984), 5–8.

- [OP] M. Ojanguren, I. Panin, Rationally trivial hermitian spaces are locally trivial, *Math. Z.* 237 (2001), 181–198.
- [OPZ] M. Ojanguren, I. Panin, K. Zainoulline, On the norm principle for quadratic forms, *J. Ramanujan Math. Soc.* 19 (2004), 1–12.
- [P09] I. Panin, Rationally isotropic quadratic spaces are locally isotropic, *Invent. Math.*, 176 (2009), 397–403.
- [Pa09] I. Panin, On Grothendieck-Serre conjecture concerning principal  $G$ -bundles over reductive group schemes: II, Preprint 2009.
- [PSV] I. Panin, A. Stavrova, N. Vavilov, On Grothendieck-Serre conjecture concerning principal  $G$ -bundles over reductive group schemes: I, Preprint 2009.
- [PPS] I. Panin, V. Petrov, A. Stavrova, On Grothendieck-Serre conjecture for simple adjoint group schemes of types  $E_6$ ,  $E_7$ , Preprint 2009.
- [PetRac] H. Petersson, M. Racine, On the invariants mod 2 of Albert Algebras, *Journal of Algebra* 174 (1995), 1049–1072.
- [PS] I. Panin, A. Suslin, On a conjecture of Grothendieck concerning Azumaya algebras, *St. Petersburg Math. J.* 9 (1998), 851–858.
- [PR94] V. Platonov, A. Rapinchuk, *Algebraic Groups and Number Theory*, Academic Press 1994.
- [Ra] M. Racine, A note on Quadratic Jordan Algebras of degree 3, *Trans. Amer. Math. Soc.* 164 (1972), 93–103.
- [R94] M. S. Raghunathan, Principal bundles admitting a rational section, *Invent. Math.* 116 (1994), 409–423.
- [R95] M. S. Raghunathan, Erratum: Principal bundles admitting a rational section, *Invent. Math.* 121 (1995), 223.
- [Ro] M. Rost, A (mod 3) invariant for exceptional Jordan algebras, *C. R. Acad. Sci. Paris, Série I* 315 (1991), 823–827.
- [S58] J.-P. Serre, Espaces fibrés algébriques, in *Anneaux de Chow et applications*, Séminaire Chevalley, 2<sup>e</sup> année, Secrétariat mathématique, Paris, 1958.
- [S93] J.-P. Serre, Cohomologie galoisienne: progrès et problèmes, *Sém. Bourbaki* 1993–1994, exposé 783.
- [Sp] T. A. Springer, The classification of reduced exceptional simple Jordan algebras, *Indag. Math.* 22 (1960), 414–422.

- [SGA3] Séminaire de Géométrie algébrique de l’I.H.E.S., 1963–1964, Schémas en groupes, dirigé par M. Demazure et A. Grothendieck, Lecture Notes in Math. 151–153, Springer (1970).
- [St] R. Steinberg, Lectures on Chevalley Groups, Yale University 1967.
- [V] V. Voskresenskii, Algebraic Groups and Their Birational Invariants, Translation of Mathematical Monographs, Volume 179, AMS.
- [Z] K. Zainoulline, On Grothendieck conjecture about principal homogeneous spaces for some classical algebraic groups, St. Petersburg Math. J. 12 (2001), 117–143.

V. Chernousov  
Department of Mathematics  
University of Alberta  
Edmonton, Alberta  
T6G 2G1, Canada  
chernous@math.ualberta.ca

