ICM 1998, Berlin, Aug. 18–27

Abstracts of Short Communications and Posters

Section:

15. Numerical Analysis and Scientific Computing

1991 MS Classification: 65Y20, 68Q25, 65H05, 65G10
Kirrinnis, Peter, Universität Bonn, Germany

Fast algorithms for partial fractions

An approximate partial fraction decomposition (APFD) of a rational function $f = q/p \in \mathbf{C}(z)$ is given by approximate factors $p_1, \ldots, p_l$ of p and corresponding numerators $q_1, \ldots, q_l$. Define $\tilde{p}$ and $\tilde{q}$ by $\tilde{p} = p_1 \cdots p_l$ and $\tilde{q}/\tilde{p} = \sum q_j/p_j$. The parameters $s, t, \gamma \in \mathbf{N}$ measure the quality of the APFD as follows: An (s,t,γ)-APFD fulfils $\|p - \tilde{p}\| < 2^{-s}$ and $\|q - \tilde{q}\| < 2^{-t}$ for a norm $\|\cdot\|$ on $\mathbf{C}[z]$, and for all j, any two roots of p_j have distance at most $2^{-\gamma}$.

We describe and analyze algorithms that compute an (s,t,γ)-APFD from the coefficients of q and p with $\deg q \le \deg p \le n$ with $O(n^3 \cdot \gamma + n \cdot (s+t))$ single precision operations (neglecting logarithmic factors). The relation to root finding algorithms is discussed.

The author's papers in *Proc. ISSAC 1992, ACM* (pp. 16–26) and *Proc. ISSAC 1998, ACM* (to appear) provide details.