

# INCOMPRESSIBILITY OF ORTHOGONAL GRASSMANNIANS

NIKITA A. KARPENKO

ABSTRACT. We prove the following conjecture due to Bryant Mathews (2008). Let  $Q$  be the orthogonal grassmannian of totally isotropic  $i$ -planes of a non-degenerate quadratic form  $q$  over an arbitrary field (where  $i$  is an integer satisfying  $1 \leq i \leq (\dim q)/2$ ). If the degree of each closed point on  $Q$  is divisible by  $2^i$  and the Witt index of  $q$  over the function field of  $Q$  is equal to  $i$ , then the variety  $Q$  is 2-incompressible.

Theorem 7, proved below, has been conjectured in the Ph.D. thesis [12, page 24] (a preprint with the conjecture appeared one year earlier).

We start with some development of the theory of canonical dimension of general projective homogeneous varieties (which might be of independent interest). We fix a prime  $p$ . Let  $G$  be a semisimple affine algebraic group over a field  $F$  such that  $G_E$  is of inner type for some finite galois field extension  $E/F$  of degree a power of  $p$  ( $E = F$  is allowed). Let  $X$  be a projective  $G$ -homogeneous  $F$ -variety. We refer to [5] for a definition and discussion of the notion of *canonical  $p$ -dimension*  $\text{cdim}_p X$  of  $X$ . Actually, canonical  $p$ -dimension is defined in the context of more general algebraic varieties. For any irreducible smooth projective variety  $X$ ,  $\text{cdim}_p X$  is the minimal dimension of a closed subvariety  $Y \subset X$  with a 0-cycle of  $p$ -coprime degree on  $Y_{F(X)}$ . Recall that a smooth projective  $X$  is  *$p$ -incompressible*, if it is irreducible and  $\text{cdim}_p X = \dim X$ .

**Proposition 1.** *For  $d := \text{cdim}_p X$ , there exist a cycle class  $\alpha \in \text{CH}^d X_{F(X)}$  (over  $F(X)$ ) of codimension  $d$  and a cycle class  $\beta \in \text{CH}_d X$  (over  $F$ ) of dimension  $d$  such that the degree of the product  $\beta_{F(X)} \cdot \alpha$  is not divisible by  $p$ .*

*Proof.* We use the Chow motives with coefficients in  $\mathbb{F}_p := \mathbb{Z}/p\mathbb{Z}$  (as defined in [3, Chapter XII]) and write  $\text{Ch}$  for the Chow group  $\text{CH}$  modulo  $p$ .

Let  $U(X)$  be the *upper* motive of  $X$ . By definition,  $U(X)$  is a direct summand of the motive  $M(X)$  of  $X$  such that  $\text{Ch}^0 U(X) \neq 0$ . By [5, Theorem 5.1 and Proposition 5.2],  $U(X)$  is also a direct summand of  $M(X)(d - m)$ , where  $m := \dim X$ . The composition

$$M(X) \rightarrow U(X) \rightarrow M(X)(d - m)$$

is given by a correspondence  $f \in \text{Ch}^d(X \times X)$ ; the composition

$$M(X)(d - m) \rightarrow U(X) \rightarrow M(X)$$

is given by a correspondence  $g \in \text{Ch}_d(X \times X)$ . The composition of correspondences  $g \circ f \in \text{Ch}_m(X \times X)$  is a projector on  $X$  such that  $U(X) = (X, g \circ f)$ . In particular, the

---

*Date:* June 2011.

*Key words and phrases.* Algebraic groups, quadratic forms, projective homogeneous varieties, Chow groups and motives. *Mathematical Subject Classification (2010):* 14L17; 14C25.

*multiplicity*  $\text{mult}(g \circ f)$  of the correspondence  $g \circ f$  is  $1 \in \mathbb{F}_p$ . Taking for  $\alpha$  the pull-back of  $f$  with respect to the morphism

$$\text{Spec } F(X) \times X \rightarrow X \times X$$

induced by the generic point of the first factor, and taking for  $\beta$  the push-forward of  $g$  with respect to the projection of  $X \times X$  onto the first factor, we get that

$$\deg(\beta_{F(X)} \cdot \alpha) = \text{mult}(g \circ f) = 1 \in \mathbb{F}_p. \quad \square$$

**Corollary 2.** *The canonical  $p$ -dimension  $\text{cdim}_p X$  of  $X$  is the minimal integer  $d$  such that there exists a cycle class  $\alpha \in \text{Ch}^d X_{F(X)}$  and a cycle class  $\beta \in \text{Ch}_d X$  with  $\deg(\beta_{F(X)} \cdot \alpha) = 1 \in \mathbb{F}_p$ .*

*Proof.* We only need to show that  $\text{cdim}_p X \leq d$ . The proof is similar to [11, Proof of  $\leq$  in Theorem 5.8]. We may assume that  $\beta = [Y]$  for a closed irreducible  $d$ -dimensional subvariety  $Y \subset X$ . Since the product  $[Y]_{F(X)} \cdot \alpha$  can be represented by a cycle on  $Y_{F(X)}$ , the variety  $Y_{F(X)}$  has a 0-cycle of  $p$ -coprime degree. Therefore  $\text{cdim}_p X \leq \dim Y = d$ .  $\square$

**Corollary 3.** *In the situation of Proposition 1, for any field extension  $L/F$ , the change of field homomorphism  $\text{Ch}_d X \rightarrow \text{Ch}_d X_L$  is non-zero.*

*Proof.* The image of  $\beta \in \text{Ch}_d X$  in  $\text{Ch}_d X_L$  is non-zero because  $\deg(\beta_{L(X)} \cdot \alpha_{L(X)}) = 1$ .  $\square$

**Remark 4.** If the variety  $X$  is *generically split* (meaning that the motive of  $X_{F(X)}$  is a sum of Tate motives which, in particular, implies that the adjoint algebraic group acting on  $X$  is of inner type), then [11, Theorem 5.8] says that  $\text{cdim}_p X$  is the minimal  $d$  with non-zero  $\text{Ch}_d X \rightarrow \text{Ch}_d X_L$  for any  $L$ . Corollary 3 can be considered as a generalization of a part of [11, Theorem 5.8] to the case of a projective  $G$ -homogeneous variety  $X$  which is not necessarily generically split with  $G$  not necessarily of inner type. Note that the statement of [11, Theorem 5.8] in whole fails in such generality. Corollary 2 is its correct replacement (giving the original statement in the case of generically split  $X$ ).

**Lemma 5.** *In the situation of Proposition 1, let  $\alpha, \alpha' \in \text{Ch}^d X_{F(X)}$  and  $\beta, \beta' \in \text{Ch}_d X$  be cycle classes with  $\deg(\beta_{F(X)} \cdot \alpha) = 1 = \deg(\beta'_{F(X)} \cdot \alpha')$ . Then*

$$\deg(\beta_{F(X)} \cdot \alpha') \neq 0 \neq \deg(\beta'_{F(X)} \cdot \alpha).$$

*Proof.* We fix an algebraically closed field containing  $F(X)$  and write  $\bar{\cdot}$  when considering a variety or a cycle class over that field. The surjectivity of the pull-back with respect to the flat morphism  $\text{Spec } F(X) \times X \rightarrow X \times X$  induced by the generic point of the first factor of the product  $X \times X$ , tells us that the group  $\text{Ch}^d(\bar{X} \times \bar{X})$  contains a *rational* (i.e., ‘‘coming from  $F$ ’’) cycle class of the form  $[\bar{X}] \times \bar{\alpha} + \cdots + \bar{\gamma} \times [\bar{X}]$  with some  $\gamma \in \text{Ch}^d X_{F(X)}$ , where  $\cdots$  is in the sum of products  $\text{Ch}^i \bar{X} \otimes \text{Ch}^j \bar{X}$  with  $0 < i, j < d$  and  $i + j = d$ . Multiplying by  $[\bar{X}] \times \bar{\beta}$ , we get a rational cycle class of the form  $[\bar{X}] \times [\mathbf{pt}] + \cdots + \bar{\gamma} \times \bar{\beta}$ , where  $\mathbf{pt}$  is a rational point on  $\bar{X}$  and  $\cdots$  is now in the sum of  $\text{Ch}^i \bar{X} \otimes \text{Ch}_i \bar{X}$  with  $0 < i < d$ . The composition of the obtained correspondence with itself equals  $[\bar{X}] \times [\mathbf{pt}] + \cdots + \deg(\gamma \cdot \beta)(\bar{\gamma} \times \bar{\beta})$ . Since an appropriate power of this correspondence is a multiplicity 1 *projector* (cf. [9, Corollary 3.2] or [7]) and  $d = \text{cdim}_p X$ , it follows by [5, Theorem 5.1] that  $\deg(\gamma \cdot \beta) \neq 0$ . Now multiplying  $[\bar{X}] \times \bar{\alpha} + \cdots + \bar{\gamma} \times [\bar{X}]$  by  $\bar{\beta} \times [\bar{X}]$ , transposing, and raising to  $(p - 1)$ th

power (by means of composition of correspondences), we get a rational cycle of the form  $[\bar{X}] \times [\mathbf{pt}] + \cdots + \bar{\alpha} \times \bar{\beta}$ .

Similarly, there is a rational cycle of the form  $[\bar{X}] \times [\mathbf{pt}] + \cdots + \bar{\alpha}' \times \bar{\beta}'$ . One of its compositions with the previous one produces  $[\bar{X}] \times [\mathbf{pt}] + \cdots + \deg(\beta' \cdot \alpha)(\bar{\alpha}' \times \bar{\beta})$ , therefore  $\deg(\beta' \cdot \alpha) \neq 0 \in \mathbb{F}_p$ . The other composition produces  $[\bar{X}] \times [\mathbf{pt}] + \cdots + \deg(\beta \cdot \alpha')(\bar{\alpha} \times \bar{\beta}')$ , so that  $\deg(\beta \cdot \alpha') \neq 0$ .  $\square$

We specify as follows the settings of Proposition 1. The prime  $p$  is now 2. The algebraic group  $G$  is now  $O^+(q)$  for a non-degenerate quadratic form  $q$  (one may take  $E = F$  if  $\dim q$  is odd or  $\text{disc } q$  is trivial, otherwise  $E$  can be the quadratic field extension of  $F$  given by the discriminant of  $q$ ). We set  $n := \dim q$ . For any integer  $i$  with  $0 \leq i \leq n/2$ , let  $Q_i$  be the variety of  $i$ -dimensional totally isotropic subspaces in  $q$ . In particular,  $Q_0 = \text{Spec } F$ . For  $i$  with  $0 < i < n/2$ ,  $Q_i$  is a projective  $G$ -homogeneous variety.

**Corollary 6.** *If  $\text{cdim}_2 Q_i = \text{cdim}_2 Q'_{i-1} = \dim Q'_{i-1}$  for some  $i$  with  $0 < i < n/2$ , where  $Q'_{i-1}$  is the orthogonal grassmannian of totally isotropic  $(i-1)$  planes of a  $(n-2)$ -dimensional quadratic form  $q'$  over  $F(Q_1)$  Witt-equivalent to  $q_{F(Q_1)}$ , then  $\deg \text{CH } Q_i \ni 2^{i-1}$ .*

*Proof.* The statement being trivial for  $i = 1$ , we may assume that  $i \geq 2$ .

For  $d := \text{cdim}_2 Q_i$ , using Proposition 1, we choose some  $\alpha \in \text{CH}^d Q_{iF(Q_i)}$  and  $\beta \in \text{CH}_d Q_i$  with odd  $\deg(\beta_{F(Q_i)} \cdot \alpha)$ . Note that  $\text{cdim}_2 Q_{iF(Q_1)} = \text{cdim}_2 Q'_{i-1} = d$ . We construct some special  $\alpha' \in \text{CH}^d Q_{iF(Q_1)(Q_i)}$  and  $\beta' \in \text{CH}_d Q_{iF(Q_1)}$  with  $\deg(\beta'_{F(Q_1)(Q_i)} \cdot \alpha') = 1$  as follows. Let  $Q_{1 \subset i}$  be the variety of  $(1, i)$ -flags of totally isotropic subspaces in  $q$  together with the evident projections  $Q_{1 \subset i} \rightarrow Q_1, Q_i$ . We define  $\beta'$  as the pull-back via  $Q_{1 \subset i F(Q_1)} \rightarrow Q_{1F(Q_1)}$  followed by the push-forward via  $Q_{1 \subset i F(Q_1)} \rightarrow Q_{iF(Q_1)}$  of the rational point class  $l_0$  on  $Q_{1F(Q_1)}$ . We define  $\alpha'$  as the product  $z_{i-1} \cdots z_1$ , where  $z_j$  is the pull-back via  $Q_{1 \subset i F(Q_1)(Q_i)} \rightarrow Q_{1F(Q_1)(Q_i)}$  followed by the push-forward via  $Q_{1 \subset i F(Q_1)(Q_i)} \rightarrow Q_{iF(Q_1)(Q_i)}$  of the class  $l_j$  of a  $j$ -dimensional projective subspace on  $Q_{1F(Q_1)(Q_i)}$ . The degree condition on  $\alpha'$  and  $\beta'$  is satisfied by [14, Statement 2.13]. Fixing an algebraically closed field containing  $F(Q_1)(Q_i)$ , we see by Lemma 5 that the product  $\bar{\beta} \cdot \bar{\alpha}'$  is an odd degree 0-cycle class on  $Q_i$ . Moreover, the class  $\bar{\beta}$  is rational. Since  $2\bar{z}_j$  is rational for every  $j < (n-2)/2$  (by the reason that  $2l_j$  is rational), the class  $2^{i-1}\bar{\beta}\bar{\alpha}$  is also rational and it follows that  $2^{i-1} \in \deg \text{CH } Q_i$ .  $\square$

We come to the main result of this note. It is known for  $i = 1$  by [4] (the proof is essentially contained already in [13]; the characteristic 2 case has been treated later on in [3]). The case of maximal  $i$ , i.e., of  $i = [n/2]$ , is also known and is discussed in the beginning of the proof. For  $i = 2$  and odd-dimensional  $q$ , it has been proved in [12] (the proof for  $i = 2$  given here is different; in particular, it does not make use of the motivic decompositions of [2] for products of projective homogeneous varieties).

**Theorem 7.** *Let  $q$  be a non-degenerate quadratic form over a field  $F$ . Let  $i$  be an integer satisfying  $1 \leq i \leq (\dim q)/2$ . If the degree of every closed point on  $Q_i$  is divisible by  $2^i$  and the Witt index of the quadratic form  $q_{F(Q_i)}$  equals  $i$ , then the variety  $Q_i$  is 2-incompressible (i.e.,  $\text{cdim}_2 Q_i = \dim Q_i$ ).*

*Proof.* We set where  $n := \dim q$ . Note that for  $i = n/2$  (and even  $n$ ) the condition on closed points on  $Q_{n/2}$  ensures that  $\text{disc } q$  is non-trivial. In particular,  $Q_{n/2}$  is irreducible.

In general, for even  $n$ , the variety  $Q_{n/2}$  is isomorphic to the orthogonal grassmannian of totally isotropic  $(n/2 - 1)$ -planes of  $q^1$  considered as a variety over  $F$ , where  $q^1$  is any 1-codimensional non-degenerate subform in  $q_K$ , and  $K$  is the quadratic étale  $F$ -algebra given by the discriminant of  $q$ . Therefore the statement of Theorem 7 for  $i = n/2$  follows from the statement for  $i = (n - 1)/2$ . By this reason, we do not consider the case of  $i = n/2$  below. In particular,  $Q_i$  below is a projective  $G$ -homogeneous variety.

We induct on  $n$ . There is nothing to prove for  $n < 3$ . Below we are assuming that  $n \geq 3$ .

Over the field  $F(Q_1)$ , the motive of  $Q_{iF(Q_1)}$  decomposes as follows (cf. [6], [8] or [1]):

$$M(Q'_{i-1}) \oplus M(Q'_i)((\dim Q_i - \dim Q'_i)/2) \oplus M(Q'_{i-1})(\dim Q_i - \dim Q'_{i-1}),$$

where, as in Corollary 6,  $Q'_j$  is the variety of  $j$ -dimensional totally isotropic subspaces of a quadratic form  $q'$  over the field  $F(Q_1)$  such that  $q_{F(Q_1)}$  is isomorphic to the orthogonal sum of  $q'$  and a hyperbolic plane. Since  $n' := \dim q' = n - 2 < n$ , the variety  $Q'_{i-1}$  is 2-incompressible by the induction hypothesis (more precisely, the induction hypothesis is applied if  $i \geq 2$ , for  $i = 1$  the statement is trivial). Indeed, since the extension  $F(Q_1)/F$  is a tower of a purely transcendental extension followed by a quadratic one, the degree of any closed point on  $Q'_{i-1}$  is divisible by  $2^{i-1}$ ; the Witt index of  $q'_{F(Q_1)(Q'_{i-1})}$  is  $i - 1$ , that is, the Witt index of  $q_{F(Q_1)(Q'_{i-1})}$  is  $i$  because the field extension  $F(Q_1)(Q'_{i-1})(Q_i)/F(Q_i)$  is purely transcendental.

By [10, Theorem 1.1] (cf. [6]), it follows that the motive of  $Q'_{i-1}$  decomposes in a direct sum of one copy of  $U(Q'_{i-1})$  and shifts of  $U(Q'_j)$  with various  $j \geq i$ . The motive of  $Q'_i$  decomposes in a direct sum of shifts of  $U(Q'_j)$  with various  $j \geq i$ . Note that the condition on the Witt index of the form  $q_{F(Q_i)}$  ensures that for any  $j \geq i$  the motive  $U(Q'_{i-1})$  is not isomorphic to  $U(Q'_j)$ . Therefore the complete motivic decomposition of  $Q_{iF(Q_1)}$  contains one copy of  $U(Q'_{i-1})$ , one copy of  $U(Q'_{i-1})(\dim Q_i - \dim Q'_{i-1})$  and no other shifts of  $U(Q'_{i-1})$ .

The complete decomposition of  $U(Q_i)_{F(Q_1)}$  contains the summand  $U(Q'_{i-1})$ . If it also contains the shift of  $U(Q'_{i-1})$ , then  $\text{cdim}_2 Q_i = \dim Q_i$  and we are done. Otherwise, by [6, Lemma 1.2 and Remark 1.3],  $\text{cdim}_2 Q_i = \text{cdim}_2 Q'_{i-1} = \dim Q'_{i-1}$ , and we get by Corollary 6 that  $Q_i$  has a closed point of degree not divisible by  $2^i$ .  $\square$

## REFERENCES

- [1] BROSNAN, P. On motivic decompositions arising from the method of Białynicki-Birula. *Invent. Math.* 161, 1 (2005), 91–111.
- [2] CHERNOUSOV, V., AND MERKURJEV, A. Motivic decomposition of projective homogeneous varieties and the Krull-Schmidt theorem. *Transform. Groups* 11, 3 (2006), 371–386.
- [3] ELMAN, R., KARPENKO, N., AND MERKURJEV, A. *The algebraic and geometric theory of quadratic forms*, vol. 56 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2008.
- [4] KARPENKO, N., AND MERKURJEV, A. Essential dimension of quadrics. *Invent. Math.* 153, 2 (2003), 361–372.
- [5] KARPENKO, N. A. Canonical dimension. Proceedings of the ICM 2010, vol. II, 146–161.

- [6] KARPENKO, N. A. Incompressibility of generic orthogonal grassmannians. *Linear Algebraic Groups and Related Structures* (preprint server) 409 (2010, Nov 23), 7 pages.
- [7] KARPENKO, N. A. Upper motives of algebraic groups and incompressibility of Severi-Brauer varieties. *Linear Algebraic Groups and Related Structures* (preprint server) 333 (2009, Apr 3, revised: 2009, Apr 24), 18 pages.
- [8] KARPENKO, N. A. Cohomology of relative cellular spaces and of isotropic flag varieties. *Algebra i Analiz* 12, 1 (2000), 3–69.
- [9] KARPENKO, N. A. Hyperbolicity of orthogonal involutions. *Doc. Math. Extra Volume: Andrei A. Suslin's Sixtieth Birthday* (2010), 371–392 (electronic). With an Appendix by Jean-Pierre Tignol.
- [10] KARPENKO, N. A. Upper motives of outer algebraic groups. In *Quadratic forms, linear algebraic groups, and cohomology*, vol. 18 of *Dev. Math.* Springer, New York, 2010, pp. 249–258.
- [11] KARPENKO, N. A., AND MERKURJEV, A. S. Canonical  $p$ -dimension of algebraic groups. *Adv. Math.* 205, 2 (2006), 410–433.
- [12] MATHEWS, B. G. *Canonical dimension of projective homogeneous varieties of inner type A and type B*. ProQuest LLC, Ann Arbor, MI, 2009. Thesis (Ph.D.)—University of California, Los Angeles.
- [13] VISHIK, A. Direct summands in the motives of quadrics. Preprint, 1999, 13 pages. Available on the web page of the author.
- [14] VISHIK, A. Fields of  $u$ -invariant  $2^r + 1$ . In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II*, vol. 270 of *Progr. Math.* Birkhäuser Boston Inc., Boston, MA, 2009, pp. 661–685.

UPMC UNIV PARIS 06, INSTITUT DE MATHÉMATIQUES DE JUSSIEU, F-75252 PARIS, FRANCE

*Web page:* [www.math.jussieu.fr/~karpenko](http://www.math.jussieu.fr/~karpenko)

*E-mail address:* [karpenko at math.jussieu.fr](mailto:karpenko@math.jussieu.fr)