Cyclotomic q–Schur algebras

Richard Dipper
Mathematische Institut B
Universität Stuttgart
Postfach 80 11 40
70550 Stuttgart
Deutschland

Gordon James
Department of Mathematics
Imperial College
Queen's Gate
London SW7 2BZ
England

Andrew Mathas
Fakultät für Mathematik
Universität Bielefeld
P.O. Box 100 131
33501 Bielefeld
Deutschland

1 Introduction

Recently, we [**9**] and, independently, Du and Scott [**11**], defined an analogue of the q–Schur algebra [**6**, **7**] for an Iwahori–Hecke algebra of type **B**. In this paper we study an analogue of the q–Schur algebra for an arbitrary Ariki–Koike algebra.

The Ariki–Koike algebra $\mathscr{H}$ is a cyclotomic algebra of type $G(r,1,n)$ [**2**], and it becomes the Iwahori–Hecke algebra of type **A** or **B** when $r = 1$ or 2 respectively. By working over a ring R which contains a primitive rth root of unity, and by specializing the parameters appropriately, the Ariki–Koike algebra turns into the group algebra $R(C_r \wr \mathfrak{S}_n)$ of the wreath product of the cyclic group C_r of order r with the symmetric group $\mathfrak{S}_n$ of degree n.

For each multicomposition λ of n, we construct a right ideal M^λ of $\mathscr{H}$ (see Definition 3.8). The cyclotomic q–Schur algebra is then defined to be $\mathscr{S} = \mathrm{End}_{\mathscr{H}}(\oplus_\lambda M^\lambda)$. (Under the specialization above where $\mathscr{H} \cong R(C_r \wr \mathfrak{S}_n)$, the module M^λ becomes a module induced from a subgroup of the form $(C_r \times \cdots \times C_r) \rtimes \mathfrak{S}_\lambda$.)

A.M.S. subject classification (1991): 16G99, 20C20, 20G05
This paper is a contribution to the DFG project on "Algorithmic number theory and algebra". The authors acknowledge support from DFG; the third author was also supported in part by SERC grant GR/J37690 and by the Sonderforschungsbereich 343 in Bielefeld.

In this paper we construct a cellular basis for the cyclotomic q–Schur algebra. As a consequence we obtain a Weyl module W^λ for each multipartition λ of n. We show that W^λ has simple head F^λ and that the set $\{F^\lambda\}$, as λ ranges over the multipartitions of n, is a complete set of non–isomorphic irreducible $\mathscr{S}$–modules. Using the cellular structure of $\mathscr{S}$, it is now easy to see that the cyclotomic q–Schur algebra is quasi–hereditary.

In order to prove these results about the cyclotomic q–Schur algebra, we need to examine the ideals M^λ in some detail. Using the cellular structure of the Ariki–Koike algebra $\mathscr{H}$ (cf. [**10**, **13**]), we obtain a basis of M^λ and a special series of submodules of M^λ, known as a Specht series. From the Specht series of M^λ we construct the cellular basis of $\mathscr{S}$.

2 The Ariki–Koike algebra

Throughout this paper, r and n will be fixed positive integers with $r \geq 1$ and $n \geq 1$.

Let R be a commutative ring with 1 and let $q, Q_1, \dots, Q_r$ be elements of R with q invertible. The Ariki–Koike algebra $\mathscr{H}$ is the associative unital R–algebra with generators $T_0, T_1, \dots, T_{n-1}$ subject to the following relations

$$\begin{aligned}
(T_0 - Q_1)\cdots(T_0 - Q_r) &= 0 && \\
T_0T_1T_0T_1 &= T_1T_0T_1T_0 && \\
(T_i + 1)(T_i - q) &= 0 && \text{for } 1 \leq i \leq n-1\\
T_{i+1}T_iT_{i+1} &= T_iT_{i+1}T_i && \text{for } 1 \leq i \leq n-2\\
T_iT_j &= T_jT_i && \text{for } 0 \leq i < j-1 \leq n-2.
\end{aligned}$$

Suppose that $\pi \in R$ is a primitive rth root of 1 and that $q = 1$ and $Q_k = \pi^k$ for $k = 1, 2, \dots, r$. Then it follows from the definition that $\mathscr{H}$ is isomorphic to $R(C_r \wr \mathfrak{S}_n)$.

Let $\mathfrak{S}_n = \mathfrak{S}(\{1, 2, \dots, n\})$ be the symmetric group on $\{1, 2, \dots, n\}$ and let $s_i = (i, i+1)$ for $1 \leq i < n$. Then $s_1, s_2, \dots, s_{n-1}$ are the standard Coxeter generators of $\mathfrak{S}_n$. If $w \in \mathfrak{S}_n$ then a word $w = s_{i_1} \dots s_{i_k}$ for w is a reduced expression for w if k is minimal; in this case we say that w has length k and write $\ell(w) = k$. Given a reduced expression $s_{i_1} \dots s_{i_k}$ for $w \in \mathfrak{S}_n$, let $T_w = T_{i_1} \dots T_{i_k}$; the relations in $\mathscr{H}$ ensure that T_w is independent of the choice of reduced expression. We denote by $\mathscr{H}(\mathfrak{S}_n)$ the subalgebra of $\mathscr{H}$ generated by $T_1, T_2, \dots, T_{n-1}$. Then $\mathscr{H}(\mathfrak{S}_n)$ has basis $\{\, T_w \,|\, w \in \mathfrak{S}_n \,\}$, and it is isomorphic to an Iwahori–Hecke algebra of type **A**.

Define elements $L_m = q^{1-m}T_{m-1} \dots T_1T_0T_1 \dots T_{m-1}$ for $m = 1, 2, \dots, n$; these are analogues of the q–Murphy operators of the Iwahori–Hecke algebras of type **A** [**5**, **15**]. An easy calculation using the relations in $\mathscr{H}$ (cf. [**5**, (2.1), (2.2)]) shows that we have the following results.

(2.1) Suppose that $1 \leq i \leq n-1$ and $1 \leq m \leq n$. Then

(i) L_i and L_m commute.

(ii) T_i and L_m commute if $i \neq m-1, m$.

(iii) T_i commutes with L_iL_{i+1} and $L_i + L_{i+1}$.

(iv) If $a \in R$ and $i \neq m$ then T_i commutes with $(L_1 - a)(L_2 - a)\dots(L_m - a)$.

Using the elements T_w and L_m defined above, Ariki and Koike gave a basis for $\mathscr{H}$ as follows.

(2.2) Theorem (Ariki–Koike [**1**, (3.10)]) *The algebra $\mathscr{H}$ is a free R–module with basis*

$$\{\, L_1^{c_1} L_2^{c_2} \dots L_n^{c_n} T_w \mid w \in \mathfrak{S}_n \quad \textit{and} \quad 0 \leq c_m \leq r-1 \quad \textit{for} \quad m = 1, 2, \dots, n \,\}.$$

In particular, $\mathscr{H}$ is free of rank $r^n n!$

(2.3) Let $*$ be the R–linear antiautomorphism of $\mathscr{H}$ determined by $T_i^* = T_i$ for all i with $0 \leq i \leq n-1$. Then $T_w^* = T_{w^{-1}}$ and $L_m^* = L_m$ for all $w \in \mathfrak{S}_n$ and $m = 1, 2, \dots, n$.

We therefore have the following result.

(2.4) $\{\, T_w L_1^{c_1} L_c^{c_2} \dots L_n^{c_n} \mid w \in \mathfrak{S}_n \quad \text{and} \quad 0 \leq c_m \leq r-1 \quad \text{for} \quad m = 1, 2, \dots, n \,\}$ is a basis of $\mathscr{H}$.

3 A cellular basis of $\mathscr{H}$

In their paper [**12**], which introduced the concept of cellular algebras, Graham and Lehrer gave a cellular basis of $\mathscr{H}$, using the Kazhdan–Lusztig basis of $\mathscr{H}(\mathfrak{S}_n)$. We require a different cellular basis, namely one similar to the basis of $\mathscr{H}(\mathfrak{S}_n)$ introduced by Murphy [**16**]. Although the construction of the cellular basis of $\mathscr{H}$ in this section is similar to that in [**10**, **13**], we are obliged to keep track of new information concerned with the cellular basis (see Corollary 3.24 below).

Consider the R–submodule of $\mathscr{H}$ which is spanned by

$$\{\, L_1^{c_1} L_2^{c_2} \dots L_n^{c_n} \mid 0 \leq c_i \leq r-1 \quad \text{for} \quad 1 \leq i \leq n \,\}.$$

When $q = 1$, this is a subalgebra of $\mathscr{H}$, but one of the main difficulties of working with the Ariki–Koike algebra is that it is not a subalgebra in general. (To see this, consider L_2^2 when $r = 2$.) We shall need certain elements $u_{\mathbf{a}}^+$ of this R–submodule of $\mathscr{H}$, and we introduce these now.

(3.1) Definition *Suppose that $\mathbf{a} = (a_1, \dots, a_r)$ is an r–tuple of integers a_i such that $0 \leq a_i \leq n$ for all i. Let $u_{\mathbf{a}}^+ = u_{\mathbf{a},1} u_{\mathbf{a},2} \dots u_{\mathbf{a},r}$ where*

$$u_{\mathbf{a},k} = \prod_{m=1}^{a_k} (L_m - Q_k) \quad \textit{for} \quad 1 \leq k \leq r.$$

(3.2) Remarks (i) Suppose that every a_k is non–zero. Then $(L_1 - Q_k)$ is a factor of each $u_{\mathbf{a},k}$; so $u^+_{\mathbf{a}}$ has a factor

$$\prod_{k=1}^{r}(L_1 - Q_k) = \prod_{k=1}^{r}(T_0 - Q_k) = 0.$$

Therefore, $u^+_{\mathbf{a}}$ is zero in this case.

(ii) Rearranging the order of $a_1, a_2, \dots, a_r$ amounts just to reordering the parameters $Q_1, Q_2, \dots, Q_r$. For example, if we define $u^-_{\mathbf{a}} = u^+_{(a_r, a_{r-1}, \dots, a_1)}$ then $u^-_{\mathbf{a}}$ is obtained from $u^+_{\mathbf{a}}$ by replacing Q_k by Q_{r-k+1} for $1 \le k \le r$.

(iii) In practice, we shall use $u^+_{\mathbf{a}}$ only for r–tuples $\mathbf{a} = (a_1, a_2, \dots, a_r)$ such that $0 = a_1 \le a_2 \le \dots \le a_r \le n$. Our last two remarks show that there is no loss in doing this, and that we could equally well work with the elements $u^-_{\mathbf{a}}$ defined in Remark (ii).

(3.3) Example Suppose that $r = 4$, $n \ge 5$ and $\mathbf{a} = (0, 2, 4, 5)$. Then

$$\begin{aligned} u^+_{\mathbf{a}} \;=\; & (L_1 - Q_2)(L_2 - Q_2) \\ & \times (L_1 - Q_3)(L_2 - Q_3)(L_3 - Q_3)(L_4 - Q_3) \\ & \times (L_1 - Q_4)(L_2 - Q_4)(L_3 - Q_4)(L_4 - Q_4)(L_5 - Q_4). \end{aligned}$$

Our first lemma relates $u^+_{\mathbf{a}}$ and $u^+_{\mathbf{b}}$ when $\mathbf{b}$ is obtained from $\mathbf{a}$ by increasing a single part by one.

(3.4) Lemma *Let* $\mathbf{a} = (a_1, a_2, \dots, a_r)$ *and assume that* $1 \le k \le r$ *and* $a_k + 1 \le n$. *Let* $\mathbf{b} = (a_1, \dots, a_{k-1}, a_k + 1, a_{k+1}, \dots, a_r)$. *Then, for some* $h_1, h_2 \in \mathscr{H}(\mathfrak{S}_n)$, *we have* $u^+_{\mathbf{a}} T_{a_k} T_{a_k - 1} \dots T_1 T_0 = u^+_{\mathbf{a}} h_1 + u^+_{\mathbf{b}} h_2$.

PROOF: The definition of $u^+_{\mathbf{b}}$ gives $u^+_{\mathbf{b}} = u^+_{\mathbf{a}}(L_{a_k+1} - Q_k)$. Hence,

$$u^+_{\mathbf{a}} T_{a_k} T_{a_k-1} \dots T_1 T_0 T_1 \dots T_{a_{k-1}} T_{a_k} = q^{a_k} u^+_{\mathbf{a}} L_{a_k+1} = q^{a_k} Q_k u^+_{\mathbf{a}} + q^{a_k} u^+_{\mathbf{b}}.$$

The desired result follows by postmultiplying by $T^{-1}_{a_k} \dots T^{-1}_1$. □

We next turn our attention to the subalgebra $\mathscr{H}(\mathfrak{S}_n)$ of $\mathscr{H}$. Here we shall need the notation and combinatorics of multipartitions.

A composition $\alpha = (\alpha_1, \alpha_2, \dots)$ is a finite sequence of non–negative integers; we denote by $|\alpha|$ the sum of this sequence. A multicomposition of n is an ordered r–tuple $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ of compositions $\lambda^{(k)}$ such that $\sum_{k=1}^{r} |\lambda^{(k)}| = n$. We call $\lambda^{(k)}$ the kth component of λ. A partition is a composition whose parts are non–increasing; a multicomposition is a multipartition if all its components are partitions.

For each composition $\alpha = (\alpha_1, \alpha_2, \dots)$ with $|\alpha| = m$ we have a Young subgroup $\mathfrak{S}_\alpha = \mathfrak{S}_{\alpha_1} \times \mathfrak{S}_{\alpha_2} \times \cdots$ of $\mathfrak{S}_m$. Similarly, to each multicomposition $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ of n we associate the Young subgroup $\mathfrak{S}_\lambda = \mathfrak{S}_{\lambda^{(1)}} \times \mathfrak{S}_{\lambda^{(2)}} \times \cdots \times \mathfrak{S}_{\lambda^{(r)}}$ of $\mathfrak{S}_n$.

We are now in a position to define certain key elements m_λ of $\mathscr{H}$. The element m_λ depends upon the Young subgroup $\mathfrak{S}_\lambda$ and also involves one of the elements $u^+_{\mathbf{a}}$ defined above.

(3.5) Definition *Suppose that λ is a multicomposition of n and define $\mathbf{a} = (a_1, a_2, \dots, a_r)$ by $a_k = \sum_{i=1}^{k-1} |\lambda^{(i)}|$. Let $x_\lambda = \sum_{w \in \mathfrak{S}_\lambda} T_w$ and set $m_\lambda = u_{\mathbf{a}}^+ x_\lambda$.*

(3.6) Example Suppose that $\lambda = ((2,1),(1^2),(2))$. Then $\mathfrak{S}_\lambda = \mathfrak{S}_2 \times \mathfrak{S}_1 \times \mathfrak{S}_1 \times \mathfrak{S}_1 \times \mathfrak{S}_2$, $\mathbf{a} = (0,3,5)$, and

$$\begin{aligned} m_\lambda \;=\; & (L_1 - Q_2)(L_2 - Q_2)(L_3 - Q_2) \\ & \times (L_1 - Q_3)(L_2 - Q_3)(L_3 - Q_3)(L_4 - Q_3)(L_5 - Q_3) \\ & \times (1 + T_1)(1 + T_6). \end{aligned}$$

(3.7) Remark If $\alpha = (|\lambda^{(1)}|, |\lambda^{(2)}|, \dots, |\lambda^{(r)}|)$ then all of the elements in $\mathscr{H}(\mathfrak{S}_\alpha)$ commute with $u_{\mathbf{a}}^+$ by (2.1)(iv). In particular, $m_\lambda = u_{\mathbf{a}}^+ x_\lambda = x_\lambda u_{\mathbf{a}}^+$. Hence, $m_\lambda^* = m_\lambda$, where $*$ is the antiautomorphism of (2.3).

The $\mathscr{H}$–modules which will be our main concern in this paper are the right ideals generated by the m_λ, as λ varies over the multicompositions of n. The cyclotomic q–Schur algebra will be built from endomorphisms between such right ideals.

(3.8) Definition *Suppose that λ is a multicomposition of n. Let $M^\lambda = m_\lambda \mathscr{H}$.*

We leave the proof of the following remarks to the reader.

(3.9) Remarks (i) If the multicomposition μ is obtained from λ by reordering the parts in each component then $M^\mu \cong M^\lambda$.

(ii) Suppose that $q = 1$ and that $Q_k = \pi^k$, for $k = 1, 2, \dots, r$, where π is a primitive rth root of unity in R. Then $\mathscr{H} \cong R(C_r \wr \mathfrak{S}_n)$. Let $\alpha_k = |\lambda^{(k)}|$ for $1 \le k \le r$. Then M^λ is induced from a module U for the subgroup $(C_r^{\alpha_1} \times C_r^{\alpha_2} \times \cdots \times C_r^{\alpha_r}) \rtimes \mathfrak{S}_\lambda$. The restriction of U to the subgroup $C_r^{\alpha_1} \times \cdots \times C_r^{\alpha_r}$ has the form $U_1^{\otimes \alpha_1} \otimes \cdots \otimes U_r^{\otimes \alpha_r}$ where U_k has rank k for $1 \le k \le r$. The restriction of U to $\mathfrak{S}_\lambda$ is the trivial module.

We shall construct a basis of M^λ, and study $\mathscr{H}$–homomorphisms between the various modules M^λ. To this end, we introduce λ–tableaux.

The diagram of a composition $\alpha = (\alpha_1, \alpha_2, \dots)$ is $\{\, (i,j) \,|\, 1 \le i \text{ and } 1 \le j \le \alpha_i \,\}$, which we regard as an array of nodes, or boxes, in the plane. The diagram of a multicomposition is the ordered r–tuple of the diagrams of its components.

Let λ be a multicomposition of n. A λ–tableau $\mathsf{t} = (\mathsf{t}^{(1)}, \dots, \mathsf{t}^{(r)})$ is obtained from the diagram of λ by replacing each node by one of the integers $1, 2, \dots, n$, allowing no repeats. We call the tableaux $\mathsf{t}^{(k)}$ the components of t.

(3.10) Definition (i) *A λ–tableau is* row standard *if the entries in each row of each component increase from left to right.*

(ii) *A λ–tableau t is* standard *if λ is a multipartition of n, t is row standard and the entries in each column of each component of t increase from top to bottom.*

(iii) *If λ is a multipartition of n, then let $\mathrm{Std}(\lambda)$ be the set of standard λ–tableaux.*

Note, particularly, that while row standard λ–tableaux are defined for all multicompositions λ, there exist standard λ–tableaux only if λ is a multipartition of n.

We require partial orders on the set of multicompositions and on the set of row standard tableaux.

If $\mathfrak{t}$ is a row standard λ–tableau and $1 \leq m \leq n$, then the entries $1, 2, \ldots, m$ in $\mathfrak{t}$ occupy the diagram of a multicomposition; let $\mathfrak{t} \downarrow m$ denote this multicomposition. For example, $\mathfrak{t} \downarrow n = \lambda$. We use this notation in our next definition.

(3.11) Definition *Suppose that* $\lambda = (\lambda^{(1)}, \ldots, \lambda^{(r)})$ *and* $\mu = (\mu^{(1)}, \ldots, \mu^{(r)})$ *are multicompositions of* n.

(i) *We say that* λ dominates μ, *and write* $\lambda \trianglerighteq \mu$, *if*

$$\sum_{i=1}^{k-1} |\lambda^{(i)}| + \sum_{i=1}^{j} \lambda_i^{(k)} \geq \sum_{i=1}^{k-1} |\mu^{(i)}| + \sum_{i=1}^{j} \mu_i^{(k)}$$

for all k *and* j *with* $1 \leq k \leq r$ *and* $j \geq 0$. *If* $\lambda \trianglerighteq \mu$ *and* $\lambda \neq \mu$ *then we write* $\lambda \triangleright \mu$.

(ii) *Suppose that* $\mathfrak{s}$ *is a row standard* λ*–tableau and that* $\mathfrak{t}$ *is a row standard* μ*–tableau. We say that* $\mathfrak{s}$ dominates $\mathfrak{t}$, *and write* $\mathfrak{s} \trianglerighteq \mathfrak{t}$ *if* $\mathfrak{s} \downarrow m \trianglerighteq \mathfrak{t} \downarrow m$ *for all* m *with* $1 \leq m \leq n$. *If* $\mathfrak{s} \trianglerighteq \mathfrak{t}$ *and* $\mathfrak{s} \neq \mathfrak{t}$ *then we write* $\mathfrak{s} \triangleright \mathfrak{t}$.

For example, if $n = r = 2$, then the multipartitions of 2 are ordered by $((2),(0)) \triangleright ((1^2),(0)) \triangleright ((1),(1)) \triangleright ((0),(2)) \triangleright ((0),(1^2))$.

Note that if $\mathfrak{s}$ is a row standard λ–tableau and $\mathfrak{t}$ is a row standard μ tableau such that $\mathfrak{s} \trianglerighteq \mathfrak{t}$ then $\lambda \trianglerighteq \mu$.

Our next definition gives another relation between tableaux.

(3.12) Definition *Suppose that* $\mathfrak{s}$ *is a tableau and that* $1 \leq j \leq n$. *We write* $\mathrm{comp}_{\mathfrak{s}}(j) = k$ *if* j *appears in the* k*th component* $\mathfrak{s}^{(k)}$ *of* $\mathfrak{s}$.

Suppose that $\mathfrak{t}$ *is another tableau. Then* $\mathrm{comp}_{\mathfrak{s}} = \mathrm{comp}_{\mathfrak{t}}$ *if* $\mathrm{comp}_{\mathfrak{s}}(j) = \mathrm{comp}_{\mathfrak{t}}(j)$ *for all* j *with* $1 \leq j \leq n$. *We also write* $\mathrm{comp}_{\mathfrak{s}} \geq \mathrm{comp}_{\mathfrak{t}}$ *if* $\mathrm{comp}_{\mathfrak{s}}(j) \geq \mathrm{comp}_{\mathfrak{t}}(j)$ *for all* $1 \leq j \leq n$; *and* $\mathrm{comp}_{\mathfrak{s}} > \mathrm{comp}_{\mathfrak{t}}$ *if* $\mathrm{comp}_{\mathfrak{s}} \geq \mathrm{comp}_{\mathfrak{t}}$ *and* $\mathrm{comp}_{\mathfrak{s}} \neq \mathrm{comp}_{\mathfrak{t}}$.

(3.13) Example Let

$$\mathfrak{s} = \left(\begin{array}{|c|c|}\hline 1&4\\\hline 5&6\\\hline\end{array}\,,\ \begin{array}{|c|c|c|}\hline 3&8&9\\\hline 2&7\\\cline{1-2}\end{array} \right), \quad \mathfrak{t} = \left(\begin{array}{|c|c|}\hline 2&3\\\hline 4&7\\\hline\end{array}\,,\ \begin{array}{|c|c|c|}\hline 1&6&8\\\hline 5&9\\\cline{1-2}\end{array} \right) \quad \text{and} \quad \mathfrak{u} = \left(\begin{array}{|c|c|c|}\hline 1&2&3\\\hline 4&6&7\\\hline\end{array}\,,\ \begin{array}{|c|c|}\hline 5&9\\\hline 8\\\cline{1-1}\end{array} \right).$$

Then $\mathfrak{s}$ is row standard, and $\mathfrak{t}$ and $\mathfrak{u}$ are standard. We have $\mathfrak{u} \triangleright \mathfrak{s}$ and $\mathfrak{u} \triangleright \mathfrak{t}$ but $\mathfrak{s}$ and $\mathfrak{t}$ are incomparable in the dominance order. Also, $\mathrm{comp}_{\mathfrak{t}} > \mathrm{comp}_{\mathfrak{u}}$, but there are no other equations or inequalities between $\mathrm{comp}_{\mathfrak{s}}$, $\mathrm{comp}_{\mathfrak{t}}$ and $\mathrm{comp}_{\mathfrak{u}}$.

Let λ be a multicomposition of n. The symmetric group $\mathfrak{S}_n$ acts from the right on the set of λ–tableaux by permuting the entries in each tableau. Let $\mathfrak{t}^\lambda$ be the λ–tableau where $1, 2, \ldots, n$ appear in order along the rows of the first component, and then along the rows

of the second component, and so on. The row stabilizer of $\mathfrak{t}^\lambda$ is the Young subgroup $\mathfrak{S}_\lambda$ of $\mathfrak{S}_n$. For example, if $\lambda = ((3,2),(1^2),(3))$ then

$$\mathfrak{t}^\lambda = \Big(\begin{array}{|c|c|c|}\hline 1&2&3\\\hline 4&5\\\cline{1-2}\end{array}\,,\quad \begin{array}{|c|}\hline 6\\\hline 7\\\cline{1-1}\end{array}\,,\quad \begin{array}{|c|c|c|}\hline 8&9&10\\\hline\end{array}\Big).$$

For a row standard λ–tableau $\mathfrak{s}$, let $d(\mathfrak{s})$ be the element of $\mathfrak{S}_n$ such that $\mathfrak{s} = \mathfrak{t}^\lambda d(\mathfrak{s})$. Then $d(\mathfrak{s})$ is a distinguished right coset representative of $\mathfrak{S}_\lambda$ in $\mathfrak{S}_n$ and we obtain, in this way, a correspondence between the set of row standard λ–tableaux and the set of right coset representatives of $\mathfrak{S}_\lambda$ in $\mathfrak{S}_n$.

Recall the antiautomorphism $*$ of $\mathscr{H}$ from (2.3).

(3.14) Definition *Suppose that λ is a multicomposition of n and that $\mathfrak{s}$ and $\mathfrak{t}$ are row standard λ–tableaux. Let $m_{\mathfrak{st}} = T^*_{d(\mathfrak{s})} m_\lambda T_{d(\mathfrak{t})}$.*

Note that $m_\lambda = m_{\mathfrak{t}^\lambda \mathfrak{t}^\lambda}$. Also, $m^*_{\mathfrak{st}} = m_{\mathfrak{ts}}$ (cf. Remark 3.7).

One of the aims of this section is to show that the elements $m_{\mathfrak{st}}$, as $(\mathfrak{s},\mathfrak{t})$ varies over the ordered pairs of standard tableaux of the same shape, give a cellular basis of $\mathscr{H}$. We shall also establish useful properties of the right ideals M^λ. Initially, though, we concentrate upon the two–sided ideal generated by m_λ.

(3.15) Lemma *Suppose that λ is a multicomposition of n and that $\mathfrak{s}$ and $\mathfrak{t}$ are row standard λ–tableaux. Let $h \in \mathscr{H}(\mathfrak{S}_n)$. Then $m_{\mathfrak{st}}h$ is a linear combination of terms of the form $m_{\mathfrak{sv}}$ where each $\mathfrak{v}$ is a row standard λ–tableau.*

PROOF: Suppose that $w \in \mathfrak{S}_n$. Then there exist $y \in \mathfrak{S}_\lambda$ and a distinguished right coset representative d of $\mathfrak{S}_\lambda$ in $\mathfrak{S}_n$ such that $w = yd$ and $\ell(w) = \ell(y) + \ell(d)$. Hence, $x_\lambda T_w = x_\lambda T_y T_d = q^{\ell(y)} x_\lambda T_d$. If $m_\lambda = u^+_{\mathbf{a}} x_\lambda$ then

$$m_{\mathfrak{st}^\lambda} T_w = T^*_{d(\mathfrak{s})} u^+_{\mathbf{a}} x_\lambda T_w = q^{\ell(y)} T^*_{d(\mathfrak{s})} u^+_{\mathbf{a}} x_\lambda T_d = q^{\ell(y)} m_{\mathfrak{sv}}$$

where the tableau $\mathfrak{v} = \mathfrak{t}^\lambda d$ is row standard.

Now, $m_{\mathfrak{st}}h = m_{\mathfrak{st}^\lambda} T_{d(\mathfrak{t})} h$, and this is a linear combination of terms of the form $m_{\mathfrak{st}^\lambda} T_w$ with $w \in \mathfrak{S}_n$. Therefore, the required result follows. □

In order to apply a result of Murphy in Proposition 3.18 below, we require a combinatorial lemma which concerns the dominance order on row standard tableaux. A preliminary definition sets the scene.

(3.16) Definition *We say that a tableau $\mathfrak{s}$ is of the* initial kind *(for λ) if* $\mathrm{comp}_{\mathfrak{s}} = \mathrm{comp}_{\mathfrak{t}^\lambda}$.

Note that a μ–tableau $\mathfrak{s}$ can be of the initial kind for λ even though $\mu \neq \lambda$.

(3.17) Lemma *Suppose that $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ is a multicomposition of n, and let $\alpha = (|\lambda^{(1)}|, \dots, |\lambda^{(r)}|)$. Suppose that w is a distinguished right coset representative of $\mathfrak{S}_\alpha$ in $\mathfrak{S}_n$ and that $\mathfrak{s}$ is a row standard λ–tableau of the initial kind. Then the following hold.*

(i) *The tableau $\mathfrak{s}w$ is row standard.*
(ii) *If $\mathfrak{s}$ is standard then $\mathfrak{s}w$ is standard.*
(iii) *If $\mathfrak{t}$ is a row standard λ–tableau of the initial kind with $\mathfrak{s} \rhd \mathfrak{t}$ then $\mathfrak{s}w \rhd \mathfrak{t}w$.*

PROOF: The fact that w is a distinguished right coset representative of $\mathfrak{S}_\alpha$ in $\mathfrak{S}_n$ implies that whenever x and y are integers such that x and y are in the same component of $\mathfrak{s}$ and $x < y$, then $xw < yw$. Hence, (i) and (ii) are true.

Now assume that $\mathfrak{t}$ is a row standard λ–tableau of the initial kind, and that $\mathfrak{s} \rhd \mathfrak{t}$. Then $d(\mathfrak{s}) \rhd d(\mathfrak{t})$ in the Bruhat–Chevalley order $\rhd$ on $\mathfrak{S}_n$, by [**4**, (1.3)]. (Our notation for the order is such that $1 \trianglerighteq v$ for all $v \in \mathfrak{S}_n$.) Since w is a distinguished right coset representative of $\mathfrak{S}_\alpha$ in $\mathfrak{S}_n$, we also have $\ell(d(\mathfrak{t})w) = \ell(d(\mathfrak{t})) + \ell(w)$. The well–known "cancellation" property of the Bruhat–Chevalley order now lets us conclude that $d(\mathfrak{s})w \rhd d(\mathfrak{t})w$. But $\mathfrak{s}w$ and $\mathfrak{t}w$ are row standard, so $d(\mathfrak{s})w = d(\mathfrak{s}w)$ and $d(\mathfrak{t})w = d(\mathfrak{t}w)$. By applying [**4**, (1.3)] again, we deduce that $\mathfrak{s}w \rhd \mathfrak{t}w$. □

(3.18) Proposition *Suppose that λ is a multicomposition of n and that $\mathfrak{s}$ and $\mathfrak{t}$ are row standard λ–tableaux. Then $m_{\mathfrak{s}\mathfrak{t}}$ is a linear combination of terms of the form $m_{\mathfrak{u}\mathfrak{v}}$ where $\mathfrak{u}$ and $\mathfrak{v}$ are standard μ–tableaux for some multipartition μ of n, and*

(i) *$\mathfrak{u} \trianglerighteq \mathfrak{s}$ and $\mathrm{comp}_{\mathfrak{u}} = \mathrm{comp}_{\mathfrak{s}}$; and,*
(ii) *$\mathfrak{v} \trianglerighteq \mathfrak{t}$ and $\mathrm{comp}_{\mathfrak{v}} = \mathrm{comp}_{\mathfrak{t}}$.*

PROOF: When $r = 1$, this is a Theorem of Murphy [**15**, (4.18)]. We deduce the general case from this.

Let $\alpha = (|\lambda^{(1)}|, \dots, |\lambda^{(r)}|)$. We may write $\mathfrak{s} = \mathfrak{s}'w_1$ and $\mathfrak{t} = \mathfrak{t}'w_2$ where $\mathfrak{s}'$ and $\mathfrak{t}'$ are row standard λ–tableau of the initial kind, and w_1 and w_2 are distinguished right coset representatives for $\mathfrak{S}_\alpha$ in $\mathfrak{S}_n$.

Let $\mathbf{a} = (a_1, \dots, a_r)$ where $a_k = \sum_{i=1}^{k-1} |\lambda^{(i)}|$, as in the definition of m_λ. We have $m_{\mathfrak{s}\mathfrak{t}} = T^*_{w_1} m_{\mathfrak{s}'\mathfrak{t}'} T_{w_2}$ and $m_{\mathfrak{s}'\mathfrak{t}'} = T^*_{d(\mathfrak{s}')} u^+_{\mathbf{a}} x_\lambda T_{d(\mathfrak{t}')} = u^+_{\mathbf{a}} T^*_{d(\mathfrak{s}')} x_\lambda T_{d(\mathfrak{t}')}$ since $d(\mathfrak{s}') \in \mathfrak{S}_\alpha$.

We may write $T^*_{d(\mathfrak{s}')} x_\lambda T_{d(\mathfrak{t}')}$ as a product of r commuting terms, one for each component of λ; say, $T^*_{d(\mathfrak{s}')} x_\lambda T_{d(\mathfrak{t}')} = x_1 x_2 \dots x_r$, where the kth term x_k involves only elements T_w with $w \in \mathfrak{S}(\{a_k + 1, a_k + 2 \dots, a_{k+1}\})$. For example, $x_1 = T^*_{d(\mathfrak{s}'_1)} x_{\lambda^{(1)}} T_{d(\mathfrak{t}'_1)}$ where $\mathfrak{s}'_1$ is the first component of $\mathfrak{s}'$ and $\mathfrak{t}'_1$ is the first component of $\mathfrak{t}'$. By applying Murphy's result [**15**, (4.18)] to the Hecke algebra $\mathscr{H}(\mathfrak{S}_{\alpha_1})$ we may write x_1 as a linear combination of terms $T^*_{d(\mathfrak{u}'_1)} x_{\mu^{(1)}} T_{d(\mathfrak{v}'_1)}$ where $\mathfrak{u}'_1$ and $\mathfrak{v}'_1$ are standard $\mu^{(1)}$–tableaux for some partition $\mu^{(1)}$ of a_1, and $\mathfrak{u}'_1 \trianglerighteq \mathfrak{s}'_1$ and $\mathfrak{v}'_1 \trianglerighteq \mathfrak{t}'_1$.

We can apply the same technique for the other factors $x_2, \dots, x_r$, to conclude that $T^*_{d(\mathfrak{s}')} x_\lambda T_{d(\mathfrak{t}')}$ is a linear combination of terms of the form $T^*_{d(\mathfrak{u}')} x_\mu T_{d(\mathfrak{v}')}$ where $\mathfrak{u}'$ and $\mathfrak{v}'$ are standard μ–tableaux for some multipartition μ of n, and $\mathfrak{u}' \trianglerighteq \mathfrak{s}'$ and $\mathfrak{v}' \trianglerighteq \mathfrak{t}'$. Also, $\mathfrak{u}'$ and $\mathfrak{v}'$

are of the initial kind, and $|\mu^{(k)}| = |\lambda^{(k)}|$ for $1 \le k \le r$. Therefore, $m_{\mathfrak{s}'\mathfrak{t}'}$ is a linear combination of terms $m_{\mathfrak{u}'\mathfrak{v}'}$ where the sum runs over the same set of pairs $(\mathfrak{u}', \mathfrak{v}')$. Consequently, $m_{\mathfrak{s}\mathfrak{t}}$ is a linear combination of terms $T^*_{w_1} m_{\mathfrak{u}'\mathfrak{v}'} T_{w_2}$.

Now, $T_{d(\mathfrak{v}')} T_{w_2} = T_{d(\mathfrak{v}')w_2} = T_{d(\mathfrak{v}'w_2)}$, so $T^*_{w_1} m_{\mathfrak{u}'\mathfrak{v}'} T_{w_2} = T^*_{w_1} m_{\mathfrak{u}'\mathfrak{v}}$ where $\mathfrak{v} = \mathfrak{v}'w_2$. By Lemma 3.17, since $\mathfrak{v}'$ is of the initial kind, $\mathfrak{v} = \mathfrak{v}'w_2$ is standard, and $\mathfrak{v}'w_2 \trianglerighteq \mathfrak{t}'w_2$; that is, $\mathfrak{v} \trianglerighteq \mathfrak{t}$. Moreover, $\mathrm{comp}_{\mathfrak{v}'w_2} = \mathrm{comp}_{\mathfrak{t}'w_2}$ since $\mathfrak{v}'$ and $\mathfrak{t}'$ are of the initial kind.

Similar remarks applied to $T^*_{w_1} T^*_{d(\mathfrak{u}')}$ now complete the proof of the Proposition. □

(3.19) Corollary *Suppose that λ is a multicomposition of n and that $\mathfrak{s}$ and $\mathfrak{t}$ are row standard λ–tableaux. If $h \in \mathscr{H}(\mathfrak{S}_n)$ then $m_{\mathfrak{s}\mathfrak{t}}h$ is a linear combination of terms of the form $m_{\mathfrak{u}\mathfrak{v}}$ where $\mathfrak{u}$ and $\mathfrak{v}$ are standard μ–tableaux for some multipartition μ of n, and $\mu \trianglerighteq \lambda$, $\mathfrak{u} \trianglerighteq \mathfrak{s}$ and* $\mathrm{comp}_{\mathfrak{u}} = \mathrm{comp}_{\mathfrak{s}}$.

PROOF: Combine Lemma 3.15 and Proposition 3.18, and recall that $\mathfrak{u} \trianglerighteq \mathfrak{s}$ implies that $\mu \trianglerighteq \lambda$. □

Corollary 3.19 provides the kind of information we need when we multiply $m_{\mathfrak{s}\mathfrak{t}}$ by elements of $\mathscr{H}(\mathfrak{S}_n)$. More complicated is our next proposition, which shows what happens when we multiply $m_{\mathfrak{s}\mathfrak{t}}$ by T_0. It is vital to the proposition that λ is a multipartition, not merely a multicomposition.

(3.20) Proposition *Suppose that λ is a multipartition of n, and that $\mathfrak{s}$ and $\mathfrak{t}$ are standard λ–tableaux. Then $m_{\mathfrak{s}\mathfrak{t}}T_0 = x_1 + x_2$ where*

(i) *x_1 is a linear combination of terms of the form $m_{\mathfrak{u}\mathfrak{v}}$ where $\mathfrak{u}$ and $\mathfrak{v}$ are standard λ–tableaux, with $\mathfrak{u} \trianglerighteq \mathfrak{s}$ and $\mathrm{comp}_{\mathfrak{u}} = \mathrm{comp}_{\mathfrak{s}}$, and*

(ii) *x_2 is a linear combination of terms of the form $m_{\mathfrak{u}\mathfrak{v}}$ where $\mathfrak{u}$ and $\mathfrak{v}$ are standard μ–tableaux for some multipartition μ of n, with $\mu \triangleright \lambda$ and $\mathrm{comp}_{\mathfrak{s}} > \mathrm{comp}_{\mathfrak{u}}$.*

PROOF: Let $\alpha = (|\lambda^{(1)}|, \dots, |\lambda^{(r)}|)$ and let $\mathbf{a} = (a_1, a_2, \dots, a_r)$ where $a_k = \sum_{i=1}^{k-1} |\lambda^{(i)}|$ for $1 \le k \le r$. Write $d(\mathfrak{t}) = yc$ with $y \in \mathfrak{S}_\alpha$ and c a distinguished right coset representative for $\mathfrak{S}_\alpha$ in $\mathfrak{S}_n$. Then the α–tableau $\mathfrak{t}^\alpha c$ is row standard. Assume that 1 is in row k of $\mathfrak{t}^\alpha c$; thus, $1 \le k \le r$.

Now, $c = (a_k, a_k+1)(a_k-1, a_k)\dots(1,2)c'$, where $\ell(c) = a_k + \ell(c')$ and c' fixes 1. Thus, $T_c = T_{a_k} T_{a_{k-1}} \dots T_1 T_{c'}$ and $T_{c'} T_0 = T_0 T_{c'}$. Let $\mathbf{b} = (a_1, \dots, a_k + 1, \dots, a_r)$. Then

$$u^+_{\mathbf{a}} T_c T_0 = u^+_{\mathbf{a}} T_{a_k} T_{a_{k-1}} \dots T_1 T_0 T_{c'} = u^+_{\mathbf{a}} h_1 + u^+_{\mathbf{b}} h_2$$

for some $h_1, h_2 \in \mathfrak{S}_n$ by Lemma 3.4. Since $y \in \mathfrak{S}_\alpha$ and $\mathfrak{t}^\lambda y$ is standard, y fixes $a_k + 1$; therefore, T_y commutes with $u^+_{\mathbf{b}}$. Hence, using Remark 3.7, we have

$$\begin{aligned} m_{\mathfrak{s}\mathfrak{t}}T_0 &= T^*_{d(\mathfrak{s})} u^+_{\mathbf{a}} x_\lambda T_{d(\mathfrak{t})} T_0 = T^*_{d(\mathfrak{s})} x_\lambda u^+_{\mathbf{a}} T_y T_c T_0 = T^*_{d(\mathfrak{s})} x_\lambda T_y u^+_{\mathbf{a}} T_c T_0 \\ &= T^*_{d(\mathfrak{s})} x_\lambda T_y (u^+_{\mathbf{a}} h_1 + u^+_{\mathbf{b}} h_2) = T^*_{d(\mathfrak{s})} x_\lambda (u^+_{\mathbf{a}} T_y h_1 + u^+_{\mathbf{b}} T_y h_2). \end{aligned}$$

The first term, $T^*_{d(\mathfrak{s})}x_\lambda u^+_{\mathbf{a}}T_y h_1$, is a linear combination of terms of the required form by Corollary 3.19. If $k = 1$ then $u^+_{\mathbf{b}} = 0$ by Remark 3.2(i), and the proof is complete in this case. Assume therefore that $k \geq 2$. We turn out attention to the term $T^*_{d(\mathfrak{s})}x_\lambda u^+_{\mathbf{b}}T_y h_2$. We now need to digress in order to prove that $T^*_{d(\mathfrak{s})}x_\lambda u^+_{\mathbf{b}}T_y h_2$ has the required form.

Define the multicomposition $\nu = (\nu^{(1)}, \dots, \nu^{(r)})$ of n by $\nu^{(i)} = \lambda^{(i)}$ for $i \neq k-1, k$, $\nu^{(k-1)} = (\lambda^{(k-1)}_1, \lambda^{(k-1)}_2, \dots, 1)$ and $\nu^{(k)} = (\lambda^{(k)}_1 - 1, \lambda^{(k)}_2, \dots)$. (Thus we reduce the first part of the kth component of λ by 1, and adjoin a new part of size 1 to the end of the $(k-1)$th component of λ.) Note that $\nu \rhd \lambda$.

Let $l = \lambda^{(k)}_1$ be the first part of $\lambda^{(k)}$. The entries in the first row of the kth component of $\mathfrak{t}^\lambda$ are then $a_k + 1, a_k + 2, \dots, a_k + l$. Let G_1 be the symmetric group on these numbers, and let G_2 be the symmetric group on $a_k + 2, \dots, a_k + l$. Let $c_1, c_2, \dots, c_l$ be the distinguished right coset representatives for G_2 in G_1, ordered in terms of increasing length. Then the tableaux $\mathfrak{t}^\nu c_i$ are row standard, and they agree with $\mathfrak{t}^\nu$ except on the last row of the $(k-1)$th component and the first row of the kth component. The last row of the $(k-1)$th component of $\mathfrak{t}^\nu c_i$ contains the single entry $a_k + i$.

We are given that $\mathfrak{s} = \mathfrak{t}^\lambda d(\mathfrak{s})$ is a row standard λ–tableau. Let $\mathfrak{u}_i = \mathfrak{t}^\nu c_i d(\mathfrak{s})$ for $1 \leq i \leq l$. Each $\mathfrak{u}_i$ agrees with $\mathfrak{s}$ except on the last row of the $(k-1)$th component of $\mathfrak{u}_i$ and on the first row of the kth component of $\mathfrak{u}_i$. Furthermore, since $(a_k + 1)d(\mathfrak{s}) < \dots < (a_k + l)d(\mathfrak{s})$, it follows that each $\mathfrak{u}_i$ is row standard.

We also have $\mathrm{comp}_{\mathfrak{u}_i}(j) = \mathrm{comp}_{\mathfrak{s}}(j)$ if $j \notin \{(a_k + 1)d(\mathfrak{s}), \dots, (a_k + l)d(\mathfrak{s})\}$ and, for $j \in \{(a_k + 1)d(\mathfrak{s}), \dots, (a_k + l)d(\mathfrak{s})\}$, we have $\mathrm{comp}_{\mathfrak{u}_i}(j) = k - 1$ or k and $\mathrm{comp}_{\mathfrak{s}}(j) = k$. Therefore, $\mathrm{comp}_{\mathfrak{s}} > \mathrm{comp}_{\mathfrak{u}_i}$.

Now x_λ has a factor

$$\sum_{w \in G_1} T_w = \sum_{i=1}^{l} T^*_{c_i} \sum_{w \in G_2} T_w.$$

Hence, $x_\lambda = \sum_{i=1}^{l} T^*_{c_i} x_\nu$. Note that $m_\nu = x_\nu u^+_{\mathbf{b}}$. Thus,

$$T^*_{d(\mathfrak{s})}x_\lambda u^+_{\mathbf{b}} = T^*_{d(\mathfrak{s})} \sum_{i=1}^{l} T^*_{c_i} x_\nu u^+_{\mathbf{b}} = \sum_{i=1}^{l} T^*_{c_i d(\mathfrak{s})} x_\nu u^+_{\mathbf{b}} = \sum_{i=1}^{l} m_{\mathfrak{u}_i \mathfrak{t}^\nu}.$$

Hence, by Corollary 3.19, $T^*_{d(\mathfrak{s})}x_\lambda u^+_{\mathbf{b}}T_y h_2$ is a linear combination of the required form.

This completes the proof of the Proposition. □

(3.21) Definition *Suppose that λ is a multicomposition of n.*

(i) *Let N^λ be the R–module spanned by*

$$\left\{ m_{\mathfrak{s}\mathfrak{t}} \;\middle|\; \begin{array}{c} \mathfrak{s} \text{ and } \mathfrak{t} \text{ are standard } \mu\text{–tableaux for some} \\ \text{multipartition } \mu \text{ of } n \text{ with } \mu \trianglerighteq \lambda \end{array} \right\}.$$

(ii) *Let* $\overline{N^\lambda}$ *be the* R*–module spanned by*

$$\left\{ m_{\mathfrak{st}} \;\middle|\; \begin{array}{c} \mathfrak{s} \text{ and } \mathfrak{t} \text{ are standard } \mu\text{–tableaux for some} \\ \text{multipartition } \mu \text{ of } n \text{ with } \mu \rhd \lambda \end{array} \right\}.$$

We now apply Propositions 3.18 and 3.20 to obtain a sequence of useful results.

(3.22) Proposition *Suppose that* λ *is a multicomposition of* n*. Then* N^λ *and* $\overline{N^\lambda}$ *are two–sided ideals of* $\mathscr{H}$*.*

PROOF: Corollary 3.19 shows that N^λ is closed under postmultiplication by $T_1, T_2, \dots, T_{n-1}$ and Proposition 3.20 shows that N^λ is also closed under postmultiplication by T_0. Because $\mathscr{H}$ is generated by $T_0, T_1, \dots, T_{n-1}$, we deduce that N^λ is a right ideal of $\mathscr{H}$. Since $m^*_{\mathfrak{st}} = m_{\mathfrak{ts}}$, by applying the antiautomorphism $*$ we see that N^λ is also a left ideal of $\mathscr{H}$.

Finally, $\overline{N^\lambda}$ is a two sided ideal of $\mathscr{H}$ since $\overline{N^\lambda} = \sum_{\mu \rhd \lambda} N^\mu$. □

A proof very similar to that of Proposition 3.22 gives our next result.

(3.23) Proposition *Suppose that* λ *is a multicomposition of* n *and that* $\mathfrak{s}$ *is a row standard* λ*–tableau. Let* $I_\mathfrak{s}$ *be the* R*–module spanned by*

$$\left\{ m_{\mathfrak{uv}} \;\middle|\; \begin{array}{c} \mathfrak{u} \text{ and } \mathfrak{v} \text{ are standard } \mu\text{–tableaux for some multipartition} \\ \mu \text{ of } n \text{ with } \mu \trianglerighteq \lambda \text{ and } \mathrm{comp}_\mathfrak{s} \geq \mathrm{comp}_\mathfrak{u} \end{array} \right\}.$$

Then $I_\mathfrak{s}$ *is a right ideal of* $\mathscr{H}$*.*

Recall that $M^\lambda = m_\lambda \mathscr{H}$.

(3.24) Corollary *Suppose that* λ *is a multicomposition of* n*. Then every element of* M^λ *is a linear combination of terms of the form* $m_{\mathfrak{uv}}$ *where* $\mathfrak{u}$ *and* $\mathfrak{v}$ *are standard* μ*–tableaux for some multipartition* μ *of* n *with* $\mu \trianglerighteq \lambda$ *and* $\mathrm{comp}_{\mathfrak{t}^\lambda} \geq \mathrm{comp}_\mathfrak{u}$*.*

PROOF: Since $m_\lambda = m_{\mathfrak{t}^\lambda \mathfrak{t}^\lambda}$, Proposition 3.18 shows that $m_\lambda \in I_{\mathfrak{t}^\lambda}$ (note that λ may not be a multipartition). Therefore, $M^\lambda \subseteq I_{\mathfrak{t}^\lambda}$ and the desired result follows by Proposition 3.23. □

(3.25) Proposition *Suppose that* λ *is a multipartition of* n *and that* $\mathfrak{t}$ *is a standard* λ*–tableau. Let* $h \in \mathscr{H}$*. Then for every standard* λ*–tableau* $\mathfrak{v}$ *there exists* $r_\mathfrak{v} \in R$ *such that, for all standard* λ*–tableau* $\mathfrak{s}$*, we have*

$$m_{\mathfrak{st}} h \equiv \sum_{\mathfrak{v} \in \mathrm{Std}(\lambda)} r_\mathfrak{v} m_{\mathfrak{sv}} \bmod \overline{N^\lambda}.$$

PROOF: Let U be the R–module spanned by $\{ m_{\mathfrak{t}^\lambda \mathfrak{t}'} \mid \mathfrak{t}' \text{ is a standard } \lambda\text{–tableau} \}$. Note that $\mathfrak{t}^\lambda$ is the unique λ–tableau $\mathfrak{u}$ such that $\mathfrak{u} \trianglerighteq \mathfrak{t}^\lambda$. Therefore, $U + \overline{N^\lambda}$ is closed under postmultiplication by $T_1, T_2, \dots, T_{n-1}$ by Proposition 3.18, and it is closed under postmultiplication by T_0 by Proposition 3.20. Hence $U + \overline{N^\lambda}$ is a right ideal of $\mathscr{H}$. Thus, for each $\mathfrak{v} \in \mathrm{Std}(\lambda)$ there exists $r_\mathfrak{v} \in R$ such that

$$m_{\mathfrak{t}^\lambda \mathfrak{t}} h \equiv \sum_{\mathfrak{v} \in \mathrm{Std}(\lambda)} r_\mathfrak{v} m_{\mathfrak{t}^\lambda \mathfrak{v}} \bmod \overline{N^\lambda}.$$

Multiply this congruence on the left by $T^*_{d(\mathfrak{s})}$, and use the fact that $\overline{N^\lambda}$ is an ideal, to obtain the congruence of the proposition. □

(3.26) Theorem ([**13**, (1.7)]) *The algebra $\mathscr{H}$ is a free R–module with basis*

$$\mathscr{M} = \left\{ m_{\mathfrak{s}\mathfrak{t}} \;\middle|\; \begin{array}{c} \mathfrak{s} \text{ and } \mathfrak{t} \text{ are standard } \lambda\text{–tableaux for some} \\ \text{multipartition } \lambda \text{ of } n \end{array} \right\}.$$

Moreover, $\mathscr{M}$ is a cellular basis of $\mathscr{H}$.

PROOF: Since $m_\lambda = 1$ when $\lambda = ((0), \dots, (0), (1^n))$, Proposition 3.25 shows that $\mathscr{M}$ spans $\mathscr{H}$. By Theorem 2.2, $\mathscr{H}$ is free of rank $r^n n!$ Since this is also the cardinality of $\mathscr{M}$, we deduce that $\mathscr{M}$ is a basis of $\mathscr{H}$. Finally, the properties that $\mathscr{M}$ must satisfy in order to be a cellular basis of $\mathscr{H}$ (as given in [**12**, (1.1)]), are covered by Proposition 3.25 and the fact that $m^*_{\mathfrak{s}\mathfrak{t}} = m_{\mathfrak{t}\mathfrak{s}}$. □

(3.27) Definition *We call $\mathscr{M}$ the* standard basis *of $\mathscr{H}$.*

We can now apply Graham and Lehrer's theory of cellular algebras [**12**] to describe the representation theory of $\mathscr{H}$.

(3.28) Definition *Suppose that λ is a multipartition of n. Let $z_\lambda = (\overline{N^\lambda} + m_\lambda)/\overline{N^\lambda}$. The* Specht module *$S^\lambda$ is the submodule of $\mathscr{H}/\overline{N^\lambda}$ given by $S^\lambda = z_\lambda \mathscr{H}$.*

As in [**12**], or directly from Proposition 3.25 and Theorem 3.26, S^λ is a free R–module with basis $\{ z_\lambda T_{d(\mathfrak{t})} \mid \mathfrak{t} \text{ a standard } \lambda\text{–tableau} \}$.

Define a bilinear form $\langle\ ,\ \rangle$ on the Specht module S^λ by

$$m_\lambda T_{d(\mathfrak{s})} T^*_{d(\mathfrak{t})} m_\lambda \equiv \langle z_\lambda T_{d(\mathfrak{s})}, z_\lambda T_{d(\mathfrak{t})} \rangle m_\lambda \bmod \overline{N^\lambda}.$$

By Proposition 3.25 (and the version of Proposition 3.25 obtained by applying the antiautomorphism $*$), the bilinear form is well defined (cf. [**12**]). Moreover, the bilinear form is symmetric and $\langle uh, v \rangle = \langle u, vh^* \rangle$ for all $u, v \in S^\lambda$ and all $h \in \mathscr{H}$ [**12**, (2.4)]. Consequently, $\operatorname{rad} S^\lambda$, the radical of the bilinear form, is an $\mathscr{H}$–module.

(3.29) Definition *Suppose that λ is a multipartition of n. Let $D^\lambda = S^\lambda/\operatorname{rad} S^\lambda$.*

(3.30) Theorem *Suppose that R is a field. Then the non–zero $\mathscr{H}$ modules in*

$$\{\, D^\lambda \mid \lambda \text{ a multipartition of } n \,\}$$

form a complete set of non–isomorphic irreducible $\mathscr{H}$–modules. Moreover, each irreducible module D^λ is absolutely irreducible.

PROOF: Since $\mathscr{M}$ is a cellular basis of $\mathscr{H}$ the Theorem follows from the general theory of cellular algebras [**12**, (3.4)]. □

The theory in [**12**] also shows that if $D^\mu \neq 0$ and D^μ is a composition factor of S^λ then $\lambda \trianglerighteq \mu$.

When $r = 1$ the partitions μ for which $D^\mu \neq 0$ have been classified in [**4**]. When $r > 1$, if $q = 1$, or if the parameters Q_k are powers of q, the multipartitions μ for which $D^\mu \neq 0$ are classified by the results of [**12**, **13**, **14**]; in type **B**, see also [**8**, **10**].

4 A Specht series for M^μ

In the last section, we used various elements m_μ of $\mathscr{H}$ to construct a cellular basis of $\mathscr{H}$. We saw that the right ideal M^μ generated by m_μ is an analogue of an induced module; indeed, if $r = 1$ and $q = 1$ then M^μ is the permutation module of $\mathfrak{S}_n$ on the Young subgroup $\mathfrak{S}_\mu$. We shall use the right ideals M^μ of $\mathscr{H}$ to construct the cyclotomic q–Schur algebra, but, before that, we study the individual modules M^μ in more detail. In particular, we shall give a semistandard basis of M^μ and construct a Specht series for M^μ.

We are going to define a new kind of λ–tableau. This will have entries which are ordered pairs (i, k) where i is a positive integer and $1 \leq k \leq r$. We denote such tableaux by capital letters.

(4.1) Definition *Suppose that λ is a multipartition of n and that μ is a multicomposition of n. A λ–tableau $\mathtt{S}$ has* type *μ if its entries are ordered pairs (i, k), as above, and for all m and k the number of times (i, k) is an entry in $\mathtt{S}$ is $\mu_i^{(k)}$.*

Next, we introduce a function which converts standard λ–tableaux (or, indeed any tableau whose entries are $1, 2, \ldots, n$) into a λ–tableau of type μ.

(4.2) Definition *Suppose that λ is a multipartition of n and that μ is a multicomposition of n. Let $\mathfrak{s}$ be a standard λ–tableau. Define $\mu(\mathfrak{s})$ to be the λ–tableau obtained from $\mathfrak{s}$ by replacing each entry m in $\mathfrak{s}$ by (i, k) if m is in row i of the kth component of $\mathfrak{t}^\mu$.*

In our examples, we shall always write i_k for the ordered pair (i, k).

(4.3) Example Suppose that $\mu = ((3,1),(2),(2,1^2))$. Then

$$\mathrm{t}^\mu = \left(\begin{array}{|c|c|c|}\hline 1&2&3\\\hline 4\\\cline{1-1}\end{array}\,,\quad \begin{array}{|c|c|}\hline 5&6\\\hline\end{array}\,,\quad \begin{array}{|c|c|}\hline 7&8\\\hline 9\\\cline{1-1}10\\\cline{1-1}\end{array}\right) \quad\text{and}\quad \mu(\mathrm{t}^\mu) = \left(\begin{array}{|c|c|c|}\hline 1_1&1_1&1_1\\\hline 2_1\\\cline{1-1}\end{array}\,,\quad \begin{array}{|c|c|}\hline 1_2&1_2\\\hline\end{array}\,,\quad \begin{array}{|c|c|}\hline 1_3&1_3\\\hline 2_3\\\cline{1-1}3_3\\\cline{1-1}\end{array}\right).$$

Suppose that

$$\mathfrak{s}_1 = \left(\begin{array}{|c|c|c|c|}\hline 1&2&3&5\\\hline 4\\\cline{1-1}\end{array}\,,\quad \begin{array}{|c|c|}\hline 6&10\\\hline\end{array}\,,\quad \begin{array}{|c|c|}\hline 7&8\\\hline 9\\\cline{1-1}\end{array}\right), \qquad \mathfrak{s}_2 = \left(\begin{array}{|c|c|c|c|}\hline 1&2&3&6\\\hline 4\\\cline{1-1}\end{array}\,,\quad \begin{array}{|c|c|}\hline 5&10\\\hline\end{array}\,,\quad \begin{array}{|c|c|}\hline 7&8\\\hline 9\\\cline{1-1}\end{array}\right)$$

and

$$\mathfrak{s}_3 = \left(\begin{array}{|c|c|c|c|}\hline 1&2&5&9\\\hline 7\\\cline{1-1}\end{array}\,,\quad \begin{array}{|c|c|}\hline 3&10\\\hline\end{array}\,,\quad \begin{array}{|c|c|}\hline 4&6\\\hline 8\\\cline{1-1}\end{array}\right).$$

Then

$$\mu(\mathfrak{s}_1) = \mu(\mathfrak{s}_2) = \left(\begin{array}{|c|c|c|c|}\hline 1_1&1_1&1_1&1_2\\\hline 2_1\\\cline{1-1}\end{array}\,,\quad \begin{array}{|c|c|}\hline 1_2&3_3\\\hline\end{array}\,,\quad \begin{array}{|c|c|}\hline 1_3&1_3\\\hline 2_3\\\cline{1-1}\end{array}\right)$$

and

$$\mu(\mathfrak{s}_3) = \left(\begin{array}{|c|c|c|c|}\hline 1_1&1_1&1_2&2_3\\\hline 1_3\\\cline{1-1}\end{array}\,,\quad \begin{array}{|c|c|}\hline 1_1&3_3\\\hline\end{array}\,,\quad \begin{array}{|c|c|}\hline 2_1&1_2\\\hline 1_3\\\cline{1-1}\end{array}\right).$$

Given two pairs (i_1,k_1) and (i_2,k_2), we write $(i_1,k_1) < (i_2,k_2)$ if $k_1 < k_2$, or $k_1 = k_2$ and $i_1 < i_2$. Note that if $\mathfrak{s}$ is a standard λ–tableau, then $\mu(\mathfrak{s})$ is a λ–tableau of type μ whose entries are weakly increasing along rows and down columns. We next single out some of these tableaux $\mu(\mathfrak{s})$.

(4.4) Definition *Suppose that λ is a multipartition of n and that μ is a multicomposition of n. Let $\mathtt{S} = (\mathtt{S}^{(1)},\dots,\mathtt{S}^{(r)})$ be a λ–tableau of type μ. Then $\mathtt{S}$ is* semistandard *if*

(i) *the entries in each row of each component $\mathtt{S}^{(k)}$ of $\mathtt{S}$ are non–decreasing; and*

(ii) *the entries in each column of each component $\mathtt{S}^{(k)}$ of $\mathtt{S}$ are strictly increasing; and*

(iii) *for each k with $1 \le k \le r$, no entry in $\mathtt{S}^{(k)}$ has the form (i,l) with $l < k$.*

Let $\mathcal{T}_0(\lambda,\mu)$ denote the set of semistandard λ–tableaux of type μ.

For example, $\mu(\mathfrak{s}_1)$ in Example 4.3 is semistandard, but $\mu(\mathfrak{s}_3)$ is not, since the third part of Definition 4.4 is not satisfied.

The point of the third condition in Definition 4.4 is the following result (cf. Corollary 3.24).

(4.5) Suppose that $\mathfrak{s}$ is a standard λ–tableau. Then $\mu(\mathfrak{s})$ satisfies Definition 4.4(iii) if and only if $\mathrm{comp}_{\mathrm{t}^\mu} \ge \mathrm{comp}_{\mathfrak{s}}$

If λ is a multipartition and $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$, then it is clear that there exists a standard λ–tableau $\mathfrak{s}$ with $\mathtt{S} = \mu(\mathfrak{s})$. In order to say more about this, we introduce another definition.

(4.6) Definition *If $\mathfrak{s}$ is a standard λ–tableau and $1 \le m \le n$ then let $\mathrm{row}_{\mathfrak{s}}(m) = (i,k)$ if m belongs to row i of the kth component of $\mathfrak{s}$.*

The definition of $\mathrm{row}_{\mathfrak{s}}$ allows us to recover $\mathfrak{s}$ from the function $\mathrm{row}_{\mathfrak{s}}$

Suppose that $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$ and let $\mathfrak{s}$ be a standard λ–tableau such that $\mu(\mathfrak{s}) = \mathtt{S}$. If there exists an integer i such that $\mathrm{row}_{\mathfrak{s}}(i) \neq \mathrm{row}_{\mathfrak{s}}(i+1)$ and i and $i+1$ are in the same row of $\mathfrak{t}^{\mu}$ then the tableau $\mathfrak{s}_1 = \mathfrak{s}(i,i+1)$ is standard and $\mu(\mathfrak{s}_1) = \mathtt{S}$. Also, $\mathfrak{s} \rhd \mathfrak{s}_1$ if $\mathrm{row}_{\mathfrak{s}}(i) < \mathrm{row}_{\mathfrak{s}}(i+1)$, and $\mathfrak{s}_1 \rhd \mathfrak{s}$ if $\mathrm{row}_{\mathfrak{s}}(i) > \mathrm{row}_{\mathfrak{s}}(i+1)$. Hence we have shown the following.

(4.7) Suppose that $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$. Then there exist standard λ–tableaux $\mathrm{first}(\mathtt{S})$ and $\mathrm{last}(\mathtt{S})$ such that

(i) $\mu(\mathrm{first}(\mathtt{S})) = \mu(\mathrm{last}(\mathtt{S})) = \mathtt{S}$; and,

(ii) if $\mathfrak{s}$ is any standard λ–tableau such that $\mu(\mathfrak{s}) = \mathtt{S}$ then $\mathrm{first}(\mathtt{S}) \trianglerighteq \mathfrak{s} \trianglerighteq \mathrm{last}(\mathtt{S})$.

(4.8) Definition *Suppose that λ is a multipartition of n and that μ is a multicomposition of n. Let $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$ and $\mathfrak{t} \in \mathrm{Std}(\lambda)$. Then $m_{\mathtt{S}\mathfrak{t}}$ is the element of $\mathscr{H}$ given by*

$$m_{\mathtt{S}\mathfrak{t}} = \sum_{\substack{\mathfrak{s}\in\mathrm{Std}(\lambda)\\ \mu(\mathfrak{s})=\mathtt{S}}} m_{\mathfrak{s}\mathfrak{t}}.$$

Note that $m_{\mathtt{S}\mathfrak{t}}$ is a sum of standard basis elements $m_{\mathfrak{s}\mathfrak{t}}$ where $\mathrm{first}(\mathtt{S}) \trianglerighteq \mathfrak{s} \trianglerighteq \mathrm{last}(\mathtt{S})$.

(4.9) Example Suppose that $\lambda = ((4,3),(2,1),(2,1))$, $\mu = ((3^2,1),(1^2),(2,1^2))$ and

$$\mathtt{S} = \Big(\begin{array}{|c|c|c|c|}\hline 1_1 & 1_1 & 1_1 & 2_1 \\ \hline 2_1 & 2_1 & 3_1 \\ \cline{1-3}\end{array}\ , \quad \begin{array}{|c|c|}\hline 1_2 & 3_3 \\ \hline 2_2 \\ \cline{1-1}\end{array}\ , \quad \begin{array}{|c|c|}\hline 1_3 & 1_3 \\ \hline 2_3 \\ \cline{1-1}\end{array} \Big).$$

Then $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$ and for any $\mathfrak{t} \in \mathrm{Std}(\lambda)$ we have $m_{\mathtt{S}\mathfrak{t}} = m_{\mathfrak{s}_1\mathfrak{t}} + m_{\mathfrak{s}_2\mathfrak{t}} + m_{\mathfrak{s}_3\mathfrak{t}}$ where

$$\mathfrak{s}_1 = \Big(\begin{array}{|c|c|c|c|}\hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & 7 \\ \cline{1-3}\end{array}\ , \quad \begin{array}{|c|c|}\hline 8 & 13 \\ \hline 9 \\ \cline{1-1}\end{array}\ , \quad \begin{array}{|c|c|}\hline 10 & 11 \\ \hline 12 \\ \cline{1-1}\end{array} \Big), \quad \mathfrak{s}_2 = \Big(\begin{array}{|c|c|c|c|}\hline 1 & 2 & 3 & 5 \\ \hline 4 & 6 & 7 \\ \cline{1-3}\end{array}\ , \quad \begin{array}{|c|c|}\hline 8 & 13 \\ \hline 9 \\ \cline{1-1}\end{array}\ , \quad \begin{array}{|c|c|}\hline 10 & 11 \\ \hline 12 \\ \cline{1-1}\end{array} \Big)$$

and

$$\mathfrak{s}_3 = \Big(\begin{array}{|c|c|c|c|}\hline 1 & 2 & 3 & 6 \\ \hline 4 & 5 & 7 \\ \cline{1-3}\end{array}\ , \quad \begin{array}{|c|c|}\hline 8 & 13 \\ \hline 9 \\ \cline{1-1}\end{array}\ , \quad \begin{array}{|c|c|}\hline 10 & 11 \\ \hline 12 \\ \cline{1-1}\end{array} \Big).$$

Here $\mathfrak{s}_1 = \mathrm{first}(\mathtt{S})$ and $\mathfrak{s}_3 = \mathrm{last}(\mathtt{S})$.

(4.10) Lemma *Suppose that λ is a multipartition of n and that μ is a multicomposition of n. Let $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$ and $\mathfrak{t} \in \mathrm{Std}(\lambda)$. Then $m_{\mathtt{S}\mathfrak{t}} \in M^{\mu}$.*

PROOF: Let $\mathbf{a} = (a_1,\dots,a_r)$ where $a_k = \sum_{i=1}^{k-1} |\mu^{(i)}|$, and $\mathbf{b} = (b_1,\dots,b_r)$ where $b_k = \sum_{i=1}^{k-1} |\lambda^{(i)}|$.

Let $\mathfrak{s}_1 = \text{first}(\mathtt{S})$ and let $d = d(\mathfrak{s}_1)$. Then, as in [4],

$$\sum_{\substack{\mathfrak{s}\in\mathrm{Std}(\lambda)\\ \mu(\mathfrak{s})=\mathtt{S}}} x_\lambda T_{d(\mathfrak{s})} = \sum_{w\in\mathfrak{S}_\lambda d\mathfrak{S}_\mu} T_w = hT_d x_\mu,$$

where $h = \sum T_v$, the sum running over certain elements $v \in \mathfrak{S}_\lambda$. Since $h \in \mathscr{H}(\mathfrak{S}_\lambda)$, we have $h^* u_{\mathbf{b}}^+ = u_{\mathbf{b}}^+ h^*$; so we obtain

$$m_{\mathfrak{St}} = \sum_{\substack{\mathfrak{s}\in\mathrm{Std}(\lambda)\\ \mu(\mathfrak{s})=\mathtt{S}}} m_{\mathfrak{st}} = \sum_{\substack{\mathfrak{s}\in\mathrm{Std}(\lambda)\\ \mu(\mathfrak{s})=\mathtt{S}}} T^*_{d(\mathfrak{s})} x_\lambda u_{\mathbf{b}}^+ T_{d(\mathfrak{t})} = x_\mu T_d^* u_{\mathbf{b}}^+ h^* T_{d(\mathfrak{t})}.$$

Therefore, since $m_\mu = x_\mu u_{\mathbf{a}}^+$, it is sufficient to prove that $T_d^* u_{\mathbf{b}}^+ \in u_{\mathbf{a}}^+ \mathscr{H}$.

Assume that $1 \le k \le r$. Recall that $\mathfrak{s}_1 = \text{first}(\mathtt{S})$, $a_k = \sum_{i=1}^{k-1} |\mu^{(i)}|$ and $b_k = \sum_{i=1}^{k-1} |\lambda^{(i)}|$ and note that $a_k \le b_k$ since $\mathtt{S}$ is semistandard. Define $\mathfrak{t}_k$ to be the standard λ–tableau such that $1, 2, \dots, a_k$ occupy the same positions in $\mathfrak{s}_1$ and $\mathfrak{t}_k$; $b_k + 1, \dots, n$ occupy the same positions in $\mathfrak{t}^\lambda$ and $\mathfrak{t}_k$; and that for $a_k + 1 \le i < j \le b_k$ we have $\mathrm{row}_{\mathfrak{t}_k}(i) \le \mathrm{row}_{\mathfrak{t}_k}(j)$ — thus the numbers $a_k+1, \dots, b_k$ are inserted into $\mathfrak{t}_k$ in "row order". The fact that $\mathtt{S}$ is semistandard ensures that $\mathfrak{t}_k$ is well defined. Note that $\mathfrak{t}_1 = \mathfrak{t}^\lambda$ and $\mathfrak{t}_r = \mathfrak{s}_1$.

Now define $w_k \in \mathfrak{S}_n$ inductively by requiring that $\mathfrak{t}_k = \mathfrak{t}^\lambda w_1 w_2 ... w_k$. Then $w_1 = 1$ and $d = w_1 w_2 ... w_r$; moreover, $\ell(d) = \ell(w_1) + ... + \ell(w_r)$. By construction, the element w_k fixes each of the numbers $1, 2, \dots, a_{k-1}$ and it also fixes each of $b_k + 1, b_k + 2, \dots, n$.

For $1 \le k \le r$, let $u_{\mathbf{a},k}$ and $u_{\mathbf{b},k}$ be as defined in Definition 3.1, and let $v_k = \prod_{m=a_k+1}^{b_k} (L_m - Q_k)$. Then $u_{\mathbf{b},k} = u_{\mathbf{a},k} v_k$.

Now, T_{w_k} commutes with $u_{\mathbf{a},l}$ for $l < k$ (since w_k fixes $1, 2, \dots, a_{k-1}$), and T_{w_k} commutes with $u_{\mathbf{b},l}$ for $l \ge k$ (since w_k fixes $b_k + 1, b_k + 2, \dots, n$); see (2.1)(iv). Therefore,

$$\begin{aligned}
T_d^* u_{\mathbf{b}}^+ &= T^*_{w_r} \dots T^*_{w_1} u_{\mathbf{b},1} u_{\mathbf{b},2} \dots u_{\mathbf{b},r} \\
&= T^*_{w_r} \dots T^*_{w_2} u_{\mathbf{b},1} u_{\mathbf{b},2} \dots u_{\mathbf{b},r} T^*_{w_1} \\
&= u_{\mathbf{a},1} T^*_{w_r} \dots T^*_{w_2} u_{\mathbf{b},2} \dots u_{\mathbf{b},r} v_1 T^*_{w_1} \\
&= u_{\mathbf{a},1} u_{\mathbf{a},2} T^*_{w_r} \dots T^*_{w_3} u_{\mathbf{b},3} \dots u_{\mathbf{b},r} v_2 T^*_{w_2} v_1 T^*_{w_1} \\
&\vdots \qquad\qquad \vdots \\
&= u_{\mathbf{a},1} \dots u_{\mathbf{a},r} v_r T^*_{w_r} v_{r-1} T^*_{w_{r-1}} \dots v_2 T^*_{w_2} v_1 T^*_{w_1}.
\end{aligned}$$

Thus, $T_d^* u_{\mathbf{b}}^+ \in u_{\mathbf{a}}^+ \mathscr{H}$, as required. □

Having shown that $m_{\mathfrak{St}} \in M^\mu$, our next aim is to prove that the set of elements of the form $m_{\mathfrak{St}}$ give a basis of M^μ.

Let $\mathbf{a} = (a_1, \dots, a_r)$ where $a_k = \sum_{i=1}^{k-1} |\mu^{(i)}|$ as in the previous proof; then $m_\mu = u_{\mathbf{a}}^+ x_\mu$. Since $u_{\mathbf{a}}^+$ and x_μ commute, we have $M^\mu = m_\mu \mathscr{H} \subseteq u_{\mathbf{a}}^+ \mathscr{H} \cap x_\mu \mathscr{H}$. We shall show that we have equality here.

We have seen that $\mathscr{H}$ has a standard basis which consists of elements $m_{\mathfrak{st}}$ with $\mathfrak{s}$ and $\mathfrak{t}$ standard λ–tableaux for some multipartition λ of n. Suppose that $h \in \mathscr{H}$ and let $h =$

$\sum_{\mathfrak{s},\mathfrak{t}} r_{\mathfrak{s}\mathfrak{t}} m_{\mathfrak{s}\mathfrak{t}}$, with $r_{\mathfrak{s}\mathfrak{t}} \in R$, be the unique expression for h in terms of the standard basis. We say that $m_{\mathfrak{s}\mathfrak{t}}$ is involved in h if $r_{\mathfrak{s}\mathfrak{t}} \neq 0$.

(4.11) Lemma *Suppose that μ is a multipartition of n and that $m_\mu = u_{\mathbf{a}}^+ x_\mu$. Let $h \in x_\mu \mathscr{H} \cap u_{\mathbf{a}}^+ \mathscr{H}$ and suppose that $(\mathfrak{s},\mathfrak{t})$ is a pair of standard tableaux of the same shape such that*

(i) *$m_{\mathfrak{s}\mathfrak{t}}$ is involved in h; and,*

(ii) *if $(\mathfrak{u},\mathfrak{v})$ is a pair of standard tableaux of the same shape such that $\mathfrak{s} \trianglerighteq \mathfrak{u}$, $\mathfrak{t} \trianglerighteq \mathfrak{v}$ and $(\mathfrak{s},\mathfrak{t}) \neq (\mathfrak{u},\mathfrak{v})$ then $m_{\mathfrak{u}\mathfrak{v}}$ is not involved in h.*

Let $\mathtt{S} = \mu(\mathfrak{s})$. Then $\mathtt{S}$ is semistandard and $\mathfrak{s} = \mathrm{last}(\mathtt{S})$.

PROOF: Since $\mathfrak{s}$ is standard, the entries in $\mathtt{S}$ are non–decreasing down rows and columns.

Suppose that i and $i+1$ belong to the same row of $\mathfrak{t}^\mu$. Then $T_i x_\mu = q x_\mu$, so $T_i h = qh$. Therefore, by [**16**, (4.19)], i and $i+1$ do not belong to the same column of $\mathfrak{s}$. Hence the entries in $\mathtt{S}$ are strictly increasing down columns.

Since $h \in u_{\mathbf{a}}^+ \mathscr{H}$ we have $\mathrm{comp}_{\mathfrak{t}^\mu} \geq \mathrm{comp}_{\mathfrak{s}}$ by Corollary 3.24 (applied to the multicomposition $((1^{\alpha_1}),(1^{\alpha_2}),\dots,(1^{\alpha_r}))$, where $\alpha = (|\mu^{(1)}|,\dots,|\mu^{(r)}|)$). Hence, by (4.5), $\mathtt{S}$ is semistandard.

Finally, suppose that $\mathfrak{s} \neq \mathrm{last}(\mathtt{S})$. Then there exist integers i and $i+1$ in the same row of $\mathfrak{t}^\mu$ such that the tableau $\mathfrak{s}' = \mathfrak{s}(i,i+1)$ is standard and $\mathfrak{s} \rhd \mathfrak{s}'$. As in [**16**, (4.19)], $m_{\mathfrak{s}'\mathfrak{t}}$ is involved in $T_i h = qh$, contradicting part (ii) of our hypothesis. Therefore, $\mathfrak{s} = \mathrm{last}(\mathtt{S})$. □

(4.12) Corollary *Suppose that μ is a multicomposition of n and $m_\mu = u_{\mathbf{a}}^+ x_\mu$. Then $u_{\mathbf{a}}^+ \mathscr{H} \cap x_\mu \mathscr{H}$ is spanned by*

$$\left\{ m_{\mathtt{S}\mathfrak{t}} \;\middle|\; \begin{array}{c} \mathtt{S} \in \mathcal{T}_0(\lambda,\mu) \text{ and } \mathfrak{t} \in \mathrm{Std}(\lambda) \text{ for some} \\ \text{multipartition } \lambda \text{ of } n \end{array} \right\}.$$

PROOF: Note that $m_{\mathtt{S}\mathfrak{t}} \in M^\mu$ by Lemma 4.10.

For each multipartition λ of n let

$$\mathcal{T}_l(\lambda,\mu) = \{\, \mathfrak{s} \in \mathrm{Std}(\lambda) \mid \mu(\mathfrak{s}) = \mathrm{last}(\mathtt{S}) \text{ for some } \mathtt{S} \in \mathcal{T}_0(\lambda,\mu) \,\}.$$

By Lemma 4.11 every non–zero element of $u_{\mathbf{a}}^+ \mathscr{H} \cap x_\mu \mathscr{H}$ involves a standard basis element $m_{\mathfrak{s}\mathfrak{t}}$ where $\mathfrak{s} \in \mathcal{T}_l(\lambda,\mu)$ and $\mathfrak{t} \in \mathrm{Std}(\lambda)$ for some multipartition λ.

Now suppose that $h \in u_{\mathbf{a}}^+ \mathscr{H} \cap x_\mu \mathscr{H}$ and write h in terms of the standard basis; say, $h = \sum r_{\mathfrak{u}\mathfrak{v}} m_{\mathfrak{u}\mathfrak{v}}$ with $r_{\mathfrak{u}\mathfrak{v}} \in R$. Let

$$h' = h - \sum_\lambda \sum_{\mathfrak{s} \in \mathcal{T}_l(\lambda,\mu)} \sum_{\mathfrak{t} \in \mathrm{Std}(\lambda)} r_{\mathfrak{s}\mathfrak{t}} m_{\mu(\mathfrak{s})\mathfrak{t}}.$$

Then $h' \in u_{\mathbf{a}}^+ \mathscr{H} \cap x_\mu \mathscr{H}$, but h' does not involve any term $m_{\mathfrak{s}\mathfrak{t}}$ for any $\mathfrak{s} \in \mathcal{T}_l(\lambda,\mu)$. Therefore, $h' = 0$, and we have obtained an expression for h as a linear combination of the required form. □

(4.13) Corollary *Suppose that μ is a multicomposition of n and $m_\mu = u_{\mathbf{a}}^+ x_\mu$. Then $M^\mu = u_{\mathbf{a}}^+ \mathscr{H} \cap x_\mu \mathscr{H}$.*

PROOF: We have that $M^\mu \subseteq u_{\mathbf{a}}^+ \mathscr{H} \cap x_\mu \mathscr{H}$. The inclusion $u_{\mathbf{a}}^+ \mathscr{H} \cap x_\mu \mathscr{H} \subseteq M^\mu$ follows from Corollary 4.12 and Lemma 4.10 □

The next theorem for Iwahori–Hecke algebras of type **A** (that is, the case $r = 1$), is due to Murphy [**16**]; for Iwahori–Hecke algebras of type **B** (that is, the case $r = 2$), the result is due to Du and Scott [**11**]

(4.14) Theorem *Suppose that μ is a multicomposition of n. Then M^μ is free as an R–module with basis*

$$\left\{ m_{\mathtt{st}} \;\middle|\; \begin{array}{c} \mathtt{S} \in \mathcal{T}_0(\lambda,\mu) \text{ and } \mathtt{t} \in \mathrm{Std}(\lambda) \text{ for} \\ \text{some multipartition } \lambda \text{ of } n \end{array} \right\}.$$

PROOF: By Lemma 4.10 each element $m_{\mathtt{st}}$ belongs to M^μ. Since distinct elements $m_{\mathtt{st}}$ involve distinct standard basis elements $m_{\mathtt{st}}$, the elements $m_{\mathtt{st}}$ are linearly independent. Finally, Corollaries 4.12 and 4.13 show that the elements $m_{\mathtt{st}}$ span M^μ. □

(4.15) Corollary *Suppose that μ is a multicomposition of n. Then there exists a filtration of M^μ,*

$$M^\mu = M_1 > M_2 > \cdots > M_{k+1} = 0$$

such that for each i with $1 \le i \le k$ there exists a multipartition λ_i of n with $M_i/M_{i+1} \cong S^{\lambda_i}$. Moreover, if λ is a multipartition of n, then the number of factors S^{λ_i} isomorphic to S^λ is equal to the number of semistandard λ–tableaux of type μ.

PROOF: Choose an ordering $\mathtt{S}_1 > \mathtt{S}_2 > \cdots > \mathtt{S}_k$ on the set of semistandard tableaux of type μ such that $j > i$ if $\lambda_i \rhd \lambda_j$ where $\mathtt{S}_i \in \mathcal{T}_0(\lambda_i,\mu)$ and $\mathtt{S}_j \in \mathcal{T}_0(\lambda_j,\mu)$. Let M_i be the R–submodule of M^μ with basis $\{ m_{\mathtt{s}_j\mathtt{t}} \mid j \ge i \text{ and } \mathtt{t} \in \mathrm{Std}(\lambda_j) \}$. Then

$$M^\mu = M_1 > M_2 > \cdots > M_{k+1} = 0,$$

and, by Proposition 3.25 and Theorem 4.14, each M_i is a right ideal of $\mathscr{H}$.

Suppose that $1 \le i \le k$. Then $M_i \cap \overline{N^{\lambda_i}} \subseteq M_{i+1}$. Hence, there is a well defined $\mathscr{H}$–homomorphism θ from S^{λ_i} onto M_i/M_{i+1} such that $\theta(z_{\lambda_i}) = m_{\mathtt{s}_i\mathtt{t}^{\lambda_i}} + M_{i+1}$. Since both S^{λ_i} and M_i/M_{i+1} have rank equal to the number of standard λ_i–tableaux, θ is an isomorphism. The Corollary now follows. □

5 The double annihilator of m_μ

The purpose of this section is to compute the double annihilator of the element m_μ of $\mathscr{H}$ for any multicomposition μ of n. This will enable us to calculate a basis for $\mathrm{Hom}_{\mathscr{H}}(M^\nu, M^\mu)$ in the next section.

(5.1) Definition *Suppose S is a subset of $\mathscr{H}$ and define $\mathrm{r}(S) = \{\, h \in \mathscr{H} \mid Sh = 0 \,\}$ and $\mathrm{l}(S) = \{\, h \in \mathscr{H} \mid hS = 0 \,\}$. The* double annihilator *of S is $\mathrm{lr}(S) = \mathrm{l}(\mathrm{r}(S))$.*

It is elementary that $\mathrm{lr}(S)$ contains the left ideal of $\mathscr{H}$ generated by S. If $\mathscr{H}$ is a quasi–Frobenius algebra, then $\mathrm{lr}(S)$ is equal to the left ideal of $\mathscr{H}$ generated by S [**3**, (61.2)]. Although $\mathscr{H}$ is quasi–Frobenius for $r \leq 2$ (see, for example, [**4**, section 2]), we do not know whether it is quasi–Frobenius for $r > 2$. If $\mathscr{H}$ is quasi–Frobenius for all r, then the main result of this section, namely Theorem 5.16, is immediate.

A connection between double annihilators and homomorphisms is provided by the following easy lemma.

(5.2) Lemma *Suppose that $m \in \mathscr{H}$ and that $\mathrm{lr}(m) = \mathscr{H}m$. Let I be a right ideal of $\mathscr{H}$.*
(i) *For all $\varphi \in \mathrm{Hom}_{\mathscr{H}}(m\mathscr{H}, I)$ there exists $h_\varphi \in \mathscr{H}$ such that $\varphi(m) = h_\varphi m$.*
(ii) $\mathrm{Hom}_{\mathscr{H}}(m\mathscr{H}, I) \cong \mathscr{H}m \cap I$.

PROOF: For all $x \in \mathrm{r}(m)$ we have $\varphi(m)x = \varphi(mx) = 0$. Therefore, $\varphi(m) \in \mathrm{lr}(m) = \mathscr{H}m$; so $\varphi(m) = h_\varphi m$ for some $h_\varphi \in \mathscr{H}$. The proof of part (ii) is now straightforward. □

The main result of this section is that $\mathrm{lr}(m_\mu) = \mathscr{H}m_\mu$. Because the Iwahori–Hecke algebra $\mathscr{H}(\mathfrak{S}_n)$ is quasi–Frobenius, our first step is easy.

(5.3) Lemma *Suppose that μ is a multicomposition of n. Then $\mathrm{lr}(x_\mu) = \mathscr{H}x_\mu$.*

PROOF: Since $\mathscr{H}(\mathfrak{S}_n)$ is a quasi–Frobenius algebra, we have $\mathrm{lr}(x_\mu) \cap \mathscr{H}(\mathfrak{S}_n) = \mathscr{H}(\mathfrak{S}_n)x_\mu$. Now suppose that $z \in \mathrm{lr}(x_\mu)$. By Theorem 2.2, we may write

$$z = \sum_{\mathbf{c} \in \mathscr{C}} L_1^{c_1}, \ldots L_n^{c_n} h_{\mathbf{c}}$$

where $\mathscr{C} = \{\, \mathbf{c} = (c_1, \ldots, c_r) \,|\, 0 \leq c_j < r \text{ for } 1 \leq j \leq n \,\}$ and $h_{\mathbf{c}} \in \mathscr{H}(\mathfrak{S}_n)$. Then, for all $y \in \mathrm{r}(x_\mu) \cap \mathscr{H}(\mathfrak{S}_n)$, we have

$$0 = zy = \sum_{\mathbf{c} \in \mathscr{C}} L_1^{c_1}, \ldots L_n^{c_n} h_{\mathbf{c}} y.$$

Therefore, by Theorem 2.2, we have $h_{\mathbf{c}} y = 0$ for all $\mathbf{c} \in \mathscr{C}$. Thus, $h_{\mathbf{c}} \in \mathrm{lr}(x_\mu) \cap \mathscr{H}(\mathfrak{S}_n) = \mathscr{H}(\mathfrak{S}_n)x_\mu$, and so $z \in \mathscr{H}x_\mu$. □

For the remainder of this section we fix a multicomposition μ and define $\mathbf{a} = (a_1, \ldots, a_r)$ by $a_k = \sum_{i=1}^{k-1} |\mu^{(i)}|$; so $0 = a_1 \leq a_2 \leq \ldots \leq a_r \leq n$. Our main task in this section is to prove that $\mathrm{lr}(u_{\mathbf{a}}^+) = \mathscr{H}u_{\mathbf{a}}^+$. To achieve this goal, we use the next result.

(5.4) Lemma *Assume that I is a left ideal of $\mathscr{H}$ and suppose that S is a subset of $\mathrm{r}(I)$ such that $\mathrm{l}(S) \subseteq I$. Then $\mathrm{lr}(I) = I$.*

PROOF: By assumption $S \subseteq \mathrm{r}(I)$ and $\mathrm{l}(S) \subseteq I$. Also, $I \subseteq \mathrm{lr}(I)$. Therefore, $\mathrm{lr}(I) \subseteq \mathrm{l}(S) \subseteq I \subseteq \mathrm{lr}(I)$; so $\mathrm{lr}(I) = I$ as required. □

Our definition of $u^+_{\mathbf{a}}$ expresses $u^+_{\mathbf{a}}$ as a double product

$$u^+_{\mathbf{a}} = \prod_{k=1}^{r} \prod_{m=1}^{a_k} (L_m - Q_k).$$

We now introduce notation which allows us to reverse the order of the multiplication here.

(5.5) Definition *Suppose that $1 \le i \le n$.*

(i) *Let $\gamma_i = k$ if k is maximal such that $a_k < i$. Let $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_n)$.*

(ii) *Let $v_i = \prod_{k=\gamma_i+1}^{r} (L_i - Q_k)$.*

Then $u^+_{\mathbf{a}} = v_1 v_2 \dots v_n$ and all of these factors commute.

(5.6) Definition *Suppose that $1 \le i \le n$. Let $y_i = T_{i-1} \dots T_2 T_1 \prod_{k=1}^{\gamma_i} (L_1 - Q_k)$.*

(5.7) Example Suppose that $r = 4$, $n = 5$ and $\mathbf{a} = (0, 2, 4, 5)$. Then

$$\begin{aligned} u^+_{\mathbf{a}} \;=\; & (L_1 - Q_2)(L_2 - Q_2) \\ & \times (L_1 - Q_3)(L_2 - Q_3)(L_3 - Q_3)(L_4 - Q_3) \\ & \times (L_1 - Q_4)(L_2 - Q_4)(L_3 - Q_4)(L_4 - Q_4)(L_5 - Q_4). \end{aligned}$$

Also $\gamma = (1, 1, 2, 2, 3)$ and, for $1 \le i \le n$, the product of the factors in the ith column of the above array is v_i. We have

$$\begin{aligned} y_1 &= (L_1 - Q_1) \\ y_2 &= T_1(L_1 - Q_1) \\ y_3 &= T_2T_1(L_1 - Q_1)(L_1 - Q_2) \\ y_4 &= T_3T_2T_1(L_1 - Q_1)(L_1 - Q_2) \\ y_5 &= T_4T_3T_2T_1(L_1 - Q_1)(L_1 - Q_2)(L_1 - Q_3). \end{aligned}$$

(5.8) Lemma *Suppose that $1 \le i \le n$. Then $v_1 v_2 \dots v_i y_i = 0$.*

PROOF: Assume that k satisfies $\gamma_i + 1 \leq k \leq r$. Then $v_1v_2\dots v_i$ has a factor $(L_1 - Q_k)(L_2 - Q_k)\dots(L_i - Q_k)$. By (2.1)(iv), this factor commutes with $T_{i-1}\dots T_2T_1$. Hence, $v_1v_2\dots v_iT_{i-1}\dots T_2T_1$ has a right factor $\prod_{k=\gamma_i+1}^{r}(L_1 - Q_k)$. However,

$$\prod_{k=1}^{r}(L_1 - Q_k) = \prod_{k=1}^{r}(T_0 - Q_k) = 0;$$

so it follows that $v_1v_2\dots v_iy_i = 0$ as claimed. □

Since $u^+_{\mathbf{a}} = v_1v_2\dots v_n$ we have the following Corollary.

(5.9) Corollary *Suppose that $1 \leq i \leq n$. Then $u^+_{\mathbf{a}}y_i = 0$.*

It turns out that $y_1, y_2, \dots, y_n$ generate $\mathrm{r}(u^+_{\mathbf{a}})$. More importantly, together with Lemma 5.4 the elements y_i will allow us to prove that $\mathrm{lr}(u^+_{\mathbf{a}}) = \mathscr{H}u^+_{\mathbf{a}}$. For our proof, we require the technical Lemma 5.13 below; for this result we need some preparation.

(5.10) Lemma *Suppose that $1 \leq i < n$ and that $a, b \in \{0, 1, \dots, r-1\}$.*

(i) *If $a \leq b$ then* $L_i^aL_{i+1}^bT_i = T_iL_i^bL_{i+1}^a + (q-1)\sum_{c=1}^{b-a} L_i^{b-c}L_{i+1}^{a+c}$.

(ii) *If $a > b$ then* $L_i^aL_{i+1}^bT_i = qT_i^{-1}L_i^bL_{i+1}^a - (q-1)\sum_{c=1}^{a-b-1} L_i^{b+c}L_{i+1}^{a-c}$.

PROOF: (i) Assume that $a \leq b$. If $b = a$ then T_i commutes with $L_i^aL_{i+1}^a$ by (2.1)(iii), so the result is correct in this case. Now suppose that $b > a$. Since $L_{i+1}T_i = T_iL_i + (q-1)L_{i+1}$ we have

$$L_i^aL_{i+1}^bT_i = L_i^aL_{i+1}^{b-1}(T_iL_i + (q-1)L_{i+1}),$$

which gives the required result by induction on $b - a$.

(ii) Either argue similarly, or apply the antiautomorphism $*$ to the result of part (i) and rearrange, interchanging a and b. □

We next generalize Lemma 5.10 as follows.

(5.11) Lemma *Suppose that $1 \leq i \leq n$ and $b_j \in \{0, 1, \dots, r-1\}$ for $j = 1, 2, \dots, n$. Then there exists an integer b and $\epsilon_1, \dots, \epsilon_{i-1} \in \{\pm 1\}$ such that*

$$L_1^{b_1}L_2^{b_2}\dots L_i^{b_i}T_{i-1}\dots T_2T_1 = x_1 + x_2,$$

where $x_1 = q^bT_{i-1}^{\epsilon_{i-1}}\dots T_2^{\epsilon_2}T_1^{\epsilon_1}L_1^{b_i}L_2^{b_1}L_3^{b_2}\dots L_i^{b_{i-1}}$ and x_2 is a linear combination of terms of the form $T_wL_1^{c_1}L_2^{c_2}\dots L_i^{c_i}$ where

(i) *$w \in \mathfrak{S}_n$; and*

(ii) $c_1, \dots, c_i \in \{0, 1, \dots, r-1\}$ *with* $c_1 + c_2 + \cdots + c_i = b_1 + b_2 + \cdots + b_i$*; and*

(iii) *either* $\prod_{j=1}^{i}(c_j+1) < \prod_{j=1}^{i}(b_j+1)$*, or* $c_1, c_2, \dots, c_i$ *is a permutation of* $b_1, b_2, \dots, b_i$ *and* $c_1 < b_i$.

PROOF: The result is true when $i = 1$. (In this case, $x_1 = L_1^{b_1}$ and $x_2 = 0$.) Assume, inductively, that $i < n$ and that the result as stated is true. By Lemma 5.10, there exists $b' \in \{0, 1\}$ and $\epsilon_i = \pm 1$ such that

$$L_i^{b_i} L_{i+1}^{b_{i+1}} T_i = q^{b'} T_i^{\epsilon_i} L_i^{b_{i+1}} L_{i+1}^{b_i} \pm (q-1) \sum L_i^{d_i} L_{i+1}^{d_{i+1}}$$

where $d_i, d_{i+1} \in \{0, 1, \dots, r-1\}$ and $d_i + d_{i+1} = b_i + b_{i+1}$ and either $(d_i+1)(d_{i+1}+1) < (b_i+1)(b_{i+1}+1)$ or $d_i = b_i < b_{i+1}$. Using this result, and (2.1)(ii), we obtain

$$\begin{aligned} L_1^{b_1} L_2^{b_2} \dots L_i^{b_i} T_{i-1} \dots T_2 T_1 \;=\; & q^{b'} T_i^{\epsilon_i} L_1^{b_1} \dots L_{i-1}^{b_{i-1}} L_i^{b_{i+1}} T_{i-1} \dots T_2 T_1 L_{i+1}^{b_i} \\ & \pm (q-1) \sum L_1^{b_1} \dots L_{i-1}^{b_{i-1}} L_i^{d_i} T_{i-1} \dots T_2 T_1 L_{i+1}^{d_{i+1}}, \end{aligned}$$

where b', d_i and d_{i+1} are as above. The result now follows from our inductive hypothesis. □

For the statement and proof of our next Lemma, we need some more notation.

(5.12) Notation Suppose that $0 \le i \le n$. Let V_i be the R–module with basis

$$\left\{ T_w L_1^{b_1} L_2^{b_2} \dots L_n^{b_n} v_1 v_2 \dots v_i \;\middle|\; \begin{array}{c} w \in \mathfrak{S}_n,\ 0 \le b_j < \gamma_j, \text{ for } 1 \le j \le i, \\ \text{and } 0 \le b_j < r, \text{ for } i+1 \le j \le n \end{array} \right\}.$$

Result (2.4) shows the elements in these sets are indeed linearly independent. Note also that $V_0 = \mathscr{H}$ and $V_n = \mathscr{H} u_{\mathbf{a}}^+$.

We shall use (2.4) extensively in the proof of Lemma 5.13.

(5.13) Lemma *Suppose that* $1 \le i \le n$ *and that* $z \in V_{i-1}$ *and* $z y_i = 0$*. Then* $z \in V_i$.

PROOF: Since $z \in V_{i-1}$ we may write z as a linear combination of linearly independent terms

$$T_w L_1^{b_1} \dots L_{i-1}^{b_{i-1}} f(L_i) L_{i+1}^{b_{i+1}} \dots L_n^{b_n} v_1 v_2 \dots v_{i-1}$$

where $w \in \mathfrak{S}_n$ and $0 \le b_j < \gamma_j$ for $1 \le j \le i-1$ and $0 \le b_j < r$ for $i+1 \le j \le n$, and $f(X)$ is a polynomial in $R[X]$ of degree less than r. Write

$$f(X) = g(X) \prod_{k=\gamma_i+1}^{r} (X - Q_k) + h(X),$$

where $g(X)$ and $h(X)$ are polynomials in $R[X]$ such that $\deg g(X) < \gamma_i$ and $\deg h(X) < r - \gamma_i$. We may apply the "Euclidean algorithm" in this way, even though $R[X]$ need not be a Euclidean domain, because the divisor is a monic polynomial.

Note that $\prod_{k=\gamma_i+1}^{r}(L_i - Q_k) = v_i$. As a consequence, we have written z as $z_1 + z_2$ where $z_1 \in V_i$ and z_2 is a linear combination of terms

$$T_w L_1^{b_1} \dots L_{i-1}^{b_{i-1}} h(L_i) L_{i+1}^{b_{i+1}} \dots L_n^{b_n} v_1 v_2 \dots v_{i-1}.$$

We shall prove that $z_2 = 0$.

Note that $z_1 y_i = 0$, by Lemma 5.8, since $z_1 \in V_i$. Therefore, because $z y_i = 0$ by assumption, it follows that $z_2 y_i = 0$.

For each n–tuple $\mathbf{c} = (c_1, \dots, c_n)$ with $0 \le c_i < r$, for all i with $1 \le i \le n$, let $U_{\mathbf{c}} = \mathscr{H}(\mathfrak{S}_n) L_1^{c_1}, \dots L_n^{c_n}$. Then we may write

$$z_2 = \sum_{\mathbf{c} \in \mathscr{C}} z_{\mathbf{c}} \quad \text{where} \quad z_{\mathbf{c}} \in U_{\mathbf{c}}$$

and $\mathscr{C} = \{\, \mathbf{c} = (c_1, \dots, c_n) \,|\, 0 \le c_j < r \text{ for } 1 \le j \le n \text{ and } c_i < r - \gamma_i \,\}$.

Assume that $z_2 \neq 0$. Among the $\mathbf{c}$ such that $z_{\mathbf{c}} \neq 0$ choose $\mathbf{d} = (d_1, \dots, d_n)$ such that

(5.14) (i) $d_1 + d_2 + \dots + d_n$ is maximal; and,
(ii) $(d_1 + 1)(d_2 + 1) \dots (d_n + 1)$ is maximal subject to (i); and,
(iii) d_i is maximal subject to (i) and (ii).

Then $z_{\mathbf{d}} = h_{\mathbf{d}} L_1^{d_1}, \dots L_n^{d_n}$ for some non–zero element $h_{\mathbf{d}}$ of $\mathscr{H}(\mathfrak{S}_n)$.

Now, $T_{i-1} \dots T_2 T_1$ commutes with each of the elements $L_{i+1}, L_{i+2}, \dots, L_n$. Therefore, by Lemma 5.11 there exists an invertible element $h \in \mathscr{H}(\mathfrak{S}_n)$ (namely, $h = q^b T_{i-1}^{\epsilon_{i-1}} \dots T_2^{\epsilon_2} T_1^{\epsilon_1}$) such that

$$z_2 T_{i-1} \dots T_2 T_1 = h_{\mathbf{d}} h L_1^{d_i} L_2^{d_1} L_3^{d_2} \dots L_i^{d_{i-1}} L_{i+1}^{d_{i+1}} \dots L_n^{d_n} + \text{ terms in the sets } U_{\mathbf{e}},$$

where $\mathbf{e} = (e_1, \dots, e_n)$ and $e_1 + \dots + e_n \le d_1 + \dots + d_n$, and either
(i) $e_1 + \dots + e_n < d_1 + \dots + d_n$ (by Lemma 5.11(ii) and (5.14)(i)); or,
(ii) $(e_1 + 1) \dots (e_n + 1) < (d_1 + 1) \dots (d_n + 1)$ (using Lemma 5.11(iii) and (5.14)(ii)); or,
(iii) $e_1, \dots, e_n$ is a permutation of $d_1, \dots, d_n$ and $e_1 < d_i$ (using Lemma 5.11(iii) and (5.14)(iii)); or,
(iv) $e_1, \dots, e_n$ is a permutation of $d_1, \dots, d_n$ but $\mathbf{e} \neq (d_i, d_1, d_2, \dots, d_{i-1}, d_{i+1}, \dots, d_n)$ (consider the term x_1 in Lemma 5.11).

In particular, no $\mathbf{e}$ on the right hand side is equal to $(d_i, d_1, d_2, \dots, d_{i-1}, d_{i+1}, \dots, d_n)$.

Next we postmultiply our expression for $z_2 T_{i-1} \dots T_2 T_1$ by $\prod_{m=1}^{\gamma_i}(L_1 - Q_m)$ to obtain $z_2 y_i$. Note that if $v_{\mathbf{e}} \in U_{\mathbf{e}}$ and k is any positive integer then $v_{\mathbf{e}} L_1^k$ is a linear combination of terms $v_{\mathbf{f}} \in U_{\mathbf{f}}$ where $\mathbf{f} = (f_1, e_2, \dots, e_n)$ and $f_1 < r$. Thus, if $e_1 + \dots + e_n \le d_1 + \dots + d_n$ and $e_1 + k \ge r$ then $v_{\mathbf{e}} L_1^k$ is a linear combination of terms $v_{\mathbf{f}}$ with $f_1 + e_2 + \dots + e_n < d_1 + \dots + d_n + k$. Therefore,

$$z_2 y_i = h_{\mathbf{d}} h L_1^{d_i + \gamma_i} L_2^{d_1} L_3^{d_2} \dots L_i^{d_{i-1}} L_{i+1}^{d_{i+1}} \dots L_n^{d_n} + \text{ terms in the sets } U_{\mathbf{f}}$$

where $\mathbf{f} \neq (d_i + \gamma_i, d_1, d_2, \dots, d_{i-1}, d_{i+1}, \dots, d_n)$.

Note that $d_i + \gamma_i < r$ since $\mathbf{d} \in \mathscr{C}$. Therefore, since $h_{\mathbf{d}} \neq 0$ and h is invertible, $z_2 y_i \neq 0$ giving a contradiction. Hence, our assumption that $z_2 \neq 0$ is false. Thus, $z = z_1 \in V_i$ as required. □

(5.15) Corollary *We have* $\mathrm{lr}(u_{\mathbf{a}}^+) = \mathscr{H} u_{\mathbf{a}}^+$.

PROOF: Let $I = \mathscr{H} u_{\mathbf{a}}^+$ and $S = \{y_1, y_1, \dots, y_n\}$. Then Corollary 5.9 shows that $S \subseteq \mathrm{r}(I)$.

Suppose that $z \in \mathscr{H}$ and $zy_i = 0$ for all i with $1 \leq i \leq n$. Since $V_0 = \mathscr{H}$ and $V_n = \mathscr{H} u_{\mathbf{a}}^+ = I$, it follows from Lemma 5.13 that $z \in I$. Thus, $\mathrm{l}(S) \subseteq I$. Lemma 5.4 now implies that $\mathrm{lr}(I) = I$; that is, $\mathrm{lr}(u_{\mathbf{a}}^+) = \mathscr{H} u_{\mathbf{a}}^+$. □

Finally, we can prove the main result of this section.

(5.16) Theorem *Suppose that μ is a multicomposition of n. Then* $\mathrm{lr}(m_\mu) = \mathscr{H} m_\mu$.

PROOF: Write $m_\mu = u_{\mathbf{a}}^+ x_\mu$ as in Definition 3.5. By applying the definitions, Corollary 5.15 and Lemma 5.3, we obtain

$$\mathscr{H} m_\mu \subseteq \mathrm{lr}(m_\mu) \subseteq \mathrm{lr}(u_{\mathbf{a}}^+) \cap \mathrm{lr}(x_\mu) = \mathscr{H} u_{\mathbf{a}}^+ \cap \mathscr{H} x_\mu.$$

However, $(u_{\mathbf{a}}^+ \mathscr{H} \cap x_\mu \mathscr{H})^* = (m_\mu \mathscr{H})^*$ by Corollary 4.13. Hence, we have equality throughout and $\mathrm{lr}(m_\mu) = \mathscr{H} m_\mu$ as claimed. □

(5.17) Corollary *Suppose that μ and ν are multicompositions of n. Then*

(i) *For every element φ in $\mathrm{Hom}_{\mathscr{H}}(M^\nu, M^\mu)$ there exists $h_\varphi \in \mathscr{H}$ such that $\varphi(m_\nu) = h_\varphi m_\nu$; in particular, $\varphi(m_\nu) \in M^{\nu *} \cap M^\mu$.*

(ii) $\mathrm{Hom}_{\mathscr{H}}(M^\nu, M^\mu) \cong M^{\nu *} \cap M^\mu$.

PROOF: By Theorem 5.16 we may apply Lemma 5.2(i) with $m = m_\nu$ to obtain that $\varphi(m_\nu) = h_\varphi m_\nu$ for some $h_\varphi \in \mathscr{H}$. Thus, $\varphi(m_\nu) \in M^{\nu *}$. It is clear that $\varphi(m_\nu) \in M^\mu$. Part (ii) follows from Lemma 5.2(ii), where an explicit isomorphism is given by $\varphi \longmapsto \varphi(m_\nu)$. □

6 The cyclotomic q–Schur algebra

In order to make our results as general as possible, we fix an integer d (or take $d = \infty$) and let $\Lambda_r(d,n)$ be the set of all multicompositions $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ of n such that each composition $\lambda^{(k)}$ has at most d parts for all k with $1 \le k \le r$. Let $\Lambda_r^+(d,n)$ be the set of multipartitions of n contained in $\Lambda_r(d,n)$. Then $\Lambda_r^+(d,n)$ contains all multipartitions of n if and only if $d \ge n$.

(6.1) Definition *The* cyclotomic q–Schur algebra *is the endomorphism algebra*

$$\mathscr{S} = \mathrm{End}_{\mathscr{H}}\Big(\bigoplus_{\mu\in\Lambda_r(d,n)} M^\mu\Big).$$

Thus, $\mathscr{S} \cong \bigoplus_{\mu,\nu\in\Lambda_r(d,n)} \mathrm{Hom}_{\mathscr{H}}(M^\nu, M^\mu)$.

Suppose that $\varphi \in \mathrm{Hom}_{\mathscr{H}}(M^\nu, M^\mu)$. Then $\varphi(m_\nu h) = \varphi(m_\nu)h$ for all $h \in \mathscr{H}$; thus φ is completely determined by $\varphi(m_\nu)$. Moreover, $\varphi(m_\nu) \in M^{\nu *} \cap M^\mu$ by Corollary 5.17. These remarks motivate us to construct a basis of $M^{\nu *} \cap M^\mu$.

(6.2) Definition *Suppose that μ and ν are multicompositions of n and that λ is a multipartition of n. Assume that $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$ and that $\mathtt{T} \in \mathcal{T}_0(\lambda,\nu)$. Let*

$$m_{\mathtt{ST}} = \sum_{\mathfrak{s},\mathfrak{t}} m_{\mathfrak{s}\mathfrak{t}}$$

where the sum is over all $\mathfrak{s},\mathfrak{t} \in \mathrm{Std}(\lambda)$ with $\mu(\mathfrak{s}) = \mathtt{S}$ and $\nu(\mathfrak{t}) = \mathtt{T}$.

Note that $m_{\mathtt{ST}}^* = m_{\mathtt{TS}}$.

(6.3) Proposition *Suppose that μ and ν are multicompositions of n. Then*

$$\Big\{ m_{\mathtt{ST}} \;\Big|\; \begin{array}{c}\mathtt{S} \in \mathcal{T}_0(\lambda,\mu) \text{ and } \mathtt{T} \in \mathcal{T}_0(\lambda,\nu) \text{ for some}\\ \text{multipartition } \lambda \text{ of } n\end{array} \Big\}$$

is a basis of $M^{\nu *} \cap M^\mu$.

PROOF: Since

$$m_{\mathtt{ST}} = \sum_{\substack{\mathfrak{s}\in\mathrm{Std}(\lambda)\\ \mu(\mathfrak{s})=\mathtt{S}}} (m_{\mathtt{Ts}})^* = \sum_{\substack{\mathfrak{t}\in\mathrm{Std}(\lambda)\\ \nu(\mathfrak{t})=\mathtt{T}}} m_{\mathtt{S}\mathfrak{t}},$$

Lemma 4.10 shows that $m_{\mathtt{ST}} \in M^{\nu *} \cap M^\mu$. Moreover, the elements $m_{\mathtt{ST}}$ are linearly independent since they involve distinct elements $m_{\mathfrak{s}\mathfrak{t}}$ of the standard basis of $\mathscr{H}$.

Now suppose that $h \in M^{\nu *} \cap M^\mu$. Since $h \in \mathscr{H}$, we may express h in terms of the standard basis $\mathscr{M}$; say

$$h = \sum_{m_{\mathfrak{s}\mathfrak{t}}\in\mathscr{M}} r_{\mathfrak{s}\mathfrak{t}} m_{\mathfrak{s}\mathfrak{t}}$$

where $r_{\mathfrak{st}} \in R$. Since $h \in M^\mu$, we have $r_{\mathfrak{st}} = r_{\mathfrak{s}'\mathfrak{t}}$ if $\mu(\mathfrak{s}) = \mu(\mathfrak{s}')$, by Theorem 4.14. Similarly, since $h \in M^{\nu *}$, we have $r_{\mathfrak{st}} = r_{\mathfrak{st}'}$ if $\nu(\mathfrak{t}) = \nu(\mathfrak{t}')$. Thus, if $\mu(\mathfrak{s}) = \mu(\mathfrak{s}')$ and $\nu(\mathfrak{t}) = \nu(\mathfrak{t}')$ then $r_{\mathfrak{st}} = r_{\mathfrak{s}'\mathfrak{t}} = r_{\mathfrak{s}'\mathfrak{t}'}$. Furthermore, $r_{\mathfrak{st}} = 0$ unless both $\mu(\mathfrak{s})$ and $\nu(\mathfrak{t})$ are semistandard. Therefore, h is a linear combination of elements $m_{\mathtt{ST}}$. This completes the proof of the Proposition. □

Proposition 6.3 shows that, in our next Definition, $\varphi_{\mathtt{ST}}$ is a well defined $\mathscr{H}$–homomorphism from M^ν into M^μ.

(6.4) Definition *Suppose that $\mu, \nu \in \Lambda_r(d,n)$ and $\lambda \in \Lambda_r^+(d,n)$. Assume that $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$ and $\mathtt{T} \in \mathcal{T}_0(\lambda,\nu)$. Define $\varphi_{\mathtt{ST}} \in \mathrm{Hom}_{\mathscr{H}}(M^\nu, M^\mu)$ by*

$$\varphi_{\mathtt{ST}}(m_\nu h) = m_{\mathtt{ST}} h$$

for all $h \in \mathscr{H}$. Extend $\varphi_{\mathtt{ST}}$ to an element of the cyclotomic q–Schur algebra $\mathscr{S}$ by defining $\varphi_{\mathtt{ST}}$ to be zero on M^κ when $\nu \neq \kappa \in \Lambda_r(d,n)$.

(6.5) Definition *Suppose that $\lambda \in \Lambda_r^+(d,n)$. Let $\overline{\mathscr{S}_\lambda}$ be the R–submodule of $\mathscr{S}$ spanned by*

$$\left\{ \varphi_{\mathtt{UV}} \;\middle|\; \begin{array}{c} \mathtt{U} \in \mathcal{T}_0(\alpha,\mu),\ \mathtt{V} \in \mathcal{T}_0(\alpha,\nu) \text{ for some} \\ \mu,\nu \in \Lambda_r(d,n) \text{ and } \alpha \in \Lambda_r^+(d,n) \text{ with } \alpha \rhd \lambda \end{array} \right\}.$$

This brings us to one of the main results of our paper. The work in Sections 4 and 5 was aimed at proving part (i) of the Theorem.

(6.6) Theorem (The Semistandard Basis Theorem)

(i) *The cyclotomic q–Schur algebra $\mathscr{S}$ is free as an R–module with basis*

$$\left\{ \varphi_{\mathtt{ST}} \;\middle|\; \begin{array}{c} \mathtt{S} \in \mathcal{T}_0(\lambda,\mu),\ \mathtt{T} \in \mathcal{T}_0(\lambda,\nu) \text{ for some} \\ \mu,\nu \in \Lambda_r(d,n) \text{ and } \lambda \in \Lambda_r^+(d,n) \end{array} \right\}.$$

(ii) *Suppose that $\mu, \nu \in \Lambda_r(d,n)$ and $\lambda \in \Lambda_r^+(d,n)$ and let $\varphi \in \mathscr{S}$. Then, for every $\kappa \in \Lambda_r(d,n)$ and every $\mathtt{T}' \in \mathcal{T}_0(\lambda,\kappa)$ there exists $r_{\mathtt{T}'} \in R$ such that for all $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$, we have*

$$\varphi_{\mathtt{ST}}\varphi \equiv \sum_{\kappa \in \Lambda_r(d,n)} \sum_{\mathtt{T}' \in \mathcal{T}_0(\lambda,\kappa)} r_{\mathtt{T}'} \varphi_{\mathtt{ST}'} \bmod \overline{\mathscr{S}_\lambda}.$$

PROOF: (i) By Corollary 5.17(ii), $\mathrm{Hom}_{\mathscr{H}}(M^\nu, M^\mu) \cong M^{\nu *} \cap M^\mu$, where an isomorphism is given by $\varphi \longmapsto \varphi(m_\nu)$. Hence, by Proposition 6.3 and Definition 6.4, the set $\{\varphi_{\mathtt{ST}}\}$ is a basis of $\mathscr{S}$.

(ii) It is sufficient to consider the case where $\varphi \in \mathrm{Hom}_{\mathscr{H}}(M^\kappa, M^\nu)$ for some $\kappa \in \Lambda_r(d,n)$. Suppose that $\varphi(m_\kappa) = m_\nu h$ where $h \in \mathscr{H}$. Then, for all $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$ and $\mathtt{T} \in \mathcal{T}_0(\lambda,\nu)$, we have

$$\varphi_{\mathtt{ST}}\varphi(m_\kappa) = m_{\mathtt{ST}} h \in M^{\kappa *} \cap M^\mu.$$

By Proposition 6.3, $m_{\mathtt{ST}}h = \sum r_{\mathtt{UV}} m_{\mathtt{UV}}$, where $r_{\mathtt{UV}} \in R$ and the sum is over $\mathtt{U} \in \mathcal{T}_0(\alpha,\mu)$ and $\mathtt{V} \in \mathcal{T}_0(\alpha,\kappa)$ for some $\alpha \in \Lambda_r^+(d,n)$. By applying Proposition 3.25 we deduce that

$$m_{\mathtt{ST}}h = \sum_{\mathtt{T}' \in \mathcal{T}_0(\lambda,\kappa)} r_{\mathtt{T}'} m_{\mathtt{ST}'} + \sum_{\mathtt{U}',\mathtt{V}'} r_{\mathtt{U}'\mathtt{V}'} m_{\mathtt{U}'\mathtt{V}'}$$

where $r_{\mathtt{T}'}, r_{\mathtt{U}'\mathtt{V}'} \in R$ and the second sum is over $\mathtt{U}' \in \mathcal{T}_0(\alpha,\mu)$ and $\mathtt{V}' \in \mathcal{T}_0(\alpha,\nu)$ for some $\alpha \in \Lambda_r^+(d,n)$ with $\alpha \rhd \lambda$. Therefore,

$$\varphi_{\mathtt{ST}}\varphi \equiv \sum_{\mathtt{T}' \in \mathcal{T}_0(\lambda,\kappa)} r_{\mathtt{T}'} \varphi_{\mathtt{ST}'} \bmod \overline{\mathscr{S}_\lambda}.$$

This completes the proof. □

(6.7) Definition *We call the basis* $\{\varphi_{\mathtt{ST}}\}$ *the* semistandard basis *of* $\mathscr{S}$.

(6.8) Remark If $d \geq n$ then $\Lambda_r^+(d,n)$ consists of all multipartitions of n. It is straightforward to prove that if $d > n$, then the algebra $\mathscr{S}$ is Morita equivalent to the algebra we obtain by taking $d = n$.

In order to show that the semistandard basis of $\mathscr{S}$ is a cellular basis we need an appropriate antiautomorphism $*$ for $\mathscr{S}$.

(6.9) Proposition *Let* $* : \mathscr{S} \longrightarrow \mathscr{S}$ *be the unique* R*–linear map such that* $\varphi_{\mathtt{ST}}^* = \varphi_{\mathtt{TS}}$ *for all elements* $\varphi_{\mathtt{ST}}$ *in the semistandard basis. Then* $*$ *is an antiautomorphism of* $\mathscr{S}$.

PROOF: Assume that $\mathtt{S} \in \mathcal{T}_0(\lambda,\mu)$ and that $\mathtt{T} \in \mathcal{T}_0(\lambda,\nu)$ for some $\mu,\nu \in \Lambda_r(d,n)$ and $\lambda \in \Lambda_r^+(d,n)$. Then

$$\varphi_{\mathtt{ST}}(m_\nu) = m_{\mathtt{ST}} = (m_{\mathtt{TS}})^* = (\varphi_{\mathtt{TS}}(m_\mu))^*.$$

The proposition now follows from the next general lemma.

(6.10) Lemma *Suppose that* $\mathscr{H}$ *is an algebra with an antiautomorphism* $*$. *Let* $\{\, m_\mu \mid \mu \in \Lambda \,\}$ *be a set of elements of* $\mathscr{H}$ *such that* $m_\mu^* = m_\mu$ *for all* $\mu \in \Lambda$ *and let*

$$S = \mathrm{End}_{\mathscr{H}}\Big(\bigoplus_{\mu\in\Lambda} m_\mu \mathscr{H}\Big).$$

Assume that, for all $\mu,\nu \in \Lambda$, *every* $\mathscr{H}$*–homomorphism from* $m_\nu\mathscr{H}$ *to* $m_\mu\mathscr{H}$ *is given by left multiplication by an element of* $\mathscr{H}$. *Then the following hold.*

(i) *For each* $\varphi \in \mathrm{Hom}_{\mathscr{H}}(m_\nu\mathscr{H}, m_\mu\mathscr{H})$ *there is a unique* $\varphi^* \in \mathrm{Hom}_{\mathscr{H}}(m_\mu\mathscr{H}, m_\nu\mathscr{H})$ *such that* $\varphi^*(m_\mu) = (\varphi(m_\nu))^*$.

(ii) *The map* $*$ *is an antiautomorphism of* $\mathscr{S}$.

PROOF: (i) Suppose that $\varphi \in \mathrm{Hom}_{\mathscr{H}}(m_\nu\mathscr{H}, m_\mu\mathscr{H})$. Then $\varphi(m_\nu) = x_1 m_\nu = m_\mu y$ for some $x_1, y \in \mathscr{H}$. Define $\varphi^* \in \mathrm{Hom}_{\mathscr{H}}(m_\mu\mathscr{H}, m_\nu\mathscr{H})$ by $\varphi^*(m_\mu h) = y^* m_\mu h$ for all $h \in \mathscr{H}$. Then φ^* is a well–defined $\mathscr{H}$–homomorphism; also, it maps into $m_\nu\mathscr{H}$, since $y^* m_\mu = m_\nu x_1^*$. Since $\varphi^*(m_\mu) = (\varphi(m_\nu))^*$, the proof of (i) is complete.

(ii) Suppose that $\psi \in \mathrm{Hom}_{\mathscr{H}}(m_\mu\mathscr{H}, m_\lambda\mathscr{H})$ for some $\lambda \in \Lambda$. Then $\psi(m_\mu) = x_2 m_\mu$ for some $x_2 \in \mathscr{H}$. We have,

$$\begin{aligned}(\psi\varphi)^*(m_\lambda) = (\psi\varphi(m_\nu))^* &= (x_2 x_1 m_\nu)^* &&= m_\nu x_1^* x_2^* \\ &= \varphi^*(m_\mu)x_2^* &&= \varphi^*(m_\mu x_2^*) \quad = \varphi^*\psi^*(m_\lambda).\end{aligned}$$

Therefore, $(\psi\varphi)^* = \varphi^*\psi^*$, and it follows that $*$ is an antiautomorphism of $\mathscr{S}$. □

(6.11) Corollary *The R–module $\overline{\mathscr{S}_\lambda}$ is a two–sided ideal of $\mathscr{S}$.*

PROOF: Theorem 6.6(ii) shows that $\overline{\mathscr{S}_\lambda}$ is a right ideal. By applying the antiautomorphism $*$, we deduce that it is also a left ideal. □

(6.12) Theorem *The semistandard basis of $\mathscr{S}$ is a cellular basis.*

PROOF: This follows at once from Theorem 6.6, Corollary 6.11 and Proposition 6.9. □

We now apply the theory of cellular algebras to the representation theory of $\mathscr{S}$, just as we treated $\mathscr{H}$ in Section 3.

Suppose that λ is a multipartition of n. Let $\mathtt{T}^\lambda = \lambda(\mathfrak{t}^\lambda)$ (see Definition 4.2). Then $\mathtt{T}^\lambda$ is the unique semistandard λ–tableau of type λ (cf. Example 4.3). Define $\varphi_\lambda = \varphi_{\mathtt{T}^\lambda\mathtt{T}^\lambda}$; then φ_λ is the identity map on M^λ.

(6.13) Definition *Suppose that $\lambda \in \Lambda_r^+(d,n)$, The* Weyl module *$W^\lambda$ is the submodule of $\mathscr{S}/\overline{\mathscr{S}_\lambda}$ given by $W^\lambda = \mathscr{S}(\varphi_\lambda + \overline{\mathscr{S}_\lambda})$.*

If $\mathtt{S}$ is a semistandard λ–tableau let $\varphi_\mathtt{S} = \varphi_{\mathtt{S}\mathtt{T}^\lambda}(\varphi_\lambda + \overline{\mathscr{S}_\lambda}) = \varphi_{\mathtt{S}\mathtt{T}^\lambda} + \overline{\mathscr{S}_\lambda}$. Then, from Theorem 6.12, we obtain the following result.

(6.14) Corollary *The Weyl module W^λ is a free R–module with basis*

$$\{\, \varphi_\mathtt{S} \mid \mathtt{S} \in \mathcal{T}_0(\lambda,\mu) \text{ for some } \mu \in \Lambda_r(d,n) \,\}.$$

Define a bilinear form $\langle\ ,\ \rangle$ on W^λ by requiring that

$$\varphi_{\mathtt{T}^\lambda\mathtt{S}}\varphi_{\mathtt{T}\mathtt{T}^\lambda} \equiv \langle \varphi_\mathtt{S}, \varphi_\mathtt{T}\rangle \varphi_\lambda \bmod \overline{\mathscr{S}_\lambda}$$

for all semistandard λ–tableaux $\mathtt{S}$ and $\mathtt{T}$. This bilinear form is well–defined and symmetric, and also satisfies $\langle su, v\rangle = \langle u, s^* v\rangle$ for all $u, v \in W^\lambda$ and all $s \in \mathscr{S}$ [**12**, (2.4)]. Consequently, $\operatorname{rad} W^\lambda = \{\, u \in W^\lambda \mid \langle u, v\rangle = 0 \text{ for all } v \in W^\lambda \,\}$ is a submodule of W^λ.

(6.15) Definition *Suppose that $\lambda \in \Lambda_r^+(d,n)$. Let $F^\lambda = W^\lambda / \operatorname{rad} W^\lambda$.*

(6.16) Theorem *Suppose that R is a field. Then*

$$\{ F^\lambda \mid \lambda \in \Lambda_r^+(d,n) \}$$

is a complete set of non–isomorphic irreducible $\mathscr{S}$–modules. Moreover, each F^λ is absolutely irreducible.

PROOF: Let $\lambda \in \Lambda_r^+(d,n)$. From the definition of the bilinear form on W^λ, we have

$$\varphi_{\mathtt{T}^\lambda \mathtt{T}^\lambda} \varphi_{\mathtt{T}^\lambda \mathtt{T}^\lambda} \equiv \langle \varphi_{\mathtt{T}^\lambda}, \varphi_{\mathtt{T}^\lambda} \rangle \varphi_\lambda \bmod \overline{\mathscr{S}_\lambda}.$$

However, $\varphi_{\mathtt{T}^\lambda \mathtt{T}^\lambda} \varphi_{\mathtt{T}^\lambda \mathtt{T}^\lambda} = \varphi_\lambda$ is the identity on M^λ; so $\langle \varphi_{\mathtt{T}^\lambda}, \varphi_{\mathtt{T}^\lambda} \rangle = 1$. Consequently, F^λ is non–zero. The Theorem now follows from [**12**, (3.4)]. □

If $\lambda, \mu \in \Lambda_r^+(d,n)$, let $d_{\lambda\mu}$ denote the composition multiplicity of F^μ as a composition factor of W^λ. Then

$$(d_{\lambda\mu})_{\lambda,\mu \in \Lambda_r^+(d,n)}$$

is the decomposition matrix of $\mathscr{S}$. The theory of cellular algebras [**12**, (3.6)] yields the following.

(6.17) Corollary *The decomposition matrix of $\mathscr{S}$ is unitriangular. That is, for $\lambda, \mu \in \Lambda_r^+(d,n)$, we have $d_{\mu\mu} = 1$ and $d_{\lambda\mu} \neq 0$ only if $\lambda \trianglerighteq \mu$.*

Finally, Theorem 6.16 combined with [**12**, (3.10)], gives us our last result.

(6.18) Corollary *The cyclotomic q–Schur algebra is quasi–hereditary.*

References

[**1**] S. ARIKI AND K. KOIKE, *A Hecke algebra of $(\mathbb{Z}/r\mathbb{Z}) \wr \mathfrak{S}_n$ and construction of its irreducible representations*, Adv. Math, **106** (1994), 216–243.

[**2**] M. BROUÉ AND G. MALLE, *Zyklotomische Heckealgebren*, Asterisque, **212** (1993), 119–189.

[**3**] C. W. CURTIS AND I. REINER, *Representation theory of finite groups and associative algebras*, Interscience, 1962.

[**4**] R. DIPPER AND G. JAMES, *Representations of Hecke algebras of general linear groups*, Proc. London Math. Soc. (3), **52** (1986), 20–52.

[**5**] ——, *Blocks and idempotents of Hecke algebras of general linear groups*, Proc. London Math. Soc. (3), **54** (1987), 57–82.

[**6**] ——, *The q–Schur algebra*, Proc. London Math. Soc. (3), **59** (1989), 23–50.

[**7**] ——, *q–Tensor space and q–Weyl modules*, Trans. AMS, **327** (1991), 251–282.

[8] ——, *Representations of Hecke algebras of type B_n*, J. Algebra, **146** (1992), 454–481.
[9] R. DIPPER, G. JAMES, AND A. MATHAS, *The (Q,q)–Schur algebra*, Proc. London Math. Soc. (3), to appear.
[10] R. DIPPER, G. JAMES, AND E. MURPHY, *Hecke algebras of type B_n at roots of unity*, Proc. London Math. Soc. (3), **70** (1995), 505–528.
[11] J. DU AND L. SCOTT, *The q–Schur^2 algebra*, preprint, (1997).
[12] J. J. GRAHAM AND G. I. LEHRER, *Cellular algebras*, Invent., **123** (1996), 1–34.
[13] A. MATHAS, *Canonical bases and the decomposition matrices of Ariki–Koike algebras*, J. Algebra, (to appear).
[14] ——, *Simple modules of Ariki–Koike algebras*, Proc. Sym. Pure Math., to appear.
[15] G. E. MURPHY, *On the representation theory of the symmetric groups and associated Hecke algebras*, J. Algebra, **152** (1982), 492–513.
[16] ——, *The representations of Hecke algebras of type $\mathbf{A_n}$*, J. Algebra, **173** (1995), 97–121.

E–mail addresses: rdipper@mathematik.uni-stuttgart.de
g.james@ic.ac.uk
sfb32@Mathematik.Uni-Bielefeld.de