FIBRE SUM FUNCTORS AND THE BIMODULE EXT

PETER DRÄXLER

1. Introduction

It is a classical technique in the representation theory of finite-dimensional algebras A to consider the A-modules as extensions of modules over smaller subcategories $\mathcal{F}$ and $\mathcal{T}$ thus identifying the module category with the representation category of the bimodule $\mathrm{Ext}^1_A(-,-)$ acting on $\mathcal{F} \times \mathcal{T}$. If $\mathcal{T} = \operatorname{add} S$ for a simple module S the category of representations of this bimodule can be identified with the subspace category of a vector space category (see [Ri1]).

In [Dr1] the fibre sum functor with respect to a module P is introduced. It relates the module with the representations of a bimodule as well. In case the endomorphism algebra of P is a field or more specially if P is a simple module, this leads to a vector space category as well.

The aim of this note is to analyse the relation between these two reduction processes to vector space categories. After recalling the first reduction concept in the next section, in the final section we will clarify the relation completely.

For simplicity we consider only finite-dimensional algebras over an algebraically closed field k which we assume to be basic. We use the term algebra for this concept. For notation and background we refer to [GR] and [Ri2].

2. Reduction to the bimodule $\mathrm{Ext}_A(-,-)$

2.1. Let us start out by recalling the concept of a bimodule. Following [GR] an aggregate is a k-additive category with finite-dimensional morphism spaces such that each object is the direct sum of subobjects with local endomorphism algebras. A typical example for an aggregate is the category A−mod of finite-dimensional (left) modules over an algebra A.

A bimodule over two aggregates $\mathcal{F}$ and $\mathcal{T}$ is a k-linear bifunctor $H\colon \mathcal{F} \times \mathcal{T} \to k$−mod which is covariant in the second and contravariant in the first argument. The category $\mathrm{rep}(H)$ of representations of H has as objects the triples (X, h, Y) where $X \in \mathcal{F}$, $Y \in \mathcal{T}$ and $h \in H(X, Y)$. A morphism from (X, h, Y) to (X', h', Y') is a pair (s, t) of morphisms $s\colon X \to X'$ in $\mathcal{F}$ and $t\colon Y \to Y'$ in $\mathcal{T}$ such that $H(X, t)(h) = H(s, Y')(h')$. The category $\mathrm{rep}(H)$ is again an aggregate.

2.2. The classical example to study the category A−mod of an algebra A by representations of a bimodule is the following: Let $\mathcal{F}$ and $\mathcal{T}$ be two full subaggregates of A−mod and $G : A\text{−mod} \to A\text{−mod}$ a subfunctor of the identity functor such that $G(X) \in \mathcal{T}$ and $X/G(X) \in \mathcal{F}$ for all X in A−mod. We consider the bimodule $H = \mathrm{Ext}^1_A(-,-)$ acting on $\mathcal{F} \times \mathcal{T}$ and obtain a full functor $R : A\text{−mod} \to \mathrm{rep}(H)$ by mapping the module X to its canonical exact sequence:

$$0 \to G(X) \xrightarrow{\iota_X} X \xrightarrow{\pi_X} X/G(X) \to 0$$

In general the functor R is neither dense nor faithful. Its kernel consists of the morphisms $f : X \to Y$ which factorise as $f = \iota_Y g \pi_X$. Since this kernel is contained in the radical of the aggregate $A-$mod we see that $A-$mod is representation equivalent to its image category inside $\operatorname{rep}(H)$. Thus we have 'reduced' the study of $A-$mod to the study of its image. It turns out that in many cases representations of a bimodule are easier to handle than the module category itself.

Let us provide some examples for choices of $\mathcal{F}$, $\mathcal{T}$ and G. We first start with a full subaggregate $\mathcal{T}$ of $A-$mod which is closed under factor modules. For an A-module X we define $G = G_{\mathcal{T}}(X)$ as the sum over all images $f(Z)$ for all Z in $\mathcal{T}$ and $f \in \operatorname{Hom}_A(Z, X)$. For $\mathcal{F}$ we choose a subaggregate of $A-$mod containing all modules $X/G(X)$. Another class of examples arises by considering an ideal I of A. We take $\mathcal{T}$ as the subcategory of A/J-modules and $\mathcal{F}$ as the subcategory of A/I-modules of $A-$mod where J is the left annihilator of I in A. The functor $G = G_I$ is defined as $G_I(X) = IX$. Dually, we can take $\mathcal{T} = A/I-$mod, $\mathcal{F} = A/J-$mod and $G = G^I$ as the annihilator of I in X where J is the right annihilator of I in A (this means $J = G^I(I)$).

Note, that the lack of density of R is repaired if one assumes that the pair $(\mathcal{F}, \mathcal{T})$ is a torsion theory or in other words G is a radical. Moreover, under this assumption R will also be faithful. Being a torsion class forces $\mathcal{T}$ to be closed under extensions and factor modules. Therefore, if $\mathcal{T} = \operatorname{add} S$, then $\operatorname{Ext}^1_A(S, S) = 0$.

2.3. Let us continue to consider the situation $\mathcal{T} = \operatorname{add} S$ where additionally S satisfies $\mathcal{T}(S, S) \cong k$. In this case $\operatorname{rep} H$ can be rewritten as a subspace category. Let us recall the relevant notation.

A vector space category is a k-additive functor $M \colon \mathcal{F} \to k-$mod where $\mathcal{F}$ is an aggregate. Its subspace category $\check{\mathcal{U}}(M)$ is an aggregate which has as objects the triples $U = (U_\omega, \gamma_U, X_U)$ where $U_\omega \in k-$mod, $U_0 \in \mathcal{F}$ and $\gamma_U \in \operatorname{Hom}_k(U_\omega, M(U_0))$. Morphisms $U \to U'$ in this category are pairs $f = (f_\omega, f_0)$ such that $f_\omega \colon U_\omega \to U'_\omega$ is k-linear, $f_0 \colon U_0 \to U'_0$ is a morphism in $\mathcal{F}$ and $M(f_0)\gamma_X = \gamma_{X'} f_\omega$.

If now H is a bimodule such that $\mathcal{T} = \operatorname{add} S$, then H is completely determined by the contravariant functor $M := H(-, S) \colon \mathcal{F} \to k-$mod which we consider as covariant functor $\mathcal{F}^{op} \to k-$mod. Moreover, any object of $\operatorname{rep}(H)$ lying in $H(X, S^n)$ may by the Yoneda lemma be identified with a morphism in $\operatorname{Hom}_{\mathcal{T}(S,S)}(\mathcal{T}(S^n, S), H(X, S))$. This identification yields an antiequivalence $\operatorname{rep}(H) \to \check{\mathcal{U}}(M)$.

Dually, one can transform $\operatorname{rep}(H)$ into a subspace category if $\mathcal{F} = \operatorname{add} S$. This happens e.g. for G^I introduced above for an ideal I satisfying $A/J \cong k$. This is used in [GNRSV] where the considered functor R usually will not be dense, but nevertheless the image can be calculated precisely.

3. Fibre sum functors

3.1. Let $(\mathcal{F}, \mathcal{T})$ be a torsion theory in $A-$mod. We define $\mathcal{K}_{\mathcal{T}}$ to be the full subaggregate of $A-$mod whose objects are the modules V satisfying $\operatorname{Ext}^1_A(V, \mathcal{T}) = 0$. The bimodule L acts on $\mathcal{T} \times \mathcal{K}_{\mathcal{T}}$ as $\operatorname{Hom}_A(-, -)$. We put $\operatorname{rep}_{mon}(L)$ to be the full subcategory of $\operatorname{rep}(L)$ given by all monomorphisms $h : X \to V$. Then the definition of $\mathcal{K}_{\mathcal{T}}$ implies that the functor $\widetilde{F} : \operatorname{rep}_{mon}(L) \to A-$mod which sends h to its cokernel is full.

As a torsion class $\mathcal{T}$ is closed under factor modules. Let us assume that $\mathcal{T}$ has a cover P i.e. $\mathcal{T} = \operatorname{fac} P$ for some module P in $A-$mod. Then we can even assume that P is a minimal cover and therefore $\operatorname{Ext}_(P, \operatorname{fac} P) = 0$. It follows from [AR] that $\widetilde{F}$ is dense.

The category $\mathrm{rep}_{mon}(L)$ seems hard to understand. To improve the situation we assume additionally that also $\mathrm{Ext}^1_A(P, \mathrm{sub}\, P) = 0$. Let C be the bimodule which acts as $\mathrm{Hom}_A(-,-)$ on $\mathrm{add}\, P \times \mathcal{K}_\mathcal{T}$. Then the functor $\Psi : \mathrm{rep}(C) \to \mathrm{rep}_{mon}(L)$ which sends $f : U \to V$ to the inclusion of $\mathrm{Im}\, f$ into V is full and dense. Altogether we derived the following result from [Dr2] where we put $F_P = \widetilde{F}\Psi$ which is said to be the fibre sum functor with respect to P.

Proposition. *Suppose the A-module P satisfies $\mathrm{Ext}^1_A(P, \mathrm{sub}\, P) = 0 = \mathrm{Ext}^1_A(P, \mathrm{fac}\, P)$. Then the fibre sum functor F_P is full and dense.*

One can also calculate that the kernel of F_P consists of the morphisms factoring through a finite collection of objects in $\mathrm{rep}(C)$.

3.2. Now we consider the special case that $\mathrm{End}_A(P) \cong k$. Then we can identify $\mathrm{rep}(C)$ with $\check{\mathcal{U}}(N)$ where N is the functor $\mathrm{Hom}_A(P,-)$ acting on $\mathcal{K}_\mathcal{T}$. Namely, a homomorphism f in $\mathrm{Hom}_A(P \otimes_k k^n, V)$ is mapped to its adjoint homomorphism $\hat{f}$ in $\mathrm{Hom}_k(k^n, \mathrm{Hom}_A(P,V))$.

If in addition $P = S$ is a simple module, then two functors relating A−mod with the subspace category of a vector space category were introduced namely $F_S : \check{\mathcal{U}}(N) \to A$−mod and the antiequivalence $R : A$−mod $\to \check{\mathcal{U}}(M)$. It is our final aim to calculate the composition RF_S. Note, that R is already an equivalence whereas F_P in general cannot be an equivalence because $F_P(k,0,0) = 0 = F(k, \mathrm{id}_k, P)$. Erasing these two objects makes also F_S into an equivalence. More precisely, we consider the full subcategory $K'_\mathcal{T}$ of $\mathcal{K}_\mathcal{T}$ whose objects do not admit a summand isomorphic to S and denote by N' the restriction of N to $K'_\mathcal{T}$. Considering only $\check{\mathcal{U}}(N')$ excludes direct summands of the form (k, id_k, S). Unfortunately, it cancels also direct summands of the form $(0,0,S)$ but, as we will see below, this is the price we have to pay to get satisfactory results. Furthermore, we replace $\check{\mathcal{U}}(N')$ by its full subcategory $\mathcal{U}(N')$ having only those objects U such that γ_U is a monomorphism. In this way we get rid of direct summands of the shape $(k,0,0)$. Now the restriction $F_S : \mathcal{U}(N') \to A$−mod$'$ is an equivalence where A−mod$'$ is the full subaggregate of modules X admitting now direct summand S. On the other hand, R maps A−mod$'$ onto $\mathcal{U}(M)^{op}$. Thus RF_S becomes an antiequivalence $\mathcal{U}(N') \to \mathcal{U}(M)$.

The best that can happen for such an antiequivalence of subspace categories is that it is induced by a suitable antiequivalence of the corresponding vector space categories. We first provide an equivalence $\mathcal{K}'_\mathcal{T} \to \mathcal{F}$.

Lemma. *Let G' be the functor sending X in A−mod to $X/G(X)$. Then G induces an equivalence $\mathcal{K}'_\mathcal{T} \to \mathcal{F}$.*

Proof. For given V in $\mathcal{K}'$ the adjoint of the inclusion of $G(V)$ into V is an object $U = (k^n, \gamma_U, V)$ satisfying $F_S(U) = G'(V)$. Therefore the density and fullness of F_S implies the required density and fullness of G'. That G' acts faithful on $\mathcal{K}'_\mathcal{T}$ follows easily because there do not exist non-zero maps from $\mathcal{K}'_\mathcal{T}$ to S. But note that $G'(S) = 0$. Here it pays out that we passed form $\mathcal{K}_\mathcal{T}$ to $\mathcal{K}'_\mathcal{T}$. □

Furthermore, we need a natural isomorphism between the involved functors. As usual $\mathrm{D} = \mathrm{Hom}_k(-,k)$ denotes the dual space functor.

Lemma. *There is a natural isomorphism $\varphi : \mathrm{D}\,\mathrm{Hom}_A(S,-) \to \mathrm{Ext}^1_A(G'(-), S)$ of functors $\mathcal{K}_\mathcal{T} \to k$−mod.*

Proof. Let us consider V in $\mathcal{K}'_{\mathcal{T}}$. The canonical exact sequence

$$0 \to G(V) \xrightarrow{\iota_V} V \xrightarrow{\pi_V} G'(V) \to 0$$

induces an exact sequence

$$\begin{aligned} \operatorname{Hom}_A(G'(V),S) &\xrightarrow{\operatorname{Hom}_A(\pi_V,S)} \operatorname{Hom}_A(V,S) \longrightarrow \operatorname{Hom}_A(G(V),S) \longrightarrow \\ &\longrightarrow \operatorname{Ext}^1_A(G'(V),S) \longrightarrow \operatorname{Ext}^1_A(V,S) = 0 \end{aligned}$$

where $\operatorname{Hom}_A(\pi_V,S)$ is actually an isomorphism because $\operatorname{Hom}_A(V,S)\iota_V = 0$. Hence we obtain an isomorphism $\operatorname{Hom}_A(G(V),S) \cong \operatorname{Ext}^1_A(G'(V),S)$. As $G(V)$ is in $\mathcal{T} = \operatorname{add} S$, we obtain $\operatorname{D}\operatorname{Hom}_A(G(V),S) \cong \operatorname{Hom}_A(S,G(V))$. Finally, $\operatorname{Hom}_A(S,G(V)) \cong \operatorname{Hom}_A(S,V)$ by the definition of G. The composition of all these isomorphisms yields the desired natural isomorphism φ. □

Using the two lemmas above and calculating RF_S we obtain:

Theorem. *The functor $\mathcal{U}(N') \to \mathcal{U}(M)$ sending an object $U = (U_\omega, \gamma_U, U_0)$ to $(\operatorname{D} U'_\omega, \varphi_{U_0}^{-1} \operatorname{D} \gamma'_U, G'(U_0))$ is an antiequivalence $\mathcal{U}(N') \to \mathcal{U}(M)$ which is isomorphic to RF_S.*

References

[AR] M. Auslander, I. Reiten, *Applications of contravariantly finite subcategories,* Adv. Math. 86 (1991), 111-152.

[C-B] W. Crawley-Boevey, *On tame algebras and bocses,* Proc. London Math. Soc. (3) 56 (1988), 451-483.

[Dr1] P. Dräxler, *$\mathcal{U}$-Fasersummen in darstellungsendlichen Algebren,* J. Algebra 113 (1988), 430-437.

[Dr2] P. Dräxler, *On the density of fiber sum functors,* Math. Z. 216 (1994), 645-656.

[Dd1] Y.A. Drozd, *Matrix problems and categories of matrices,* Zap. Nauchn. Sem. LOMI 28 (1972), 144-153

[Dd2] Ju. A. Drozd: *Tame and wild matrix problems,* Lecture Notes in Math. 832 (1980), 242-258.

[GNRSV] P. Gabriel, L.A. Nazarova, A.V. Roiter, V.V. Sergejchuk, D. Vossieck, *Tame and wild subspace problems,* Ukr. Math. J. 45 (1993), 313-352.

[GR] P. Gabriel, A.V. Roiter, *Representations of finite-dimensional algebras,* Encyclopedia of the Mathematical Sciences, Vol. 73, Algebra VIII, A.I. Kostrikin and I.V. Shafarevich (Eds.), Berlin, Heidelberg, New York, 1992.

[NR] L.A. Nazarova, A.V. Roiter, *Kategorielle Matrizenprobleme und die Brauer-Thrall-Vermutung,* Mitt. Math. Sem. Giessen 115 (1975).

[Ri1] C.M. Ringel, *Report on the Brauer-Thrall conjectures,* Lecture Notes in Math. 831 (1980), 104-136.

[Ri2] C.M. Ringel, *Tame algebras and integral quadratic forms,* Lecture Notes in Math. 1099 (1984).

Fakultät für Mathematik, Universität Bielefeld, POBox 100131, D-33501 Bielefeld, Germany