

# Mathematical Modelling and Simulation with Comsol Multiphysics II

## Winter term 2015/2016

### Exercise 4

Dr. Denny Otten



**Bearbeitung:** Montag, 02.11.2015, 12:30-14:00 Uhr (während der Übung).

#### Exercise 4 (FitzHugh-Nagumo system: Traveling Front).

Consider the **FitzHugh-Nagumo system**

$$\begin{pmatrix} u_{1,t} \\ u_{2,t} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & D \end{pmatrix} \begin{pmatrix} u_{1,xx} \\ u_{2,xx} \end{pmatrix} + \begin{pmatrix} u_1 - \zeta u_1^3 - u_2 + \alpha \\ \beta(\gamma u_1 - \delta u_2 + \varepsilon) \end{pmatrix}, \quad x \in \mathbb{R}, t \geq 0$$

for some  $D \geq 0$ ,  $\alpha, \beta, \gamma, \delta, \varepsilon, \zeta \in \mathbb{R}$ ,  $\zeta \neq 0$  and  $u_i = u_i(x, t) \in \mathbb{R}$  for  $i = 1, 2$ . Using the notation

$$u = (u_1, u_2) \in \mathbb{R}^2 \quad \text{and} \quad f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f(u) = \begin{pmatrix} u_1 - \zeta u_1^3 - u_2 + \alpha \\ \beta(\gamma u_1 - \delta u_2 + \varepsilon) \end{pmatrix},$$

the FitzHugh-Nagumo system can also be written as

$$u_t = \begin{pmatrix} 1 & 0 \\ 0 & D \end{pmatrix} u_{xx} + f(u), \quad x \in \mathbb{R}, t \geq 0.$$

a) Solve the **nonfrozen FitzHugh-Nagumo system**

$$(1) \quad \begin{aligned} \begin{pmatrix} u_{1,t} \\ u_{2,t} \end{pmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & D \end{pmatrix} \begin{pmatrix} u_{1,xx} \\ u_{2,xx} \end{pmatrix} + \begin{pmatrix} u_1 - \zeta u_1^3 - u_2 + \alpha \\ \beta(\gamma u_1 - \delta u_2 + \varepsilon) \end{pmatrix}, \quad x \in \Omega, t \in (0, T_1], \\ \begin{pmatrix} u_1(0) \\ u_2(0) \end{pmatrix} &= \begin{pmatrix} u_0^{(1)} \\ u_0^{(2)} \end{pmatrix}, \quad x \in \bar{\Omega}, t = 0, \\ \begin{pmatrix} u_{1,x} \\ u_{2,x} \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad x \in \partial\Omega, t \in [0, T_1], \end{aligned}$$

on the spatial domain  $\Omega = (-60, 60)$  for end time  $T_1 = 80$ , asymptotic states

$$u_- = \begin{pmatrix} 1.18769696080266 \\ 0.629232320266825 \end{pmatrix}, \quad u_+ = \begin{pmatrix} -1.56443178284120 \\ -0.288143927613547 \end{pmatrix},$$

initial data

$$u_0^{(1)}(x) = \begin{cases} u_-^{(1)} & , x < 0 \\ u_+^{(1)} & , x \geq 0 \end{cases}, \quad u_0^{(2)}(x) = \begin{cases} u_-^{(2)} & , x < 0 \\ u_+^{(2)} & , x \geq 0 \end{cases},$$

and parameters

$$D = \frac{1}{10}, \quad \alpha = 0, \quad \beta = \frac{2}{25}, \quad \gamma = 1, \quad \delta = 3, \quad \varepsilon = \frac{7}{10}, \quad \zeta = \frac{1}{3}.$$

For the space discretization use linear Lagrange elements with maximal element size  $\Delta x = 0.1$ . For the time discretization use the BDF method of maximum order 5 with intermediate time steps, time stepsize  $\Delta t = 0.1$ , relative tolerance  $rtol = 10^{-2}$  and absolute tolerance  $atol = 10^{-4}$  with global method set to be unscaled. The nonlinear equations should be solved by the Newton method. i.e. automatic (Newton).

b) Solve the **nonfrozen FitzHugh-Nagumo system**

$$(2) \quad \begin{aligned} \begin{pmatrix} \hat{v}_{1,t} \\ \hat{v}_{2,t} \end{pmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & D \end{pmatrix} \begin{pmatrix} \hat{v}_{1,xx} \\ \hat{v}_{2,xx} \end{pmatrix} + \begin{pmatrix} \hat{v}_1 - \zeta \hat{v}_1^3 - \hat{v}_2 + \alpha \\ \beta(\gamma \hat{v}_1 - \delta \hat{v}_2 + \varepsilon) \end{pmatrix}, \quad x \in \Omega, t \in (0, T_2], \\ \begin{pmatrix} \hat{v}_1(0) \\ \hat{v}_2(0) \end{pmatrix} &= \begin{pmatrix} \hat{v}_0^{(1)} \\ \hat{v}_0^{(2)} \end{pmatrix}, \quad x \in \bar{\Omega}, t = 0, \\ \begin{pmatrix} \hat{v}_{1,x} \\ \hat{v}_{2,x} \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad x \in \partial\Omega, t \in [0, T_2], \end{aligned}$$

on the spatial domain  $\Omega = (-60, 60)$  for end time  $T_2 = 2$ ,  $\hat{v}_0^{(1)}(x) = u_0^{(1)}(x)$ ,  $\hat{v}_0^{(2)}(x) = u_0^{(2)}(x)$  and parameters

$$D = \frac{1}{10}, \quad \alpha = 0, \quad \beta = \frac{2}{25}, \quad \gamma = 1, \quad \delta = 3, \quad \varepsilon = \frac{7}{10}, \quad \zeta = \frac{1}{3}.$$

For the space discretization use linear Lagrange elements with maximal element size  $\Delta x = 0.1$ . For the time discretization use the BDF method of maximum order 5 with intermediate time steps, time stepsize  $\Delta t = 0.1$ , relative tolerance  $rtol = 10^{-2}$  and absolute tolerance  $atol = 10^{-4}$  with global method set to be unscaled. The nonlinear equations should be solved by the Newton method. i.e. automatic (Newton).

c) Solve the **frozen FitzHugh-Nagumo system**

$$(3) \quad \begin{aligned} \begin{pmatrix} v_{1,t} \\ v_{2,t} \end{pmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & D \end{pmatrix} \begin{pmatrix} v_{1,\xi\xi} \\ v_{2,\xi\xi} \end{pmatrix} + \mu \begin{pmatrix} v_1 - \zeta v_1^3 - v_2 + \alpha \\ v_2 - \delta v_2 + \varepsilon \end{pmatrix}, \quad \xi \in \Omega, t \in (0, T_3], \\ \begin{pmatrix} v_1(0) \\ v_2(0) \end{pmatrix} &= \begin{pmatrix} v_0^{(1)} \\ v_0^{(2)} \end{pmatrix}, \quad \xi \in \bar{\Omega}, t = 0, \\ \begin{pmatrix} v_{1,\xi} \\ v_{2,\xi} \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad \xi \in \partial\Omega, t \in [0, T_3], \\ 0 &= (v_1 - \hat{v}_1, \hat{v}_{1,\xi})_{L^2(\Omega, \mathbb{R})} + (v_2 - \hat{v}_2, \hat{v}_{2,\xi})_{L^2(\Omega, \mathbb{R})}, \quad t \in [0, T_3], \\ \gamma_t &= \mu, \quad t \in (0, T_3], \\ \gamma(0) &= 0, \quad t = 0 \end{aligned}$$

on the spatial domain  $\Omega = (-60, 60)$  for end time  $T_3 = 100$ , initial data  $v_0^{(1)}(\xi) = u_0^{(1)}(\xi)$ ,  $v_0^{(2)}(\xi) = u_0^{(2)}(\xi)$ , reference function  $\hat{v}(\xi)$  as the solution of (2) at end time  $T_2$  and parameters

$$D = \frac{1}{10}, \quad \alpha = 0, \quad \beta = \frac{2}{25}, \quad \gamma = 1, \quad \delta = 3, \quad \varepsilon = \frac{7}{10}, \quad \zeta = \frac{1}{3}.$$

For the space discretization use linear Lagrange elements with maximal element size  $\Delta x = 0.1$ . For the time discretization use the BDF method of maximum order 2 with intermediate time steps, time stepsize  $\Delta t = 0.1$ , relative tolerance  $rtol = 10^{-2}$  and absolute tolerance  $atol = 10^{-5}$  with global method set to be unscaled. The nonlinear equations should be solved by the Newton method (automatic (Newton)).

d) Solve the **eigenvalue problem** for the linearization of the FitzHugh-Nagumo system

$$(4) \quad \begin{aligned} \lambda \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & D \end{pmatrix} \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} + \mu_* \begin{pmatrix} w_{1,\xi} \\ w_{2,\xi} \end{pmatrix} + Df(v_*(\xi)) \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad \xi \in \Omega, \\ \begin{pmatrix} w_{1,\xi} \\ w_{2,\xi} \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad \xi \in \partial\Omega \end{aligned}$$

on the spatial domain  $\Omega = (-60, 60)$ , where  $Df(u)$  denotes the derivative of  $f$  given by

$$Df(u) = \begin{pmatrix} 1 - 3\zeta u_1^2 & -1 \\ \beta\gamma & -\beta\delta \end{pmatrix}.$$

For  $v_*$  and  $\mu_*$  use the solutions  $v$  and  $\mu$  of (3) at the end time  $T_3 = 100$ . Determine  $\text{neigs} = 400$  eigenvalues  $\lambda$  and corresponding eigenfunctions  $w = (w_1, w_2)^T$ . The eigenvalues should be closest in absolute value around the shift  $-1$ .

e) **Postprocessing and Visualization** of results: Create the following plots to visualize the results of the computations:

- [Traveling Front, View 1](#): Plot the solution  $u_1$  of (1) at time  $t = 0, 20, 40$  and  $60$ .
- [Traveling Front, View 1](#): Plot the solution  $u_2$  of (1) at time  $t = 0, 20, 40$  and  $60$ .
- [Traveling Front, View 2](#): Create a time-space plot for the solution  $u_1$  of (1).
- [Traveling Front, View 2](#): Create a time-space plot for the solution  $u_2$  of (1).
- [Reference function](#): Plot the template solutions  $\hat{v}_1$  and  $\hat{v}_2$  of (2) at time  $T_2$ .
- [Profile, View 1](#): Plot the solution  $v_1$  and  $v_2$  of (3) at the end time  $T_3$ .
- [Profile, View 2](#): Create a time-space plot for the solution  $v_1$  of (3).
- [Profile, View 2](#): Create a time-space plot for the solution  $v_2$  of (3).
- [Velocity](#): Plot the velocity  $\mu$  of (3) for time  $t$  from  $0$  to  $T_3$ .
- [Position](#): Plot the position  $\gamma$  of (3) for time  $t$  from  $0$  to  $T_3$ .
- [Convergence indicator](#): Plot  $\|v_t(t)\|_{L^2(\Omega, \mathbb{R}^2)}$  and  $|\mu_t(t)|$  for time  $t$  from  $0$  to  $T_3$ .
- [Eigenvalues and Spectrum](#): Plot the eigenvalues  $\lambda$  of (4).
- [Eigenfunctions](#): Plot the eigenfunction  $w$  of (4) belonging to the zero eigenvalue.