Empty polygons in Penrose tilings

Dirk Frettlöh

Joint work with Alexey Garber


Technische Fakultät
Universität Bielefeld

Kolloquium über Kombinatorik

TU Berlin, 14^{th} Nov 2026


Empty lattice polygons:

("polygon" here always means convex polygon)



Empty lattice polygons:

("polygon" here always means convex polygon)



Several interesting results and applications.

One: Helly number of point lattices.


Helly numbers


Theorem (Radon '21, König '22, Helly '23)

Let $K_1, ..., K_n$ be convex subsets of $\mathbb{R}^d$, with $n \geqslant d+1$. If any $d+1$ of these sets have a common point, then all have a common point; that is

$$\bigcap_{j=1}^{n} K_j \neq \varnothing.$$

Hence $d+1$ is the **Helly number** $h(\mathbb{R}^d)$ of $\mathbb{R}^d$.





For instance, $h(\mathbb{R}^2) = 3$.

Helly numbers

Let us generalize Helly numbers to other sets than $\mathbb{R}^d$.

Definition

Let $S \subseteq \mathbb{R}^d$ be nonempty. The S-Helly number $h(S)$ of S is the minimal number $n > d$ such that the following statement holds:

Let $K_1, ..., K_n$ be convex subsets of S. If any $h(S)$ of these sets have a common point, then all have a common point.

Helly numbers

Let us generalize Helly numbers to other sets than $\mathbb{R}^d$.

Definition

Let $S \subseteq \mathbb{R}^d$ be nonempty. The S-Helly number $h(S)$ of S is the minimal number $n > d$ such that the following statement holds:

Let $K_1, ..., K_n$ be convex subsets of S. If any $h(S)$ of these sets have a common point, then all have a common point.


One result in this context:

Theorem (Doignon '73)

$h(\mathbb{Z}^d) = 2^d$.

Helly numbers

For instance, $h(\mathbb{Z}^2) = 4$.




Any three triangles have a common point, but not all four.

Helly numbers and empty polygons

Theorem (Averkov '13)

Let $S \subseteq \mathbb{R}^d$ be discrete. Then the Helly number $h(S)$ equals the maximal number of vertices of an empty S-polytope.

For instance $h(\mathbb{Z}^2) = 4$:




Helly numbers and empty polygons

Theorem (Averkov '13)

Let $S \subseteq \mathbb{R}^d$ be discrete. Then the Helly number $h(S)$ equals the maximal number of vertices of an empty S-polytope.


For instance $h(\mathbb{Z}^2) = 4$:



Which interesting discrete sets S we may study?


Penrose tilings

A famous aperiodic tiling: the Penrose Tiling




Penrose tilings

Consider its vertex set P:



Empty polygons in Penrose tilings

...and determine the empty polygons.



Empty polygons in Penrose tilings

By Averkov's result the maximal number of vertices yields the Helly number $h(P)$.

So far we have:

Empty polygons in Penrose tilings

By Averkov's result the maximal number of vertices yields the Helly number $h(P)$.

So far we have:

Theorem[Garber '16] $h(P) \leqslant 32$.

...and of course $h(P) \geqslant 6$.

Empty polygons in Penrose tilings

By Averkov's result the maximal number of vertices yields the Helly number $h(P)$.

So far we have:

Theorem[Garber '16] $h(P) \leqslant 32$.

...and of course $h(P) \geqslant 6$.

Our guess: $h(P) = 6$.

Empty polygons in Penrose tilings

By Averkov's result the maximal number of vertices yields the Helly number $h(P)$.

So far we have:

Theorem[Garber '16] $h(P) \leqslant 32$.


...and of course $h(P) \geqslant 6$.

Our guess: $h(P) = 6$.

Problem: there are arbitrarily long, thin empty polygons in P.


Empty polygons in Penrose tilings

Long and thin quadrangles:



Empty polygons in Ammann-Beenker tilings

Another semi-famous example:



Let A denote the vertex set of the Ammann-Beenker tiling.

Theorem[Garber '16] $h(A) \leqslant 16$.

...and of course $h(A) \geqslant 6$.

Our guess: $h(A) = 6$.

Let A denote the vertex set of the Ammann-Beenker tiling.

Theorem[Garber '16] $h(A) \leqslant 16$.

...and of course $h(A) \geqslant 6$.

Our guess: $h(A) = 6$.

So what do we have?

- An interesting topic,
- a simple proof of Averkov's theorem (not shown),
- some guesses,

Let A denote the vertex set of the Ammann-Beenker tiling.

Theorem[Garber '16] $h(A) \leqslant 16$.

...and of course $h(A) \geqslant 6$.

Our guess: $h(A) = 6$.

So what do we have?

- An interesting topic,
- a simple proof of Averkov's theorem (not shown),
- some guesses,
- a visiting postdoc who will work on this,
- AI?

Let A denote the vertex set of the Ammann-Beenker tiling.

Theorem[Garber '16] $h(A) \leqslant 16$.

...and of course $h(A) \geqslant 6$.

Our guess: $h(A) = 6$.

So what do we have?

- An interesting topic,
- a simple proof of Averkov's theorem (not shown),
- some guesses,
- a visiting postdoc who will work on this,
- AI?

Thank you!