

# Bi-Lipschitz equivalence and bounded distance equivalence of Delone sets

Dirk Frettlöh

Technische Fakultät  
Universität Bielefeld

3.12.2015

joint work with Alexey Garber

- ▶ Basics
- ▶ Dimension 1
- ▶ Higher dimensions

*Delone set:* point set  $\Lambda$  in  $\mathbb{R}^d$ , with  $R > r > 0$  such that


- ▶ each ball of radius  $r$  contains at most one point of  $\Lambda$   
(*uniformly discrete*)
- ▶ each ball of radius  $R$  contains at least one point of  $\Lambda$   
(*relatively dense*)

(Aka “separated nets”. Can also live in  $\mathbb{H}^d$ ,  $(\mathbb{Q}_p)^d$  ...)


**Delone set:** point set  $\Lambda$  in  $\mathbb{R}^d$ , with  $R > r > 0$  such that

- ▶ each ball of radius  $r$  contains at most one point of  $\Lambda$  (*uniformly discrete*)
- ▶ each ball of radius  $R$  contains at least one point of  $\Lambda$  (*relatively dense*)

(Aka “separated nets”. Can also live in  $\mathbb{H}^d$ ,  $(\mathbb{Q}_p)^d$  ...)



crystallographic



disordered

**Two relations** between Delone sets:

$\Lambda \stackrel{\text{bil}}{\sim} \Lambda'$  (*bilipschitz equivalent*):

There is  $f : \Lambda \rightarrow \Lambda'$  bijective with

$$\exists c > 0 \quad \forall x, y \in \Lambda \quad \frac{1}{c}|x - y| \leq |f(x) - f(y)| \leq c|x - y|$$

(i.e.  $f$  and  $f^{-1}$  are Lipschitz continuous)

**Two relations** between Delone sets:

$\Lambda \overset{\text{bil}}{\sim} \Lambda'$  (*bilipschitz equivalent*):

There is  $f : \Lambda \rightarrow \Lambda'$  bijective with

$$\exists c > 0 \quad \forall x, y \in \Lambda \quad \frac{1}{c}|x - y| \leq |f(x) - f(y)| \leq c|x - y|$$


(i.e.  $f$  and  $f^{-1}$  are Lipschitz continuous)

$\Lambda \overset{\text{bd}}{\sim} \Lambda'$  (*bounded distance equivalent*):


There is  $g : \Lambda \rightarrow \Lambda'$  bijective with

$$\exists C > 0 \quad \forall x \in \Lambda : \quad |x - g(x)| < C$$

**Example:** Two rectangular lattices  $\Lambda, \Lambda'$ . Is  $\Lambda \stackrel{\text{bd}}{\approx} \Lambda'$ ?




**Example:** Two rectangular lattices  $\Lambda, \Lambda'$ . Is  $\Lambda \stackrel{\text{bd}}{\approx} \Lambda'$ ?







**Example:** Two rectangular lattices  $\Lambda, \Lambda'$ . Is  $\Lambda \stackrel{\text{bd}}{\approx} \Lambda'$ ?



**Example:** Two rectangular lattices  $\Lambda, \Lambda'$ . Is  $\Lambda \stackrel{\text{bd}}{\approx} \Lambda'$ ?



**Example:** Two rectangular lattices  $\Lambda, \Lambda'$ . Is  $\Lambda \stackrel{\text{bd}}{\approx} \Lambda'$ ?



## Some basic results

### Lemma (1)

*Bilipschitz equivalence and bounded distance equivalence are equivalence relations.*

### Lemma (2)

*Let  $\Lambda, \Lambda'$  be Delone sets in  $\mathbb{R}^d$ . If  $\Lambda \stackrel{\text{bd}}{\sim} \Lambda'$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \Lambda'$ .*

# Warmup: Dimension 1

## Lemma (3)

Let  $\Lambda, \Lambda'$  be Delone sets in  $\mathbb{R}$ . Then  $\Lambda \stackrel{\text{bil}}{\sim} \Lambda'$ .


# Warmup: Dimension 1

## Lemma (3)

Let  $\Lambda, \Lambda'$  be Delone sets in  $\mathbb{R}$ . Then  $\Lambda \stackrel{\text{bil}}{\sim} \Lambda'$ .

*Proof (by image):* Show  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}$ .

Let  $\Lambda = \{\dots, \lambda_{-1}, \lambda_0, \lambda_1, \lambda_2, \dots\}$ ,  $\lambda_i < \lambda_{i+1}$



Let  $\Lambda, \Lambda' \subset \mathbb{R}$ . When is  $\Lambda \stackrel{\text{bd}}{\sim} \Lambda'$ ? Always? No:

Examples:

- ▶  $\{\dots - 3, -2, -1, 0, 1, 2, 3, \dots\} \not\stackrel{\text{bd}}{\sim} \{\dots, -6, -4, -2, 0, 2, 4, 6, \dots\}$
- ▶  $\{\dots - 3, -2, -1, 0, 2, 4, 6, \dots\} \not\stackrel{\text{bd}}{\sim} \{\dots - 6, -4, -2, 0, 1, 2, 3, \dots\}$

Let  $\Lambda, \Lambda' \subset \mathbb{R}$ . When is  $\Lambda \stackrel{\text{bd}}{\sim} \Lambda'$ ? Always? No:

Examples:

- ▶  $\{\dots - 3, -2, -1, 0, 1, 2, 3, \dots\} \not\stackrel{\text{bd}}{\sim} \{\dots, -6, -4, -2, 0, 2, 4, 6, \dots\}$
- ▶  $\{\dots - 3, -2, -1, 0, 2, 4, 6, \dots\} \not\stackrel{\text{bd}}{\sim} \{\dots - 6, -4, -2, 0, 1, 2, 3, \dots\}$

Density matters. Preliminary definition:

$$\text{dens}(\Lambda) := \lim_{r \rightarrow \infty} \frac{1}{2r} \#(\Lambda \cap [-r, r]),$$

if it exists.



Let  $\Lambda, \Lambda' \subset \mathbb{R}$ . When is  $\Lambda \stackrel{\text{bd}}{\sim} \Lambda'$ ? Always? No:


Examples:

- ▶  $\{\dots - 3, -2, -1, 0, 1, 2, 3, \dots\} \not\stackrel{\text{bd}}{\sim} \{\dots, -6, -4, -2, 0, 2, 4, 6, \dots\}$
- ▶  $\{\dots - 3, -2, -1, 0, 2, 4, 6, \dots\} \stackrel{\text{bd}}{\sim} \{\dots - 6, -4, -2, 0, 1, 2, 3, \dots\}$

Density matters. Preliminary definition:

$$\text{dens}(\Lambda) := \lim_{r \rightarrow \infty} \frac{1}{2r} \#(\Lambda \cap [-r, r]),$$

if it exists. Does not need to exist:



Oscillates between  $\frac{2}{3}$  and  $\frac{5}{6}$ .

**Question:** If  $\text{dens}(\Lambda) = \text{dens}(\Lambda')$ , is  $\Lambda \stackrel{\text{bd}}{\sim} \Lambda'$ ?

**Lemma**

*Let  $\Lambda, \Lambda'$  be periodic. Then  $\text{dens}(\Lambda) = \text{dens}(\Lambda')$  implies  $\Lambda \stackrel{\text{bd}}{\sim} \Lambda'$ .*

**Interesting examples are non-periodic.**

**Question:** If  $\text{dens}(\Lambda) = \text{dens}(\Lambda')$ , is  $\Lambda \stackrel{\text{bd}}{\sim} \Lambda'$ ?

**Lemma**

Let  $\Lambda, \Lambda'$  be periodic. Then  $\text{dens}(\Lambda) = \text{dens}(\Lambda')$  implies  $\Lambda \stackrel{\text{bd}}{\sim} \Lambda'$ .

**Interesting examples are non-periodic.**

**Theorem (Kesten 1966)**

Let  $\xi \in [0, 1]$ ,  $0 \leq a < b \leq 1$  and define


$$\Lambda := \{k \in \mathbb{Z} \mid a \leq (k\xi \bmod 1) < b\}.$$

Then  $D(n) := \#(\Lambda \cap [1, n]) - n(b - a)$  is bounded, if and only if  $b - a = k\xi \bmod 1$  for some  $k \in \mathbb{Z}$ .

Choose  $\xi \in [0, 1]$  irrational, let  $0 < b \leq 1$  and define


$$\Lambda_b := \{k \in \mathbb{Z} \mid 0 \leq (k\xi \bmod 1) < b\}.$$

Then  $D(n) := \#(\Lambda \cap [1, n]) - nb$  is bounded, if and only if  $b = k\xi \bmod 1$  for some  $k \in \mathbb{Z}$ .




The image shows  $\{(k, k\xi \bmod 1) \mid k = 0, 1, 2, \dots\}$ .

*Proof (by image) of if-part:* (F-Gähler 2011, Duneau-Oguey 1990):



*Proof (by image) of if-part:* (F-Gähler 2011, Duneau-Oguey 1990):



In particular Kesten yields Delone sets  $\Lambda_b$  that are not bounded distance equivalent to any  $c\mathbb{Z}$ . Even when  $\text{dens}(\Lambda_b)$  exists!

# Higher dimensional spaces

## Theorem (Bogopolski 1997)

*Any two Delone sets in  $\mathbb{H}^d$  ( $d \geq 2$ ) are bounded distance equivalent, hence bilipschitz equivalent.*

## Theorem (Burago-Kleiner, C. McMullen 1998)

*There are Delone sets  $\Lambda$  in  $\mathbb{R}^d$  such that  $\Lambda \not\stackrel{\text{bil}}{\sim} \mathbb{Z}^d$ .*

# Higher dimensional spaces

## Theorem (Bogopolski 1997)

*Any two Delone sets in  $\mathbb{H}^d$  ( $d \geq 2$ ) are bounded distance equivalent, hence bilipschitz equivalent.*

## Theorem (Burago-Kleiner, C. McMullen 1998)


*There are Delone sets  $\Lambda$  in  $\mathbb{R}^d$  such that  $\Lambda \not\stackrel{\text{bil}}{\sim} \mathbb{Z}^d$ .*

Cool! Let us study some problems in this field. E.g.

- ▶ Are the vertices of the Penrose tiling bounded distance equivalent to some lattice?
- ▶ How many equivalence classes wrt  $\stackrel{\text{bd}}{\sim}$  resp.  $\stackrel{\text{bil}}{\sim}$  ?



**Recall:** Interesting examples are non-periodic.  
Like the Penrose tiling:



## Theorem (F-Garber 2011 unpublished)

If  $\Lambda$  is a linearly repetitive Delone set in  $\mathbb{R}^2$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}^2$ .

**Repetitive:** there is  $R$  such that congruent copies of any local patch in  $\Lambda$  occur in every ball of radius  $R$ .

**Linear repetitive:**  $R$  depends linearly on the diameter of the patch.

## Theorem (F-Garber 2011 unpublished)

If  $\Lambda$  is a linearly repetitive Delone set in  $\mathbb{R}^2$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}^2$ .

**Repetitive:** there is  $R$  such that congruent copies of any local patch in  $\Lambda$  occur in every ball of radius  $R$ .

**Linear repetitive:**  $R$  depends linearly on the diameter of the patch.

## Theorem (Aliste-Prieto, Coronel, Gambaudo 2011)

If  $\Lambda$  is a linearly repetitive Delone set in  $\mathbb{R}^d$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}^d$ .

## Theorem (F-Garber 2011 unpublished)

If  $\Lambda$  is a linearly repetitive Delone set in  $\mathbb{R}^2$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}^2$ .

**Repetitive:** there is  $R$  such that congruent copies of any local patch in  $\Lambda$  occur in every ball of radius  $R$ .

**Linear repetitive:**  $R$  depends linearly on the diameter of the patch.

## Theorem (Aliste-Prieto, Coronel, Gambaudo 2011)

If  $\Lambda$  is a linearly repetitive Delone set in  $\mathbb{R}^d$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}^d$ .

## Corollary (F-Garber 2011 unpublished)

Let  $\Lambda_P$  be the vertices of the Penrose tiling.  $\Lambda_P \stackrel{\text{bil}}{\sim} \mathbb{Z}^2$ .

### Theorem (F-Garber 2011 unpublished)

If  $\Lambda$  is a linearly repetitive Delone set in  $\mathbb{R}^2$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}^2$ .

**Repetitive:** there is  $R$  such that congruent copies of any local patch in  $\Lambda$  occur in every ball of radius  $R$ .

**Linear repetitive:**  $R$  depends linearly on the diameter of the patch.

### Theorem (Aliste-Prieto, Coronel, Gambaudo 2011)

If  $\Lambda$  is a linearly repetitive Delone set in  $\mathbb{R}^d$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}^d$ .

### Corollary (F-Garber 2011 unpublished)

Let  $\Lambda_P$  be the vertices of the Penrose tiling.  $\Lambda_P \stackrel{\text{bil}}{\sim} \mathbb{Z}^2$ .

### Theorem (Solomon 2007)

$\Lambda_P \stackrel{\text{bd}}{\sim} c\mathbb{Z}^2$ .

### Theorem (F-Garber 2011 unpublished)

If  $\Lambda$  is a linearly repetitive Delone set in  $\mathbb{R}^2$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}^2$ .

**Repetitive:** there is  $R$  such that congruent copies of any local patch in  $\Lambda$  occur in every ball of radius  $R$ .

**Linear repetitive:**  $R$  depends linearly on the diameter of the patch.

### Theorem (Aliste-Prieto, Coronel, Gambaudo 2011)

If  $\Lambda$  is a linearly repetitive Delone set in  $\mathbb{R}^d$ , then  $\Lambda \stackrel{\text{bil}}{\sim} \mathbb{Z}^d$ .

### Corollary (F-Garber 2011 unpublished)

Let  $\Lambda_P$  be the vertices of the Penrose tiling.  $\Lambda_P \stackrel{\text{bil}}{\sim} \mathbb{Z}^2$ .


### Theorem (Solomon 2007)

$\Lambda_P \stackrel{\text{bd}}{\sim} c\mathbb{Z}^2$ .

### Theorem (Deuber-Simonovits-Sos 1995)


$\Lambda_P \stackrel{\text{bd}}{\sim} c\mathbb{Z}^2$ .

Well, then let's generalise Kesten to  $\mathbb{R}^d$  (at least "if"-part)



(cut-and-project sets, aka model sets, "mathematical quasicrystals")

Well, then let's generalise Kesten to  $\mathbb{R}^d$  (at least "if"-part)




(cut-and-project sets, aka model sets, "mathematical quasicrystals")

Other colleagues had the same idea: Haynes 2013,  
Haynes-Kxxx-Weiss 2014.



Well, then let's generalise Kesten to  $\mathbb{R}^d$  (at least "if"-part)



(cut-and-project sets, aka model sets, "mathematical quasicrystals")

Other colleagues had the same idea: Haynes 2013,  
Haynes-Kxxx-Weiss 2014.

Last October I've learned from Alan Haynes that this was done already in

C. Godrèche and C. Oguey:

Construction of average lattices for quasiperiodic structures by the section method, *J. Phys. France* 51 (1990) 21-37

Regarding question 2:

How many equivalence classes wrt  $\overset{\text{bd}}{\sim}$  resp.  $\overset{\text{bil}}{\sim}$  ?

**Theorem (Magazinov 2010)**

*There are  $|\mathbb{R}|$  equivalence classes of Delone sets in  $\mathbb{R}^d$  wrt  $\overset{\text{bil}}{\sim}$ .*

**Theorem (Garber 2009)**

*There are  $|\mathbb{R}|$  equivalence classes of Delone sets in  $\mathbb{R}^d$  wrt  $\overset{\text{bd}}{\sim}$ .*

Regarding question 2:

How many equivalence classes wrt  $\overset{\text{bd}}{\sim}$  resp.  $\overset{\text{bil}}{\sim}$  ?

**Theorem (Magazinov 2010)**

*There are  $|\mathbb{R}|$  equivalence classes of Delone sets in  $\mathbb{R}^d$  wrt  $\overset{\text{bil}}{\sim}$ .*

**Theorem (Garber 2009)**

*There are  $|\mathbb{R}|$  equivalence classes of Delone sets in  $\mathbb{R}^d$  wrt  $\overset{\text{bd}}{\sim}$ .*

*Proof.* It is easy to show that each Delone set in  $\mathbb{R}^d$  is bounded distance equivalent to some subset of  $r\mathbb{Z}^d$ , where  $r$  is the radius of uniformly discreteness.

$(|\mathbb{R}| \text{ many values of } r) \times (|\mathbb{R}| \text{ many subsets of } \mathbb{Z}^d) = |\mathbb{R}|.$

Further research:

- ▶ "Only if"-part of Kesten's Theorem
- ▶ Let  $\Lambda_1 \stackrel{\text{bil}}{\sim} \Lambda_2$ . Is  $\Lambda_2 \stackrel{\text{bil}}{\sim} \Lambda_1 \cup \Lambda_2$ ? Under which conditions?
- ▶ Let  $\Lambda \stackrel{\text{bd}}{\sim} \mathbb{Z}^2$ ,  $\Lambda = \Lambda_1 \cup \Lambda_2$ ,  $\Lambda_1 \stackrel{\text{bd}}{\sim} \Lambda_2$ . Is  $\Lambda_1 \stackrel{\text{bd}}{\sim} 2\mathbb{Z}^d$ ?

The background of the slide is a complex fractal pattern, resembling a Sierpinski triangle or a similar self-similar structure. It is composed of many small, interconnected shapes in various colors including blue, red, yellow, and white, set against a light gray background. The text "Thank you." is centered in a large, bold, brown font.

**Thank you.**