

# VOLUME GROWTH AND ESCAPE RATE OF BROWNIAN MOTION ON A CARTAN-HADAMARD MANIFOLD

ALEXANDER GRIGOR'YAN AND ELTON P. HSU

ABSTRACT. We prove an upper bound for the escape rate of Brownian motion on a Cartan-Hadamard manifold in terms of the volume growth function. One of the ingredients of the proof is the Sobolev inequality on such manifolds.

## CONTENTS

1. Introduction	1
2. Heat equation solution estimates	4
3. Escape rate of Brownian motion	7
4. Escape rate on model manifolds	10
4.1. Constant curvature	10
4.2. General model manifolds	11
References	14

## 1. INTRODUCTION

Let  $M$  be a geodesically complete noncompact Riemannian manifold. We denote by  $d(x, y)$  the geodesic distance between  $x$  and  $y$  and by  $\mu$  the Riemannian volume measure. We use  $\mathbb{P}_x$  to denote the diffusion measure generated by the Laplace-Beltrami operator  $\Delta$ . Let  $X = \{X_t, t \in \mathbb{R}_+\}$  be the coordinate process on the path space  $W(M) = C(\mathbb{R}_+, M)$ . By definition,  $\mathbb{P}_x$  is a probability measure on  $W(M)$  under which  $X$  is a Brownian motion starting from  $x$ .

Fix a reference point  $z \in M$  and let  $\rho(x) = d(x, z)$ . We say that a function  $R(t)$  is an *upper rate function* for Brownian motion on  $M$  if

$$\mathbb{P}_z \{\rho(X_t) \leq R(t) \text{ for all sufficiently large } t\} = 1.$$

The purpose of this paper is to study the rate of escape of Brownian motion on  $M$  in terms the volume growth function. Let us first point out that the notion of an upper rate function makes sense only if the lifetime of Brownian motion is infinite. In this case, the manifold  $M$  is called stochastically complete. The stochastic completeness is equivalent to the identity

$$\int_M p(t, x, y) d\mu(y) = 1,$$

---

2000 *Mathematics Subject Classification.* Primary 58J65; secondary 60J60, 35K05.

*Key words and phrases.* Escape rate, Brownian motion, Cartan-Hadamard manifold, volume growth.

where  $p(t, x, y)$  is the minimal heat kernel on  $M$ , which is also the transition density function of Brownian motion on  $M$ .

Let  $B(z, R)$  be the geodesic ball of radius  $R$  centered at  $z$ . It was proved in [3] that  $M$  is stochastically complete if

$$(1.1) \quad \int^{\infty} \frac{r \, dr}{\log \mu(B(z, r))} = \infty.$$

The integral in (1.1) will be used in this paper to construct an upper rate function. Before we state the result, let us briefly survey the existing estimates of escape rate.

- The classical Khinchin's law of the iterated logarithm says that for a Brownian motion in  $\mathbb{R}^n$  with probability 1

$$\limsup_{t \rightarrow \infty} \frac{\rho(X_t)}{\sqrt{4t \log \log t}} = 1$$

(the factor 4 instead of the classical 2 appears because in our setting a Brownian motion is generated by  $\Delta$  rather than  $\frac{1}{2}\Delta$ ). It follows that, for any  $\varepsilon > 0$ ,

$$(1.2) \quad R(t) = \sqrt{(4 + \varepsilon)t \log \log t}$$

is an upper rate function.

- If  $M$  has non-negative Ricci curvature then (1.2) is again an upper rate function on  $M$  – see [9, Theorem 1.3] and [7, Theorem 4.2].
- If the volume growth function is at most polynomial, that is,

$$\mu(B(z, r)) \leq Cr^D$$

for large enough  $r$  and for some positive constants  $C, D$  then the function

$$(1.3) \quad R(t) = \text{const} \sqrt{t \log t}$$

is an upper rate function – see [15, Theorem 5.1], [7, Theorem 1.1], and [9, Theorem 1.1]. Note that the logarithm in (1.3) is single in contrast to (1.2), and in general cannot be replaced by the iterated logarithm (cf. [1] and [10]).

- If the volume growth function admits a sub-Gaussian exponential estimate

$$\mu(B(z, r)) \leq \exp(Cr^\alpha),$$

where  $0 < \alpha < 2$ , then the function

$$R(t) = \text{const} t^{\frac{1}{2-\alpha}}$$

is an upper rate function – see [7, Theorem 4.1].

Note that (1.1) is satisfied if the volume growth function admits the Gaussian exponential estimate:

$$(1.4) \quad \mu(B(z, r)) \leq \exp(Cr^2)$$

(under the condition (1.4), the stochastic completeness was also proved by different methods in [15], [12], and [2]. However, none of the existing results provided any estimates of escape rate under the condition (1.4), let alone under the volume growth function  $\exp(Cr^2 \log r)$  and the like.

We will construct an upper rate function under the most general condition (1.1). However, we assume in addition that  $M$  is a Cartan-Hadamard manifold, that is,

a geodesically complete simply connected Riemannian manifold of non-positive sectional curvature. The property of Cartan-Hadamard manifolds that we use is the Sobolev inequality: if  $N = \dim M$  then, for any function  $f \in C_0^\infty(M)$ ,

$$(1.5) \quad \left( \int_M |f|^{\frac{N}{N-1}} d\mu \right)^{\frac{N-1}{N}} \leq C_N \int_M |\nabla f| d\mu,$$

where  $C_N$  is a constant depending only on  $N$  – see [11]. The Sobolev inequality allows us to carry through the Moser iteration argument in [14] and prove a mean value estimate for solutions of the heat equation on  $M$ , which is one of the ingredients of our proof.

Now we state our main result.

**Theorem 1.** *Let  $M$  be a Cartan-Hadamard manifold. Assume that the following volume estimate holds for a fixed point  $z \in M$  and all sufficiently large  $R$ :*

$$(1.6) \quad \mu(B(z, R)) \leq \exp(f(R)),$$

where  $f(R)$  is a positive, strictly increasing, and continuous function on  $[0, +\infty)$  such that

$$(1.7) \quad \int_0^\infty \frac{r dr}{f(r)} = \infty.$$

Let  $\phi(t)$  be the function on  $\mathbb{R}_+$  defined by

$$(1.8) \quad t = \int_0^{\phi(t)} \frac{r dr}{f(r)}.$$

Then  $R(t) = \phi(Ct)$  is an upper rate function for Brownian motion on  $M$  for some absolute constant  $C$  (for example, for any  $C > 128$ ).

If we set  $f(R) = \log \mu(B(z, R))$  for large  $R$  then the condition (1.7) becomes identical to (1.1). Under this condition, Theorem 1 guarantees the existence of an upper rate function  $R(t)$ . This in particular means that in a finite time Brownian motion stays with probability one in a bounded set, which implies that the life time of Brownian motion is infinite almost surely. Hence, the manifold  $M$  is stochastically complete. This recovers the cited above result that (1.1) on geodesically complete manifolds implies the stochastic completeness, although under the additional assumption that  $M$  is Cartan-Hadamard.

Let us show some examples.

- If

$$(1.9) \quad \mu(B(z, R)) \leq CR^D$$

for some constant  $C$  and  $D$  then (1.6) holds with

$$f(R) = D \log R + \text{const},$$

and (1.8) yields

$$t \simeq \frac{\phi^2}{2D \log \phi}.$$

It follows that  $\log t \simeq \log \phi^2$  and

$$\phi(t) \simeq \sqrt{Dt \log t}.$$

Hence, the following function

$$R(t) = \sqrt{CDt \log t}$$

is a upper rate function, which matches the above cited results of [7], [9], [15].

- If  $\mu(B(z, R)) \leq \exp(Cr^\alpha)$  for some  $0 < \alpha < 2$  then (1.6) holds with  $f(R) = Cr^\alpha$ , and (1.8) yields  $t \simeq \phi(t)^{2-\alpha}$  whence we obtain the upper rate function

$$R(t) = Ct^{\frac{1}{2-\alpha}},$$

which matches the above cited result of [7].

- If

$$\mu(B(z, R)) \leq \exp(CR^2),$$

then  $f(R) = CR^2$ . Then (1.8) yields  $t \simeq \ln \phi(t)$ . Hence, we obtain the upper rate function

$$R(t) = \exp(Ct).$$

This result is new. Similarly, if

$$\mu(B(z, R)) \leq \exp(CR^2 \log R)$$

then (1.8) yields  $t \simeq \log \log \phi$  whence

$$R(t) = \exp(\exp(Ct)).$$

The hypothesis that  $M$  is Cartan-Hadamard can be replaced by the requirement that the Sobolev inequality (1.5) holds on  $M$ . Furthermore, the method goes through also in the setting of weighted manifolds, when measure  $\mu$  is not necessarily the Riemannian measure but has a smooth positive density, say  $\sigma(x)$ , with respect to the Riemannian measure. Then instead of the Laplace-Beltrami operator, one should consider the weighted Laplace operator

$$\Delta_\mu = \frac{1}{\sigma} \operatorname{div}(\sigma \nabla),$$

which is symmetric with respect to  $\mu$ . Theorem 1 extends to the weighted manifolds that are geodesically complete and satisfy the Sobolev inequality (1.5).

This paper is organized as follows. Section 2 contains the proof of an upper bound for certain positive solutions of the heat equation. In Section 3 we prove the main result stated above. In Section 4 we compute the sharp upper rate function on model manifold and show that, for a certain range of volume growth functions, the upper rate function of Theorem 1 is sharp up to a constant factor in front of  $t$ .

## 2. HEAT EQUATION SOLUTION ESTIMATES

In this section we prove a pointwise upper bound of certain solutions of the heat equation on a Cartan-Hadamard manifold  $M$  (Theorem 4). It is an easy consequence of an  $L^2$ -bound for a general complete manifold and a mean value type inequality for a Cartan-Hadamard manifold. These two upper bounds are known and we will state them as lemmas.

For any set  $A \subset M$ , let  $A_r$  be the open  $r$ -neighborhood of  $A$  in  $M$ .

**Lemma 2.** *Let  $M$  be a geodesically complete Riemannian manifold. Suppose that  $u(x, t)$  is a smooth subsolution to the heat equation in the cylinder  $A_r \times [0, T]$ , where  $A \subset M$  is a compact set and  $r, T > 0$  (see FIG. 1). Assume also that  $0 \leq u(x, t) \leq 1$  and  $u(x, 0) = 0$  on  $A_r$ . Then, for any  $t \in (0, T]$ ,*

$$\int_A u^2(x, t) d\mu(x) \leq \mu(A_r) \max\left(1, \frac{r^2}{2t}\right) \exp\left(-\frac{r^2}{2t} + 1\right).$$




FIGURE 1. Illustration to Lemma 2

For the proof see [6, Theorem 3] (cf. also [7, Proposition 3.6]). Note that no geometric assumption about  $M$  is made except for the geodesic completeness. The proof exploits essentially the property of the geodesic distance function that  $|\nabla d| \leq 1$ .

Takeda [15] proved a similar estimate for  $\int_A u(x, t) d\mu(x)$ , using a different probabilistic argument. However, it is more convenient for us to work with the  $L^2$  rather than with  $L^1$  estimate in view of the following lemma.

**Lemma 3.** *Let  $M$  be a Cartan-Hadamard manifold of dimension  $N$ . Suppose that  $u(x, t)$  is a smooth nonnegative subsolution to the heat equation in a cylinder  $B(y, r) \times [0, T]$ , where  $r, T > 0$  (see FIG. 2). Then*

$$(2.1) \quad u(y, T)^2 \leq \frac{C_N}{\min(\sqrt{T}, r)^{N+2}} \int_0^T \int_{B(y, r)} u^2(x, t) d\mu(x) dt,$$

where  $C_N$  is a constant depending only on  $N$ .

*Proof.* As was already mentioned above, a Cartan-Hadamard manifold admits the Sobolev inequality (1.5). By a standard argument, (1.5) implies the Sobolev-Moser inequality

$$\int_M |f|^{2+\frac{4}{N}} d\mu \leq C_N \left( \int_M |f|^2 d\mu \right)^{2/N} \int_M |\nabla f|^2 d\mu,$$

which leads by the Moser iteration argument [14] to the mean value inequality (2.1). Note that the value of  $C_N$  may be different in all the above inequalities.




FIGURE 2. Illustration to Lemma 3

An alternative proof of the implication (1.5) $\Rightarrow$ (2.1) can be found in [4] (cf. also [5, Theorem 3.1 and (3.4)]).  $\square$

With these two preliminary results, we prove the main inequality in this section. This inequality will give an upper bound for probabilities of escaping times not exceeding a given upper bound.

**Theorem 4.** *Let  $M$  be a Cartan-Hadamard manifold of dimension  $N$ . Suppose that  $u(x, t)$  is a smooth subsolution to the heat equation in a cylinder  $B(y, 2r) \times [0, T]$ , where  $r, T > 0$ . If  $0 \leq u \leq 1$  in this cylinder and  $u(x, 0) = 0$  on  $B(y, 2r)$ , then*

$$(2.2) \quad u(y, T) \leq C_N \sqrt{\mu(B(y, 2r))} \frac{\max(\sqrt{T}, r)}{\min(\sqrt{T}, r)^{1+N/2}} \exp\left(-\frac{r^2}{4T}\right).$$

*Proof.* We use Lemma 2 with  $A = \overline{B(y, r)}$ . Then  $A_r = B(y, 2r)$  and, for any  $0 < t \leq T$ ,

$$\int_{B(y, r)} u^2(x, t) d\mu(x) \leq \mu(B(y, 2r)) \max\left(1, \frac{r^2}{2t}\right) \exp\left(-\frac{r^2}{2t} + 1\right)$$

(see FIG. 3), whence

$$\begin{aligned} \int_{T/2}^T \int_{B(y, r)} u^2(x, t) d\mu(x) dt \\ \leq 2\mu(B(y, 2r)) T \max\left(1, \frac{r^2}{T}\right) \exp\left(-\frac{r^2}{2T}\right). \end{aligned}$$

Applying Lemma 3 in the cylinder  $B(y, r) \times [T/2, T]$ , we obtain

$$u(y, T)^2 \leq C_N \mu(B(y, 2r)) \frac{\max(T, r^2)}{\min(\sqrt{T}, r)^{N+2}} \exp\left(-\frac{r^2}{2T}\right),$$

whence (2.2) follows.  $\square$




FIGURE 3. Illustration to the proof of Theorem 4

### 3. ESCAPE RATE OF BROWNIAN MOTION

We first explain the main idea of the proof. For any open set  $\Omega \subset M$ , denote by  $\tau_\Omega$  the first exit time from  $\Omega$ , that is,

$$\tau_\Omega = \inf \{t > 0 : X_t \notin \Omega\}.$$

Recall that  $B(x, r)$  denotes the geodesic ball of radius  $r$  centered at  $x$ . Fix a reference point  $z \in M$  and set  $\rho(x) = d(x, z)$ .

Let  $\{R_n\}_{n=1}^\infty$  be a sequence of strictly increasing radii to be fixed later such that  $\lim_{n \rightarrow \infty} R_n = \infty$  and consider the following sequence of stopping times

$$\tau_n = \tau_{B(z, R_n)}.$$

Then  $\tau_n - \tau_{n-1}$  is the amount of time the Brownian motion  $X_t$  takes to cross from  $\partial B(z, R_{n-1})$  to  $\partial B(z, R_n)$  for the first time (if  $n = 0$  then set  $R_0 = 0$  and  $\tau_0 = 0$ ). Let  $\{c_n\}_{n=1}^\infty$  be a sequence of positive numbers to be fixed later. Suppose that we can show that

$$(3.1) \quad \sum_{n=1}^{\infty} \mathbb{P}_z \{\tau_n - \tau_{n-1} \leq c_n\} < \infty.$$

Then, by the Borel-Cantelli lemma, with  $\mathbb{P}_z$ -probability 1 we have

$$(3.2) \quad \tau_n - \tau_{n-1} > c_n, \quad \text{for all large enough } n.$$

For any  $n \geq 1$ , set

$$T_n = \sum_{k=1}^n c_k.$$

It follows from (3.2) that, for all sufficiently large  $n$ ,

$$\tau_n > T_n - T_0,$$

where  $T_0$  is a large enough (random) number. In other words, we have the implication

$$(3.3) \quad t \leq T_n - T_0 \Rightarrow \rho(X_t) \leq R_n, \text{ if } n \text{ is large enough.}$$

Let  $\psi$  be an increasing bijection of  $\mathbb{R}_+$  onto itself such that

$$(3.4) \quad T_{n-1} - \psi(R_n) \rightarrow +\infty \text{ as } n \rightarrow \infty.$$

We claim that  $\psi^{-1}$  is an upper rate function. Indeed, for large enough  $t$ , choose  $n$  such that

$$T_{n-1} - T_0 < t \leq T_n - T_0.$$

If  $t$  is large enough then also  $n$  is large enough so that by (3.3)

$$\rho(X_t) \leq R_n$$

and by (3.4)

$$T_{n-1} - \psi(R_n) > T_0.$$

It follows that

$$t > T_{n-1} - T_0 > \psi(R_n),$$

whence

$$\rho(X_t) \leq R_n < \psi^{-1}(t),$$

which proves that  $\psi^{-1}$  is an upper rate function.

Now let us find  $c_n$  such that (3.1) is true. By the strong Markov property of Brownian motion we have

$$(3.5) \quad \mathbb{P}_z \{ \tau_n - \tau_{n-1} \leq c_n \} = \mathbb{E}_z \mathbb{P}_{X_{\tau_{n-1}}} \{ \tau_n \leq c_n \}.$$

Note that  $X_{\tau_{n-1}} \in \partial B(z, R_{n-1})$ . If a Brownian motion starts from a point  $y \in \partial B(z, R_{n-1})$ , then it has to travel no less than distance

$$r_n = R_n - R_{n-1}$$

before it reaches  $\partial B(z, R_n)$  (see FIG. 4), hence

$$\mathbb{P}_y \{ \tau_n \leq c_n \} \leq \mathbb{P}_y \{ \tau_{B(y, r_n)} \leq c_n \}, \quad y \in \partial B(z, R_{n-1}).$$




FIGURE 4. Brownian motion  $X_t$  exits the ball  $B(y, r_n)$  before  $B(z, R_n)$



From the above inequality and (3.5) we obtain

$$(3.6) \quad \mathbb{P}_z \{ \tau_n - \tau_{n-1} \leq c_n \} \leq \sup_{y \in \partial B(z, R_{n-1})} \mathbb{P}_y \{ \tau_{B(y, r_n)} \leq c_n \}.$$

For a fixed  $y \in \partial B(z, R_{n-1})$ , consider the function

$$u(x, t) = \mathbb{P}_x \{ \tau_{B(y, r)} \leq t \}.$$

Clearly,  $u(x, t)$  is the solution of the heat equation in the cylinder  $B(y, r) \times \mathbb{R}_+$ . Furthermore,  $0 \leq u \leq 1$  and

$$u(x, 0) = 0 \quad \text{for } x \in B(y, r).$$

The probability we wanted to estimate is the value of the solution at the center of the ball:

$$\mathbb{P}_y \{ \tau_{B(y, r_n)} \leq c_n \} = u(y, c_n).$$

Applying the estimate (2.2) of Theorem 4 in the cylinder  $B(y, r_n) \times [0, c_n]$  and noting that  $B(y, r_n) \subset B(z, R_n)$  so that

$$\mu(B(y, r_n)) \leq \exp(f(R_n)),$$

we obtain

$$(3.7) \quad u(y, c_n) \leq C_N \exp(f(R_n)/2) \frac{\max(\sqrt{c_n}, r_n)}{\min(\sqrt{c_n}, r_n)^{1+N/2}} \exp\left(-\frac{r_n^2}{16c_n}\right).$$

Now we choose  $c_n$  to satisfy the identity

$$\frac{r_n^2}{16c_n} = f(R_n)$$

that is,

$$c_n = \frac{1}{16} \frac{r_n^2}{f(R_n)}.$$

Noticing that  $c_n < r_n^2$  for large enough  $n$ , we obtain from (3.7)

$$\begin{aligned} \mathbb{P}_y \{ \tau_{B(y, r_n)} \leq c_n \} &\leq C_N \frac{r_n}{\sqrt{c_n}^{1+N/2}} \exp(-f(R_n)/2) \\ &= C_N r_n^{-N/2} f(R_n)^{\frac{2+N}{4}} \exp(-f(R_n)/2) \\ &\leq C_N r_n^{-N/2}. \end{aligned}$$

Set now  $R_n = 2^n$  so that  $r_n = 2^{n-1}$ . The above estimate together with (3.6) obviously implies

$$\sum_{n=1}^{\infty} \mathbb{P}_z \{ \tau_n - \tau_{n-1} \leq c_n \} \leq C_N \sum_{n=1}^{\infty} r_n^{-N/2} < \infty,$$

that is (3.1).

Knowing the sequences  $\{R_n\}$  and  $\{c_n\}$ , we can now determine a function  $\psi$  that satisfies (3.4). Indeed, we have

$$\begin{aligned} T_n &= c_1 + \dots + c_n \\ &= \frac{1}{16} \sum_{k=1}^n \frac{r_k^2}{f(R_k)} \\ &= \frac{1}{128} \sum_{k=1}^n \frac{R_{k+1}(R_{k+1} - R_k)}{f(R_k)} \\ &\geq \frac{1}{128} \sum_{k=1}^n \int_{R_k}^{R_{k+1}} \frac{r dr}{f(r)} \\ &= \frac{1}{128} \int_{R_1}^{R_{n+1}} \frac{r dr}{f(r)}. \end{aligned}$$

Setting

$$\psi(r) = c \int_0^r \frac{r dr}{f(r)},$$

where  $c < \frac{1}{128}$ , and using (1.7), we obtain that

$$T_n - \psi(R_{n+1}) \rightarrow \infty \text{ as } n \rightarrow \infty,$$

which is equivalent to (3.4). Therefore,  $\psi^{-1}$  is an upper rate function. Clearly,  $\psi^{-1}(t) = \phi(Ct)$  where  $\phi$  is defined by (1.8) and  $C = c^{-1}$ , which finishes the proof of our main result Theorem 1.

#### 4. ESCAPE RATE ON MODEL MANIFOLDS

In this section we compute sharp upper rate function on model manifolds and compare it to the one from Theorem 1. We first illustrate the method in a simple case when  $M$  is a hyperbolic space.

**4.1. Constant curvature.** Let  $M$  be the hyperbolic space  $\mathbb{H}_K^N$  of dimension  $N \geq 2$  and of the constant sectional curvature  $-K^2$ , where  $K > 0$ . Then  $\mu(B(z, r)) \leq C e^{(N-1)Kr}$  so that we can take  $f(r) = (N-1)Kr$ . Theorem 1 yields the following upper rate function

$$R(t) = CK(N-1)t.$$

In this case a sharp upper rate function can be computed as follows. The radial process  $r_t = \rho(X_t)$  satisfies the identity (see [13])

$$r_t = \sqrt{2} W_t + (N-1) \int_0^t K \coth Kr_s ds,$$

where  $W_t$  is a one-dimensional Brownian motion. We have

$$r_t \rightarrow \infty \text{ and } \frac{W_t}{t} \rightarrow 0$$

as  $t \rightarrow \infty$ , whence

$$\frac{r_t}{t} \rightarrow (N-1)K.$$

Hence, a sharp upper rate function is

$$R(t) = (1 + \varepsilon) K (N - 1) t,$$

where  $\varepsilon > 0$ .

**4.2. General model manifolds.** Here  $M$  is not necessarily Cartan-Hadamard, but we do assume that  $M$  has a pole  $z$ , that is, the exponential map  $\exp : T_z M \rightarrow M$  is a diffeomorphism. Then the polar coordinates  $(\rho, \theta)$  are defined on  $M \setminus \{z\}$ . The manifold  $M$  is said to be a *model* if the Riemannian metric of  $M$  is spherically symmetric, that is, has the form

$$(4.1) \quad ds^2 = dr^2 + h(r)^2 d\theta^2,$$

where  $h(r)$  is a smooth positive function of  $r > 0$  and  $d\theta^2$  is the canonical metric on  $\mathbb{S}^{N-1}$  (note that  $\theta$  varies in  $\mathbb{S}^{N-1}$ ). For example,  $\mathbb{R}^N$  is a model with  $h(r) = r$  and the hyperbolic space  $\mathbb{H}_K^N$  is a model with  $h(r) = K^{-1} \sinh Kr$ . The volume growth function of the metric (4.1) is

$$V(r) := \mu(B(z, r)) = \omega_N \int_0^r h(s)^{N-1} ds,$$

where  $\omega_N$  is the  $(N - 1)$ -volume of the unit sphere  $\mathbb{S}^{N-1}$ . The Laplace operator of the metric (4.1) is represented in the polar coordinates as follows:

$$\Delta = \frac{\partial^2}{\partial r^2} + m(r) \frac{\partial}{\partial r} + \frac{1}{h^2(r)} \Delta_{\mathbb{S}^{n-1}},$$

where  $\Delta_{\mathbb{S}^{n-1}}$  is the Laplacian in the variable  $\theta$  with respect to the canonical metric of  $\mathbb{S}^{N-1}$  and

$$m(r) := (N - 1) \frac{h'}{h} = \frac{V''}{V'}.$$

The function  $m(r)$  plays an important role in what follows. Clearly,  $m$  satisfies the identity

$$(4.2) \quad V'(r) = V'(r_0) \exp\left(\int_{r_0}^r m(s) ds\right)$$

for all  $r > r_0 > 0$ . We assume in the sequel that

$$(4.3) \quad m(r) > 0 \quad \text{and} \quad m'(r) \geq 0 \quad \text{for large enough } r$$

and

$$(4.4) \quad \int^\infty \frac{dr}{m(r)} = \infty.$$

For example, we have  $m(r) = \frac{N-1}{r}$  in  $\mathbb{R}^N$  and  $m(r) = (N - 1) K \coth Kr$  in  $\mathbb{H}_K^N$ . In neither case is the hypothesis (4.3) satisfied. On the other hand, if  $V'(r) = \exp(r^\alpha)$  then  $m(r) = \alpha r^{\alpha-1}$ , and both (4.3) and (4.4) are satisfied provided  $1 \leq \alpha \leq 2$ . If  $V'(r) = \exp(r^2 \log^\beta r)$  then (4.3) and (4.4) hold for all  $0 \leq \beta \leq 1$ .

We claim that under the condition (4.3) Brownian motion on  $M$  is transient, and under the conditions (4.3)-(4.4)  $M$  is stochastically complete. We use the following well-known results (cf. [8]) that for model manifolds the recurrence is equivalent to

$$(4.5) \quad \int^{\infty} \frac{dr}{V'(r)} = \infty,$$

and the stochastic completeness is equivalent to

$$(4.6) \quad \int^{\infty} \frac{V(r)}{V'(r)} dr = \infty.$$

Clearly, (4.3) implies that  $m(r) \geq c$  for some positive constant  $c$  and for all large enough  $r$ . It follows from (4.2) that  $V'(r)$  grows at least exponentially as  $r \rightarrow \infty$ , which implies (4.5). To prove (4.6), observe that, for large enough  $r > r_0$  we have

$$\begin{aligned} V(r) - V(r_0) &= \int_{r_0}^r V'(s) ds = \int_{r_0}^r \frac{V''(s)}{m(s)} ds \\ &\geq \frac{1}{m(r)} \int_{r_0}^r V''(s) ds = \frac{1}{m(r)} (V'(r) - V'(r_0)), \end{aligned}$$

whence

$$\frac{1}{m(r)} \leq \frac{V(r) - V(r_0)}{V'(r) - V'(r_0)} \sim \frac{V(r)}{V'(r)} \text{ as } r \rightarrow \infty.$$

Hence, (4.6) follows from (4.4).

Let us define the function  $r(t)$  by the identity

$$(4.7) \quad t = \int_0^{r(t)} \frac{ds}{m(s)}.$$

Our main result in this section is as follows.

**Theorem 5.** *Under the above assumptions, the function  $r((1 + \varepsilon)t)$  is the upper rate function for Brownian motion on  $M$  for any  $\varepsilon > 0$ , and is not for any  $\varepsilon < 0$ .*

Let us compare the function  $r(t)$  with the upper rate function  $R(t)$  given by Theorem 1, which is defined by the identity

$$\int_0^{R(t)} \frac{r dr}{\log V(r)} = Ct.$$

For “nice” functions  $V(r)$ , one has

$$(4.8) \quad \frac{V''}{V'} \simeq \frac{V'}{V} = (\log V)' \simeq \frac{\log V(r)}{r},$$

which means that the functions  $r(t)$  and  $R(t)$  are comparable up to a constant multiple in front of  $t$ . For example, (4.8) holds for functions like  $V(r) = \exp(r^\alpha)$  and  $V(r) = \exp(r^\alpha \log^\beta r)$ , where  $\alpha > 0$ , etc. On the other hand, it is easy to construct an example of  $V(r)$  when  $r(t)$  may be significantly less than  $R(t)$ , because one can modify a “nice” function  $V(r)$  to make the second derivative  $V''(r)$  very small in some intervals without affecting too much the values of  $V'$  and  $V$ . Then the function  $r(t)$  in (4.7) will drop significantly, while  $R(t)$  will not change very much.

*Proof of Theorem 5.* By the Ito decomposition, the radial process  $r_t = \rho(X_t)$  satisfies the identity

$$(4.9) \quad r_t = \sqrt{2}W_t + \int_0^t m(r_s) ds,$$

where  $W_t$  is a one-dimensional Brownian motion (see [13]). Since the process  $X_t$  is transient,  $r_t \rightarrow \infty$  as  $t \rightarrow \infty$  with probability 1. Hence,  $m(r_t) \geq c$  for large enough  $t$  so that the second term in the right hand side of (4.9) grows at least linearly in  $t$ . Since  $W_t = o(t)$  as  $t \rightarrow \infty$ , we have with probability 1,

$$(4.10) \quad r_t \sim \int_0^t m(r_s) ds \text{ as } t \rightarrow \infty.$$

Consider the function

$$u(t) = \int_0^t m(r_s) ds$$

It follows from (4.10) that, for any  $C > 1$  and for large enough  $t$ ,

$$(4.11) \quad r_t \leq Cu(t)$$

whence by the monotonicity of  $m$ ,

$$m(r_t) \leq m(Cu(t)).$$

Since  $\frac{du}{dt}(t) = m(r_t)$ , we obtain the differential inequality for  $u(t)$ :

$$\frac{du}{dt} \leq m(Cu(t)).$$

Solving it by separation of variables, we obtain, for large enough  $t_0$  and for all  $t > t_0$ ,

$$\int_{Cu(t_0)}^{Cu(t)} \frac{d\xi}{m(\xi)} \leq C(t - t_0),$$

whence

$$(4.12) \quad \int_0^{Cu(t)} \frac{d\xi}{m(\xi)} \leq Ct + C_0,$$

where  $C_0$  is a large enough (random) constant. Comparing (4.12) with (4.7) and using again (4.11), we obtain

$$r_t \leq Cu(t) \leq r(Ct + C_0) \leq r(C^2t)$$

for large enough  $r$  with probability 1. Since  $C > 1$  was arbitrary, this proves that  $r((1 + \varepsilon)t)$  is an upper rate function for any  $\varepsilon > 0$ . In the same way one proves that  $r_t \geq r(C^{-2}t)$  for large enough  $t$  so that  $r((1 - \varepsilon)t)$  is not an upper rate function.  $\square$

**ACKNOWLEDGEMENTS.** The research of the first author was supported by SFB 701 of the German Research Council. The research of the second author was supported in part by the NSF grant DMS-0407819. This work was done during a visit of the second author at the University of Bielefeld, which was supported by a grant of SFB 701.

## REFERENCES

- [1] **Barlow M.T., Perkins E.A.**, Symmetric Markov chains in  $\mathbb{Z}^d$ : how fast can they move?, *Probab. Th. Rel. Fields*, **82** (1989) 95–108.
- [2] **Davies E.B.**, Heat kernel bounds, conservation of probability and the Feller property, *J. d'Analyse Math.*, **58** (1992) 99-119.
- [3] **Grigor'yan A.**, On stochastically complete manifolds, (in Russian) *DAN SSSR*, **290** (1986) no.3, 534-537. Engl. transl.: *Soviet Math. Dokl.*, **34** (1987) no.2, 310-313.
- [4] **Grigor'yan A.**, The heat equation on non-compact Riemannian manifolds, (in Russian) *Matem. Sbornik*, **182** (1991) no.1, 55-87. Engl. transl.: *Math. USSR Sb.*, **72** (1992) no.1, 47-77.
- [5] **Grigor'yan A.**, Heat kernel upper bounds on a complete non-compact manifold, *Revista Matemática Iberoamericana*, **10** (1994) no.2, 395-452.
- [6] **Grigor'yan A.**, Integral maximum principle and its applications, *Proc. Roy. Soc. Edinburgh*, **124A** (1994) 353-362.
- [7] **Grigor'yan A.**, Escape rate of Brownian motion on weighted manifolds, *Applicable Analysis*, **71** (1999) no.1-4, 63-89.
- [8] **Grigor'yan A.**, Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds, *Bull. Amer. Math. Soc.*, **36** (1999) 135-249.
- [9] **Grigor'yan A., Kelbert M.**, Range of fluctuation of Brownian motion on a complete Riemannian manifold, *Ann. Probab.*, **26** (1998) 78-111.
- [10] **Grigor'yan A., Kelbert M.**, On Hardy-Littlewood inequality for Brownian motion on Riemannian manifolds, *J. London Math. Soc. (2)*, **62** (2000) 625-639.
- [11] **Hoffman D., Spruck J.**, Sobolev and isoperimetric inequalities for Riemannian submanifolds, *Comm. Pure Appl. Math.*, **27** (1974) 715–727. See also “A correction to: Sobolev and isoperimetric inequalities for Riemannian submanifolds”, *Comm. Pure Appl. Math.*, **28** (1975) no.6, 765–766.
- [12] **Hsu, Elton P.**, Heat semigroup on a complete Riemannian manifold, *Ann. Probab.*, **17** (1989) 1248-1254.
- [13] **Hsu, Elton P.**, “Stochastic Analysis on Manifolds”, Graduate Texts in Mathematics 38, 2002.
- [14] **Moser J.**, A Harnack inequality for parabolic differential equations, *Comm. Pure Appl. Math.*, **17** (1964) 101-134. Correction: *Comm. Pure Appl. Math.*, **20** (1967) 231-236.
- [15] **Takeda M.**, On a martingale method for symmetric diffusion process and its applications, *Osaka J. Math*, **26** (1989) 605-623.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BIELEFELD, 33501 BIELEFELD, GERMANY  
*E-mail address:* grigor@math.uni-bielefeld.de

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, IL 60208  
*E-mail address:* ehshu@math.northwestern.edu