

INTENSE TRIPLES

Mima Stanojkovski

Universiteit Leiden – Università degli Studi di Padova

31 July 2016

PhD project supervised by

Prof. Hendrik Lenstra and Prof. Andrea Lucchini

Some of the results are joint work with

Jon González Sánchez (EHU)

Intense automorphisms

Let G be a finite group. An automorphism α of G is **intense** if for all $H \leq G$ there exists $g \in G$ such that $\alpha(H) = gHg^{-1}$.
Write $\alpha \in \text{Int}(G)$.

Intense automorphisms

Let G be a finite group. An automorphism α of G is **intense** if for all $H \leq G$ there exists $g \in G$ such that $\alpha(H) = gHg^{-1}$. Write $\alpha \in \text{Int}(G)$.

Motivation: Intense automorphisms appear naturally as solutions to a certain cohomological problem. They (surprisingly!) give rise to a very rich theory.

Example:

- Every automorphism of a cyclic group is intense.
- Inner automorphisms are intense.

Intensity

Let p be a prime number and let G be a finite p -group. Then

$$\text{Int}(G) = P \rtimes C$$

where

- P is a p -group.
- C is cyclic of order dividing $p - 1$.

Intensity

Let p be a prime number and let G be a finite p -group. Then

$$\text{Int}(G) = P \rtimes C$$

where

- P is a p -group.
- C is cyclic of order dividing $p - 1$.

The group C is unique up to conjugation under elements of P .
The **intensity** of G is $\text{int}(G) = \#C$.

Intense triples

An **intense triple** is a triple (p, G, α) such that

- p is a prime number.
- G is a finite p -group.
- α is conjugate to a non-trivial element of C .

Intense triples

An **intense triple** is a triple (p, G, α) such that

- p is a prime number.
- G is a finite p -group.
- α is conjugate to a non-trivial element of C .

Intense triples are quite rare: if a group occurs in an intense triple, then its structure is almost uniquely determined by p and its *class*.

There are no intense triples with $p = 2$.

Equivalent triples

Example:

Let p be an odd prime and let $n \in \mathbb{Z}_{>0}$.

For all $\alpha \in \mathbb{F}_p^* \setminus \{1\}$, the triple $(p, \mathbb{F}_p^n, \alpha)$ is intense.

Equivalent triples

Example:

Let p be an odd prime and let $n \in \mathbb{Z}_{>0}$.

For all $\alpha \in \mathbb{F}_p^* \setminus \{1\}$, the triple $(p, \mathbb{F}_p^n, \alpha)$ is intense.

Two intense triples (p, G, α) and (q, G', β) are **equivalent** if there exists an isomorphism $\sigma : G \rightarrow G'$ such that $\beta = \sigma\alpha\sigma^{-1}$. It follows that $p = q$.

Let $\mathcal{T} = \{[p, G, \alpha] \mid p, G, \alpha \dots\}$ denote the set of equivalence classes of intense triples.

Abelian groups

Let p be a prime number and let \mathbb{Z}_p denote the ring of p -adic integers. Define $\omega(\mathbb{F}_p^*) = \{\alpha \in \mathbb{Z}_p^* \mid \alpha^{p-1} = 1\}$.

Note that $\omega(\mathbb{F}_p^*) \cong \mathbb{F}_p^*$ and that every abelian p -group has a natural structure of \mathbb{Z}_p -module.

Abelian groups

Let p be a prime number and let \mathbb{Z}_p denote the ring of p -adic integers. Define $\omega(\mathbb{F}_p^*) = \{\alpha \in \mathbb{Z}_p^* \mid \alpha^{p-1} = 1\}$.

Note that $\omega(\mathbb{F}_p^*) \cong \mathbb{F}_p^*$ and that every abelian p -group has a natural structure of \mathbb{Z}_p -module.

Proposition

Assume that:

- p is odd.
- $G \neq 1$ is a finite abelian p -group.
- $\alpha \in \omega(\mathbb{F}_p^*) \setminus \{1\}$. (Example: $\alpha = -1$)

Then $[p, G, \alpha] \in \mathcal{T}$ and $\text{int}(G) = p - 1$.

The lower central series

The **lower central series** of G is given by

- $G_1 = G$.
- $G_{i+1} = [G, G_i]$.

If G is a p -group, there exists k such that $G_k = 1$ and the **(nilpotency) class** of G is

$$c = \#\{i \mid G_i \neq G_{i+1}\} = -1 + \min\{k \mid G_k = 1\}.$$

Strategy

For all $c \in \mathbb{Z}_{\geq 0}$, let $\mathcal{T}[c] = \{[\rho, G, \alpha] \in \mathcal{T} \mid G \text{ has class } c\}$.

Then:

- $\mathcal{T} = \bigsqcup_c \mathcal{T}[c]$.
- $\mathcal{T}[0] = \emptyset$.
- $\mathcal{T}[1] = \{[\rho, G, \alpha] \text{ as in the Proposition}\}$.
- $\mathcal{T}[c]$ for $c = 2$?
- $\mathcal{T}[c]$ for $c \geq 3$?

Class 2

Let p be an odd prime and let $n \in \mathbb{Z}_{>0}$. Define $(\text{ES}(p, n), *)$ as

- $\text{ES}(p, n) = \mathbb{F}_p \times \mathbb{F}_p^n \times \mathbb{F}_p^n$.
- $(z_1, y_1, x_1) * (z_2, y_2, x_2) = (z_1 + z_2 + x_1 \cdot y_2, y_1 + y_2, x_1 + x_2)$.

Class 2

Let p be an odd prime and let $n \in \mathbb{Z}_{>0}$. Define $(\text{ES}(p, n), *)$ as

- $\text{ES}(p, n) = \mathbb{F}_p \times \mathbb{F}_p^n \times \mathbb{F}_p^n$.
- $(z_1, y_1, x_1) * (z_2, y_2, x_2) = (z_1 + z_2 + x_1 \cdot y_2, y_1 + y_2, x_1 + x_2)$.

Exercise:

- $(\text{ES}(p, n), *)$ has order p^{2n+1} and class 2.
- Let $\lambda \in \mathbb{F}_p^*$. Then $\alpha_\lambda : (z, y, x) \mapsto (\lambda^2 z, \lambda y, \lambda x)$ is an intense automorphism of $(\text{ES}(p, n), *)$.

Class 2

Let p be an odd prime and let $n \in \mathbb{Z}_{>0}$. Define $(\text{ES}(p, n), *)$ as

- $\text{ES}(p, n) = \mathbb{F}_p \times \mathbb{F}_p^n \times \mathbb{F}_p^n$.
- $(z_1, y_1, x_1) * (z_2, y_2, x_2) = (z_1 + z_2 + x_1 \cdot y_2, y_1 + y_2, x_1 + x_2)$.

Exercise:

- $(\text{ES}(p, n), *)$ has order p^{2n+1} and class 2.
- Let $\lambda \in \mathbb{F}_p^*$. Then $\alpha_\lambda : (z, y, x) \mapsto (\lambda^2 z, \lambda y, \lambda x)$ is an intense automorphism of $(\text{ES}(p, n), *)$.

Proposition

$\mathcal{T}[2] = \{[p, (\text{ES}(p, n), *), \alpha_\lambda] \mid p \text{ is odd}, n \in \mathbb{Z}_{>0}, \lambda \in \mathbb{F}_p^* \setminus \{1\}\}$.

Class at least 3

Given a p -group G , let $(G_i)_{i \geq 1}$ be its lower central series. Let $(f_i)_{i \geq 1}$ be the sequence, with values in $\mathbb{Z}_{\geq 0}$, such that the order of G_i/G_{i+1} is equal to p^{f_i} .

Class at least 3


Given a p -group G , let $(G_i)_{i \geq 1}$ be its lower central series. Let $(f_i)_{i \geq 1}$ be the sequence, with values in $\mathbb{Z}_{\geq 0}$, such that the order of G_i/G_{i+1} is equal to p^{f_i} .

Proposition

Let $c \geq 3$ and assume $[p, G, \alpha] \in \mathcal{T}[c]$. The following hold.

- The order of α is equal to 2 and $\text{int}(G) = 2$.
- For all i , the quotient G_i/G_{i+1} is a vector space over \mathbb{F}_p and α induces multiplication by $(-1)^i$ on it.
- $(f_i)_{i \geq 1} = (2, 1, 2, 1, \dots, 2, 1, f, 0, 0, 0, \dots)$ with $f \in \{0, 1, 2\}$.

Normal subgroups structure



An intense graph

Fix p and define $\mathcal{T}_p = \{[p, G, \alpha] \mid G, \alpha, \dots\}$.

There is a well-defined sequence of sets

$$\dots \longrightarrow \mathcal{T}_p[c+1] \xrightarrow{\pi_{c+1}} \mathcal{T}_p[c] \xrightarrow{\pi_c} \mathcal{T}_p[c-1] \longrightarrow \dots$$

where, for all c , the map π_c is defined by

$$\pi_c : [p, G, \alpha] \mapsto [p, G/G_c, \bar{\alpha}].$$

An intense graph

Fix p and define $\mathcal{T}_p = \{[p, G, \alpha] \mid G, \alpha, \dots\}$.

There is a well-defined sequence of sets

$$\dots \longrightarrow \mathcal{T}_p[c+1] \xrightarrow{\pi_{c+1}} \mathcal{T}_p[c] \xrightarrow{\pi_c} \mathcal{T}_p[c-1] \longrightarrow \dots$$


where, for all c , the map π_c is defined by

$$\pi_c : [p, G, \alpha] \mapsto [p, G/G_c, \bar{\alpha}].$$


We define a graph $\mathcal{G}_p = (E_p, V_p)$, where

- $V_p = \mathcal{T}_p$.
- $(v, w) \in E_p$ if there exists c such that $\pi_c(v) = w$.

The graph for $p = 3$



The graph for $p > 3$



The infinite case

Theorem

Let p be an odd prime and let $c \in \mathbb{Z}_{>0}$. Then the following hold.

- If $c \geq 3$, then $\mathcal{T}_p[c]$ is finite.
- $\mathcal{T}_p[c] = \emptyset \iff p = 3$ and $c \geq 5$.
- If $p > 3$, then $\# \varprojlim_c \mathcal{T}_p[c] = 1$.

The infinite case

Theorem

Let p be an odd prime and let $c \in \mathbb{Z}_{>0}$. Then the following hold.

- If $c \geq 3$, then $\mathcal{T}_p[c]$ is finite.
- $\mathcal{T}_p[c] = \emptyset \iff p = 3$ and $c \geq 5$.
- If $p > 3$, then $\# \varprojlim_c \mathcal{T}_p[c] = 1$.

If $\varprojlim_c \mathcal{T}_p[c] = \{[p, G^{(c)}, \alpha^{(c)}]\}_{c>0}$, we want to determine the pro- p -group $G_{lim} = \varprojlim_c G^{(c)}$ and the automorphism α_{lim} of G_{lim} that is induced by the automorphisms $\alpha^{(c)}$.

A profinite example

Let $p > 3$ be a prime and let $t \in \mathbb{Z}_p$ satisfy $\left(\frac{t}{p}\right) = -1$. Set $A_p = \mathbb{Z}_p + \mathbb{Z}_p i + \mathbb{Z}_p j + \mathbb{Z}_p ij$ with defining relations $i^2 = t$, $j^2 = p$, and $ji = -ij$. Then A_p is a non-commutative local ring such that $A_p/jA_p \cong \mathbb{F}_{p^2}$. The involution $\bar{\cdot} : A_p \rightarrow A_p$ is defined by


$$a = s + ti + uj + vij \mapsto \bar{a} = s - ti - uj - vij.$$

Let $G = \{a \in A_p^* \mid a\bar{a} = 1 \text{ and } a \equiv 1 \pmod{jA_p}\}$ and, for all $a \in G$, define $\alpha(a) = iai^{-1}$.

Theorem

G is a pro- p -group and α is topologically intense, i.e. for any closed subgroup H of G there exists $g \in G$ such that $\alpha(H) = gHg^{-1}$.

Moreover, $(G, \alpha) \cong (G_{\text{lim}}, \alpha_{\text{lim}})$.



THANK
YOU

