

# INTENSE TRIPLES

Mima Stanojkovski

Universiteit Leiden

DIAMANT SYMPOSIUM

27 November 2015

PhD project supervised by  
Prof. Hendrik Lenstra

# Intense automorphisms

Let  $G$  be a finite group. An automorphism  $\alpha$  of  $G$  is **intense** if for all  $H \leq G$  there exists  $g \in G$  such that  $\alpha(H) = gHg^{-1}$ .  
Write  $\alpha \in \text{Int}(G)$ .

# Intense automorphisms

Let  $G$  be a finite group. An automorphism  $\alpha$  of  $G$  is **intense** if for all  $H \leq G$  there exists  $g \in G$  such that  $\alpha(H) = gHg^{-1}$ . Write  $\alpha \in \text{Int}(G)$ .

**Motivation:** Intense automorphisms appear naturally as solutions to a certain cohomological problem. They (surprisingly!) give rise to a very rich theory.

Example:

- Every automorphism of a cyclic group is intense.
- Inner automorphisms are intense.

# Intensity

Let  $p$  be a prime number and let  $G$  be a finite  $p$ -group. Then

$$\text{Int}(G) = P \rtimes C$$

where

- $P$  is a  $p$ -group.
- $C$  is cyclic of order dividing  $p - 1$ .

# Intensity

Let  $p$  be a prime number and let  $G$  be a finite  $p$ -group. Then

$$\text{Int}(G) = P \rtimes C$$

where

- $P$  is a  $p$ -group.
- $C$  is cyclic of order dividing  $p - 1$ .

The group  $C$  is unique up to conjugation under elements of  $P$ .  
The **intensity** of  $G$  is  $\text{int}(G) = \#C$ .

# Intense triples

An **intense triple** is a triple  $(p, G, \alpha)$  such that

- $p$  is a prime number.
- $G$  is a finite  $p$ -group.
- $\alpha$  is conjugate to a non-trivial element of  $C$ .

# Intense triples

An **intense triple** is a triple  $(p, G, \alpha)$  such that

- $p$  is a prime number.
- $G$  is a finite  $p$ -group.
- $\alpha$  is conjugate to a non-trivial element of  $C$ .

Intense triples are quite rare: if a group occurs in an intense triple, then its structure is almost uniquely determined by  $p$  and its *class*.

There are no intense triples with  $p = 2$ .

# Equivalent triples

Example:

Let  $p$  be an odd prime and let  $n \in \mathbb{Z}_{>0}$ .

For all  $\alpha \in \mathbb{F}_p^* \setminus \{1\}$ , the triple  $(p, \mathbb{F}_p^n, \alpha)$  is intense.



# Equivalent triples

## Example:

Let  $p$  be an odd prime and let  $n \in \mathbb{Z}_{>0}$ .

For all  $\alpha \in \mathbb{F}_p^* \setminus \{1\}$ , the triple  $(p, \mathbb{F}_p^n, \alpha)$  is intense.

Two intense triples  $(p, G, \alpha)$  and  $(q, G', \beta)$  are **equivalent** if there exists an isomorphism  $\sigma : G \rightarrow G'$  such that  $\beta = \sigma\alpha\sigma^{-1}$ . It follows that  $p = q$ .

Let  $\mathcal{T} = \{[p, G, \alpha] \mid p, G, \alpha \dots\}$  denote the set of equivalence classes of intense triples.

# Abelian groups

Let  $p$  be a prime number and let  $\mathbb{Z}_p$  denote the ring of  $p$ -adic integers. Define  $\omega(\mathbb{F}_p^*) = \{\alpha \in \mathbb{Z}_p^* \mid \alpha^{p-1} = 1\}$ .

Note that  $\omega(\mathbb{F}_p^*) \cong \mathbb{F}_p^*$  and that every abelian  $p$ -group has a natural structure of  $\mathbb{Z}_p$ -module.

# Abelian groups

Let  $p$  be a prime number and let  $\mathbb{Z}_p$  denote the ring of  $p$ -adic integers. Define  $\omega(\mathbb{F}_p^*) = \{\alpha \in \mathbb{Z}_p^* \mid \alpha^{p-1} = 1\}$ .

Note that  $\omega(\mathbb{F}_p^*) \cong \mathbb{F}_p^*$  and that every abelian  $p$ -group has a natural structure of  $\mathbb{Z}_p$ -module.

## Proposition

Assume that:

- $p$  is odd.
- $G \neq 1$  is a finite abelian  $p$ -group.
- $\alpha \in \omega(\mathbb{F}_p^*) \setminus \{1\}$ . ( Example:  $\alpha = -1$  )

Then  $[p, G, \alpha] \in \mathcal{T}$  and  $\text{int}(G) = p - 1$ .

# The lower central series

Let  $G$  be a finite group.

- If  $x, y \in G$ , then  $[x, y] = xyx^{-1}y^{-1}$ .
- If  $H, K \leq G$ , then  $[H, K] = \langle [x, y] \mid x \in H, y \in K \rangle$ .

The **lower central series** of  $G$  is given by

- $G_1 = G$ .
- $G_{i+1} = [G, G_i]$ .

If  $G$  is a  $p$ -group, there exists  $k$  such that  $G_k = 1$  and the **(nilpotency) class** of  $G$  is

$$c = \#\{i \mid G_i \neq G_{i+1}\} = -1 + \min\{k \mid G_k = 1\}.$$

# Strategy

For all  $c \in \mathbb{Z}_{\geq 0}$ , let  $\mathcal{T}[c] = \{[\rho, G, \alpha] \in \mathcal{T} \mid G \text{ has class } c\}$ .

Then:

- $\mathcal{T} = \bigsqcup_c \mathcal{T}[c]$ .
- $\mathcal{T}[0] = \emptyset$ .
- $\mathcal{T}[1] = \{[\rho, G, \alpha] \text{ as in the Proposition}\}$ .
- $\mathcal{T}[c]$  for  $c = 2$  ?
- $\mathcal{T}[c]$  for  $c \geq 3$  ?

# Class 2

Let  $p$  be an odd prime and let  $n \in \mathbb{Z}_{>0}$ . Define  $(\text{ES}(p, n), *)$  as

- $\text{ES}(p, n) = \mathbb{F}_p \times \mathbb{F}_p^n \times \mathbb{F}_p^n$ .
- $(z_1, y_1, x_1) * (z_2, y_2, x_2) = (z_1 + z_2 + x_1 \cdot y_2, y_1 + y_2, x_1 + x_2)$ .

# Class 2

Let  $p$  be an odd prime and let  $n \in \mathbb{Z}_{>0}$ . Define  $(\text{ES}(p, n), *)$  as

- $\text{ES}(p, n) = \mathbb{F}_p \times \mathbb{F}_p^n \times \mathbb{F}_p^n$ .
- $(z_1, y_1, x_1) * (z_2, y_2, x_2) = (z_1 + z_2 + x_1 \cdot y_2, y_1 + y_2, x_1 + x_2)$ .

Exercise:

- $(\text{ES}(p, n), *)$  has order  $p^{2n+1}$  and class 2.
- Let  $\lambda \in \mathbb{F}_p^*$ . Then  $\alpha_\lambda : (z, y, x) \mapsto (\lambda^2 z, \lambda y, \lambda x)$  is an intense automorphism of  $(\text{ES}(p, n), *)$ .

# Class 2

Let  $p$  be an odd prime and let  $n \in \mathbb{Z}_{>0}$ . Define  $(\text{ES}(p, n), *)$  as

- $\text{ES}(p, n) = \mathbb{F}_p \times \mathbb{F}_p^n \times \mathbb{F}_p^n$ .
- $(z_1, y_1, x_1) * (z_2, y_2, x_2) = (z_1 + z_2 + x_1 \cdot y_2, y_1 + y_2, x_1 + x_2)$ .

Exercise:

- $(\text{ES}(p, n), *)$  has order  $p^{2n+1}$  and class 2.
- Let  $\lambda \in \mathbb{F}_p^*$ . Then  $\alpha_\lambda : (z, y, x) \mapsto (\lambda^2 z, \lambda y, \lambda x)$  is an intense automorphism of  $(\text{ES}(p, n), *)$ .

## Proposition

$\mathcal{T}[2] = \{[p, (\text{ES}(p, n), *), \alpha_\lambda] \mid p \text{ is odd}, n \in \mathbb{Z}_{>0}, \lambda \in \mathbb{F}_p^* \setminus \{1\}\}$ .



# Class at least 3

Given a  $p$ -group  $G$ , let  $(G_i)_{i \geq 1}$  be its lower central series. Let  $(f_i)_{i \geq 1}$  be the sequence, with values in  $\mathbb{Z}_{\geq 0}$ , such that the order of  $G_i/G_{i+1}$  is equal to  $p^{f_i}$ .

# Class at least 3


Given a  $p$ -group  $G$ , let  $(G_i)_{i \geq 1}$  be its lower central series. Let  $(f_i)_{i \geq 1}$  be the sequence, with values in  $\mathbb{Z}_{\geq 0}$ , such that the order of  $G_i/G_{i+1}$  is equal to  $p^{f_i}$ .

## Proposition

Let  $c \geq 3$  and assume  $[p, G, \alpha] \in \mathcal{T}[c]$ . The following hold.

- The order of  $\alpha$  is equal to 2 and  $\text{int}(G) = 2$ .
- For all  $i$ , the quotient  $G_i/G_{i+1}$  is a vector space over  $\mathbb{F}_p$  and  $\alpha$  induces multiplication by  $(-1)^i$  on it.
- $(f_i)_{i \geq 1} = (2, 1, 2, 1, \dots, 2, 1, f, 0, 0, 0, \dots)$  with  $f \in \{0, 1, 2\}$ .

# Normal subgroups structure



# An intense graph

Fix  $p$  and define  $\mathcal{T}_p = \{[p, G, \alpha] \mid G, \alpha, \dots\}$ .

There is a well-defined sequence of sets

$$\dots \longrightarrow \mathcal{T}_p[c+1] \xrightarrow{\pi_{c+1}} \mathcal{T}_p[c] \xrightarrow{\pi_c} \mathcal{T}_p[c-1] \longrightarrow \dots$$

where, for all  $c$ , the map  $\pi_c$  is defined by

$$\pi_c : [p, G, \alpha] \mapsto [p, G/G_c, \bar{\alpha}].$$

# An intense graph

Fix  $p$  and define  $\mathcal{T}_p = \{[p, G, \alpha] \mid G, \alpha, \dots\}$ .

There is a well-defined sequence of sets

$$\dots \longrightarrow \mathcal{T}_p[c+1] \xrightarrow{\pi_{c+1}} \mathcal{T}_p[c] \xrightarrow{\pi_c} \mathcal{T}_p[c-1] \longrightarrow \dots$$


where, for all  $c$ , the map  $\pi_c$  is defined by

$$\pi_c : [p, G, \alpha] \mapsto [p, G/G_c, \bar{\alpha}].$$


We define a graph  $\mathcal{G}_p = (E_p, V_p)$ , where

- $V_p = \mathcal{T}_p$ .
- $(v, w) \in E_p$  if there exists  $c$  such that  $\pi_c(v) = w$ .

# The graph for $p = 3$



# The graph for $p > 3$



# The infinite case

## Theorem

Let  $p$  be an odd prime and let  $c \in \mathbb{Z}_{>0}$ . Then the following hold.

- If  $c \geq 3$ , then  $\mathcal{T}_p[c]$  is finite.
- $\mathcal{T}_p[c] = \emptyset \iff p = 3$  and  $c \geq 5$ .
- If  $p > 3$ , then  $\# \varprojlim_c \mathcal{T}_p[c] = 1$ .



# The infinite case

## Theorem

Let  $p$  be an odd prime and let  $c \in \mathbb{Z}_{>0}$ . Then the following hold.

- If  $c \geq 3$ , then  $\mathcal{T}_p[c]$  is finite.
- $\mathcal{T}_p[c] = \emptyset \iff p = 3$  and  $c \geq 5$ .
- If  $p > 3$ , then  $\# \varprojlim_c \mathcal{T}_p[c] = 1$ .

If  $\varprojlim_c \mathcal{T}_p[c] = \{[p, G^{(c)}, \alpha^{(c)}]\}_{c>0}$ , we want to determine the pro- $p$ -group  $G_{lim} = \varprojlim_c G^{(c)}$  and the automorphism  $\alpha_{lim}$  of  $G_{lim}$  that is induced by the automorphisms  $\alpha^{(c)}$ .

# A profinite example

Let  $p > 3$  be a prime and let  $t \in \mathbb{Z}_p$  satisfy  $\left(\frac{t}{p}\right) = -1$ . Set  $A_p = \mathbb{Z}_p + \mathbb{Z}_p i + \mathbb{Z}_p j + \mathbb{Z}_p ij$  with defining relations  $i^2 = t$ ,  $j^2 = p$ , and  $ji = -ij$ . Then  $A_p$  is a non-commutative local ring such that  $A_p/jA_p \cong \mathbb{F}_{p^2}$ . The involution  $\bar{\cdot} : A_p \rightarrow A_p$  is defined by

$$a = s + ti + uj + vij \mapsto \bar{a} = s - ti - uj - vij.$$

Let  $G = \{a \in A_p^* \mid a\bar{a} = 1 \text{ and } a \equiv 1 \pmod{jA_p}\}$  and, for all  $a \in G$ , define  $\alpha(a) = iai^{-1}$ .

## Theorem

*$G$  is a pro- $p$ -group and  $\alpha$  is topologically intense, i.e. for any closed subgroup  $H$  of  $G$  there exists  $g \in G$  such that  $\alpha(H) = gHg^{-1}$ .*

*Moreover,  $(G, \alpha) \cong (G_{\text{lim}}, \alpha_{\text{lim}})$ .*

THANK  
YOU

