

THE GENERALIZED FRANCHETTA CONJECTURE FOR SOME HYPER-KÄHLER VARIETIES

LIE FU, ROBERT LATERVEER, AND CHARLES VIAL,
with an appendix joint with MINGMIN SHEN

ABSTRACT. We study the generalized Franchetta conjecture for holomorphic symplectic varieties. The conjecture predicts that the restriction of an algebraic cycle on the universal family of certain polarized hyper-Kähler varieties to a fiber is rationally equivalent to zero if and only if its cohomology class vanishes. We provide the following evidences: (1) The Beauville–Donagi family of Fano varieties of lines on cubic fourfolds; (2) The relative square, relative cube, relative Hilbert square and relative Hilbert cube of the universal families of K3 surfaces which are complete intersections in (weighted) projective spaces; (3) The relative product of the relative r_1, \dots, r_m -th Hilbert powers of the universal family of quartic K3 surfaces, where $r_1 + \dots + r_m \leq 5$; (4) The relative square and relative Hilbert square of the universal families of K3 surfaces of genera 6, 7, 8, 9, 10 and 12; (5) Relative square of the universal Fano variety of lines of the universal family of cubic fourfolds; (6) Zero-cycles and codimension 2 cycles for the Lehn–Lehn–Sorger–van Straten family of hyper-Kähler eightfolds. We also draw many consequences in the direction of the Beauville–Voisin conjecture as well as Voisin’s refinement for coisotropic subvarieties. In the appendix, we establish a new relation among tautological cycles on the square of the Fano variety of lines of a smooth cubic fourfold and provide some applications.

CONTENTS

1. Introduction	2
2. General remarks	5
3. Fano varieties of lines of cubic fourfolds	7
4. Hilbert squares of complete intersection K3 surfaces	9
5. Some more cases of Hilbert schemes of K3 surfaces	11
6. Lehn-Lehn-Sorger-van Straten hyper-Kähler eightfolds	22
Appendix A. On a new tautological relation	26
References	32

Date: October 25, 2017.

Lie Fu is supported by the Agence Nationale de la Recherche through ECOVA (ANR-15-CE40-0002), HodgeFun (ANR-16-CE40-0011), LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program *Investissements d’Avenir* (ANR-11-IDEX-0007), and *Projet Inter-Laboratoire* 2017 by Fédération de Recherche en Mathématiques Rhône-Alpes/Auvergne CNRS 3490.

1. INTRODUCTION

The original Franchetta conjecture [13] (proved in [19], see also [25] and [2]) states the following:

Theorem 1.1 ([13], [19], [25], [2]). *Let $g \geq 2$ be an integer and \mathcal{M}_g be the moduli space of smooth projective curves of genus g . Denote by $\mathcal{M}_g^\circ \subset \mathcal{M}_g$ the Zariski open subset parameterizing those curves without nontrivial automorphisms and by $C \rightarrow \mathcal{M}_g^\circ$ the universal curve. Then for any line bundle L on C and any $b \in \mathcal{M}_g^\circ$, the restriction of L to the fiber C_b is a multiple of the canonical bundle of C_b .*

In the case of the universal family of K3 surfaces $\mathcal{S} \rightarrow \mathcal{F}_g^\circ$, where \mathcal{F}_g° is the moduli space of polarized K3 surfaces of genus g without non-trivial automorphisms, O’Grady proposed in [32] the following analogue of the Franchetta conjecture. Recall that the Beauville–Voisin class ([7]) of a projective K3 surface S is the 0-cycle class \circ_S represented by any point on a rational curve of the K3 surface. It enjoys the property that the intersections of any two divisors, as well as the second (Chow-theoretic) Chern class of S , are multiples of \circ_S .

Conjecture 1.2 (O’Grady [32]). *Notation is as above. Then for any algebraic cycle $z \in \text{CH}^2(\mathcal{S})$ and any point $b \in \mathcal{F}_g^\circ$, the restriction of z to the fiber K3 surface S_b is a multiple of the Beauville–Voisin class of S_b .*

Using Mukai models, Conjecture 1.2 is verified in [33] for K3 surfaces with genus $g \leq 10$ and $g = 12, 13, 16, 18, 20$. Otherwise, Conjecture 1.2 is still wide open.

The main goal of the paper is to investigate the following higher dimensional analogue of O’Grady’s Conjecture 1.2 concerning projective hyper-Kähler varieties. Recall that a smooth projective variety is called *hyper-Kähler* or *irreducible holomorphic symplectic*, if it is simply connected and $H^{2,0}$ is generated by a nowhere degenerate holomorphic 2-form.

Conjecture 1.3 (Generalized Franchetta Conjecture, cf. [8]). *Let \mathcal{F} be the moduli space of polarized hyper-Kähler varieties of a given type, \mathcal{F}° be its open subset parametrizing hyper-Kähler varieties with trivial automorphism groups and $X^\circ \rightarrow \mathcal{F}^\circ$ be the universal family. For any $z \in \text{CH}(X^\circ)_{\mathbb{Q}}$, if its restrictions to the very general fibers are homologically trivial then its restriction to any fiber is (rationally equivalent to) zero.*

One could also be optimistic and expect that Conjecture 1.3 holds more generally for self-products of hyper-Kähler varieties, *i.e.*, for any $z \in \text{CH}(X^\circ \times_{\mathcal{F}^\circ} \cdots \times_{\mathcal{F}^\circ} X^\circ)_{\mathbb{Q}}$, if its restrictions to the very general fibers are homologically trivial then its restriction to any fiber is (rationally equivalent to) zero. We provide some evidence in Theorems 1.4, 1.5, 1.11 and 1.12 below.

Recently, Bergeron and Li [8] proved the cohomological version of the generalized Franchetta Conjecture 1.3 for relative 0-cycles when the second Betti number is sufficiently large, which is an important support in favor of the conjecture.

Let us also mention that Conjecture 1.3 is closely related to the so-called Beauville–Voisin conjecture and its refinement (see Conjecture 2.3 and 2.4). On the one hand, the proof of some of our main results actually uses some known cases of Beauville–Voisin conjecture (especially [41]); on the other hand, the Generalized Franchetta Conjecture implies the part of Beauville–Voisin conjecture involving only Chern classes and the polarization, see Proposition 2.5.

We outline the main results of the paper, which provide more evidences for the Generalized Franchetta Conjecture 1.3.

1.1. Powers and Hilbert powers of some K3 surfaces. We can establish Conjecture 1.3 for the relative squares and cubes, as well as the relative Hilbert squares and Hilbert cubes, of the universal family of K3 surfaces which are complete intersections in projective spaces.

Theorem 1.4. *Let B° be the parameter space of smooth K3 surfaces of genus $g = 3, 4$ or 5 , and let $\mathcal{S}^\circ \rightarrow B^\circ$ be the universal family. Let \mathcal{X}° be $\mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ$, $\mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ$, $\text{Hilb}_{B^\circ}^2 \mathcal{S}^\circ$, $\mathcal{S}^\circ \times_{B^\circ} \text{Hilb}_{B^\circ}^2 \mathcal{S}^\circ$ or $\text{Hilb}_{B^\circ}^3 \mathcal{S}^\circ$. For any cycle $z \in \text{CH}(\mathcal{X}^\circ)_{\mathbf{Q}}$ and any $b \in B^\circ$, the restriction of z to the fiber X_b is zero if and only if it is numerically trivial.*

The proof will be given in §4 for squares and Hilbert squares and in §5.2 for the other cases. We note that, thanks to the result of de Cataldo and Migliorini [11], the crucial cases are the self-products $\mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ$, $\mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ$. This is in line with our optimistic expectation that the generalized Franchetta conjecture should hold for self-products of hyper-Kähler varieties.

By pushing our techniques further (cf. §5.1), we can also treat some other cases of (Hilbert) powers of K3 surfaces :

Theorem 1.5. *The Generalized Franchetta Conjecture 1.3 holds for*

- (i) $\mathcal{S} \times_B \mathcal{S}$, $\text{Hilb}_B^2 \mathcal{S}$, $\mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S}$, $\mathcal{S} \times_B \text{Hilb}_B^2 \mathcal{S}$ and $\text{Hilb}_B^3 \mathcal{S}$, where $\mathcal{S} \rightarrow B$ is the universal family of smooth K3 surfaces of genus 2 (double planes).
- (ii) $\text{Hilb}_B^{r_1} \mathcal{S} \times_B \cdots \times_B \text{Hilb}_B^{r_m} \mathcal{S}$, where $\mathcal{S} \rightarrow B$ is the universal family of smooth quartic K3 surfaces and $r_1 + \cdots + r_m \leq 5$.
- (iii) The relative square and relative Hilbert square of the universal family of K3 surfaces of genera 6, 7, 8, 9, 10, 12.

The proof will be given in §5.3, where these results are just special cases of the more general but more technical Theorem 5.6. See also Remark 5.7 which explains that the ranges in Theorems 1.4 and 1.5 above are, at least most of them, already at the limit of our method.

As immediate consequences, we obtain some partial confirmation of Voisin's refinement of the Beauville–Voisin conjecture involving coisotropic subvarieties (Conjecture 2.4) :

Corollary 1.6. *Let S be a general K3 surface of genus $g \leq 10$ or 12 , and let X be the Hilbert square $X = S^{[2]}$. Let $R(X) \subset \text{CH}(X)_{\mathbf{Q}}$ denote the \mathbf{Q} -subalgebra generated by the polarization class h , the Chern classes c_i , and the Lagrangian surface $T \subset X$ constructed in [20, Proposition 4]. Then $R^*(X)$ injects into cohomology by the cycle class map.*

Corollary 1.7. *Let $S \subset \mathbf{P}^3$ be a quartic K3 surface, and let $X = S^{[5]}$, $S^{[2]} \times S^{[2]} \times S$, $S^{[2]} \times S^3$ or $S^{[2]} \times S^{[3]}$. Let $R(X) \subset \text{CH}(X)_{\mathbf{Q}}$ denote the \mathbf{Q} -subalgebra generated by the polarization class h , the Chern classes c_i , the coisotropic subvarieties E_μ of [44, 4.1 item 1)], the Lagrangian surface $T \subset S^{[2]}$ constructed in [20, Proposition 4], and the surface of bitangents $U \subset S^{[2]}$. Then $R^*(X)$ injects into cohomology by the cycle class map.*

Corollary 1.8. *Let $S \subset \mathbf{P}^4$ be a general K3 surface of degree 6, and let $X = S^{[3]}$. Let $R(X) \subset \text{CH}(X)_{\mathbf{Q}}$ denote the \mathbf{Q} -subalgebra generated by the polarization class h , the Chern classes c_i , the coisotropic subvarieties E_μ of [44, 4.1 item 1)], and the indeterminacy locus $I \subset X$ of the birational anti-symplectic involution of Beauville [3] (so $I \cong \mathbf{P}^3$). Then $R^*(X)$ injects into cohomology by the cycle class map.*

These three corollaries are proven in §5.3 and also partially extended to products of Hilbert schemes in Corollary 5.8. A similar application to a 19-dimensional family of double EPW sextics is given in §5.5.

Another consequence, whose proof as well as the background is in §5.6, concerns the Bloch conjecture for the anti-symplectic involution on Hilbert squares of quartic surfaces constructed by Beauville [3]:

Corollary 1.9. *Let $X = S^{[2]}$ be the Hilbert square of a quartic K3 surface S , and let $\iota: X \rightarrow X$ be the anti-symplectic involution of Beauville [3]. Then*

$$\begin{aligned} \iota^* = -\text{id}: \quad \text{CH}^i(X)_{(2)} &\rightarrow \text{CH}^i(X)_{(2)} \quad (i = 2, 4), \\ \iota^* = \text{id}: \quad \text{CH}^4(X)_{(j)} &\rightarrow \text{CH}^4(X)_{(j)} \quad (j = 0, 4). \end{aligned}$$

(Here, the notation $\text{CH}^*(X)_{(*)}$ refers to the Fourier decomposition of $\text{CH}(X)_{\mathbf{Q}}$ constructed by Shen–Vial [35].)

1.2. Beauville–Donagi family. For the universal family of Fano varieties of lines of cubic fourfolds, which form a complete family of projective hyper-Kähler fourfolds of $K3^{[2]}$ -type ([6]), we have the following slightly stronger result than predicted by Conjecture 1.3:

Theorem 1.10. *Let B° be the parameter space of smooth cubic fourfolds, $\mathcal{X}^\circ \rightarrow B^\circ$ be the universal family and $\mathcal{F}^\circ \rightarrow B^\circ$ be the universal family of Fano varieties of lines of the fibers of $\mathcal{X}^\circ/B^\circ$. Then for any $i \in \mathbf{N}$, any $z \in \text{CH}^i(\mathcal{F}^\circ)_{\mathbf{Q}}$ and any $b \in B^\circ$, the restriction of z to the fiber F_b is numerically trivial if and only if it is (rationally equivalent to) zero.¹*

In order to study the next case (Theorem 1.12), we also prove the following analogous result on the relative square of the universal family of Fano varieties of lines:

Theorem 1.11. *Notation is as in Theorem 1.10. Then for $z \in \text{CH}^i(\mathcal{F}^\circ \times_{B^\circ} \mathcal{F}^\circ)_{\mathbf{Q}}$ and any $b \in B^\circ$, the restriction of z to the fiber $F_b \times F_b$ is numerically trivial if and only if it is (rationally equivalent to) zero.²*

The proof of Theorem 1.10 (resp. Theorem 1.11) consists of two steps. First we show that cycles that belong to the image of the restriction map $\text{CH}^i(\mathcal{F}^\circ)_{\mathbf{Q}} \rightarrow \text{CH}^i(F_b)_{\mathbf{Q}}$ (resp. $\text{CH}^i(\mathcal{F}^\circ \times_{B^\circ} \mathcal{F}^\circ)_{\mathbf{Q}} \rightarrow \text{CH}^i(F_b \times F_b)_{\mathbf{Q}}$) are tautological in the sense of Remark 3.3 (resp. Definition 6.2). Second we show that relations among tautological cycles modulo numerical equivalence in fact hold modulo rational equivalence. More precisely, we determine completely in terms of generators and relations the rings of tautological cycles for F_b and $F_b \times F_b$. In the case of $F_b \times F_b$, all relations but one had been established in [41] and [35]. The remaining relation is established in a joint appendix with Mingmin Shen, where we also draw some consequences concerning the multiplicative properties of the Chow motive of F_b .

1.3. Lehn–Lehn–Sorger–van Straten family. Similarly to the Fano varieties of lines of cubic fourfolds, Lehn–Lehn–Sorger–van Straten (LLSvS) consider in [24] the twisted cubic curves on a cubic fourfold not containing a plane and show that the base of the maximal rationally connected (MRC) quotient of the moduli space of such curves is a hyper-Kähler eightfold. Later Addington and M. Lehn show in [1] that this hyper-Kähler eightfold is of $K3^{[4]}$ -deformation type. For the universal family of LLSvS hyper-Kähler eightfolds, we have the following result, which confirms the zero-cycle and codimension-2 cycle cases of the Generalized Franchetta Conjecture 1.3.

¹In fact, we show that the restriction of $\text{CH}^*(\mathcal{F}^\circ)_{\mathbf{Q}}$ to $\text{CH}^*(F_b)_{\mathbf{Q}}$ is the *tautological* subring, which is defined as the \mathbf{Q} -subalgebra generated by the Plücker polarization of F_b and by the Chern classes of F_b , see Remark 3.3.

²We actually show that the restriction of $\text{CH}^i(\mathcal{F}^\circ \times_{B^\circ} \mathcal{F}^\circ)_{\mathbf{Q}}$ to $\text{CH}^i(F_b \times F_b)_{\mathbf{Q}}$ is the *tautological* subring, which is defined as the \mathbf{Q} -subalgebra generated by the tautological subrings of the two factors together with the classes of the diagonal and the incidence subvariety; see Proposition 6.3.

Theorem 1.12. *Let $B^{\circ\circ}$ be the parameter space of smooth cubic fourfolds not containing a plane and let $\mathcal{Z} \rightarrow B^{\circ\circ}$ be the universal family of LLSvS hyper-Kähler eightfolds ([24]). Then*

- (i) *for any $b \in B^{\circ\circ}$ and for any $\gamma \in \text{CH}^8(\mathcal{Z})$ which is fiber-wise of degree 0, the restriction of γ to the fiber Z_b is (rationally equivalent to) zero.*
- (ii) *for any $b \in B^{\circ\circ}$ and for any $\gamma \in \text{CH}^2(\mathcal{Z})_{\mathbf{Q}}$, its restriction to the fiber Z_b is zero if and only if its cohomology class vanishes.*

As a consequence, we deduce a part of the Beauville–Voisin Conjecture 2.3 as well as the refined Conjecture 2.4 for LLSvS eightfolds :

Corollary 1.13. *Given any smooth cubic fourfold X which does not contain a plane, let Z be the LLSvS hyper-Kähler eightfold associated to X . Denote by h the polarization class. Then the classes*

$$h^8, c_2h^6, c_2^2h^4, c_2^3h^2, c_2^4, c_4h^4, c_2c_4h^2, c_2^2c_4, c_6h^2, c_2c_6, c_4^2, c_8 \in \text{CH}_0(\mathcal{Z})_{\mathbf{Q}}$$

are all proportional, where $c_i := c_i(T_Z)$ is the i -th (Chow-theoretic) Chern class of the tangent bundle of Z . We call the generator of degree 1 in this one-dimensional subspace the canonical 0-cycle class or the Beauville–Voisin class of Z , denoted by ν_Z .

More strongly, let $R(\mathcal{Z})$ be the \mathbf{Q} -subalgebra generated by the polarization class h , the Chern classes c_i together with the following classes of coisotropic subvarieties of Z :

- *the embedded cubic fourfold $X \subset Z$ ([24]) ;*
- *the space of twisted cubics contained in a general hyperplane section of X ([37]) ;*
- *the coisotropic subvarieties of codimension 1, 2, 3, 4 constructed by Voisin ([44, Corollary 4.9]) ;*
- *the fixed locus of the anti-symplectic involution ι of Z ([22]) ;*
- *the images by ι of all the above subvarieties.*

Then $R^8(\mathcal{Z}) = \mathbf{Q} \cdot \nu_Z$.

Conventions. All algebraic varieties are over the field of complex numbers. We work with Chow groups with rational coefficients. For the m -th Hilbert scheme of a surface S , the two notations $S^{[m]}$ and $\text{Hilb}^m(S)$ are used interchangeably and similarly for the relative situation.

Acknowledgements. The authors want to thank Nicolas Addington, Zhiyuan Li, Renjie Lyu, Nicolas Ressayre, Qizheng Yin for their interest and helpful comments and discussions.

2. GENERAL REMARKS

2.1. Generic fiber vs. geometric fibers. There is the following slightly different version of the generalized Franchetta conjecture for hyper-Kähler varieties :

Conjecture 2.1. *Let \mathcal{F} be the moduli space of polarized hyper-Kähler varieties and let $\pi : \mathcal{X} \rightarrow \mathcal{F}$ be the universal family. Denote by \mathcal{X}_η the generic fiber of π , where $\eta = \text{Spec}(\mathbf{C}(\mathcal{F}))$. Then the group $\text{CH}^*(\mathcal{X}_\eta)_{\text{hom}}$ is torsion.*

Here homological equivalence is with respect to some classical Weil cohomology, for instance, étale cohomology or de Rham cohomology.

Lemma 2.2. *Conjecture 1.3 and Conjecture 2.1 are equivalent.*

Proof. Let us start by assuming Conjecture 1.3. Since $\text{CH}(\mathcal{X}_\eta) = \varinjlim \text{CH}(\mathcal{X}_U)$, where U runs through all non-empty Zariski open subset of the moduli space \mathcal{F} , for a given cycle $z_\eta \in \text{CH}(\mathcal{X}_\eta)$,

we can assume it comes from the universal family over a Zariski open subset \mathcal{F}° of \mathcal{F} , namely, $z \in \text{CH}(\mathcal{X}^\circ)$. Using [38, Lemma 2.1], the hypothesis that the restriction of z to the geometric generic fiber is homologically trivial implies that the restriction of z to every very general geometric fiber is also trivial. Now the conclusion of Conjecture 1.3 says that the restriction of z to a very general geometric fiber is (rationally equivalent to) zero. By the standard argument of decomposition of the diagonal ([9], [40], [43]), this implies the existence of a Zariski open dense subset $U \subset \mathcal{F}$, such that $z|_{\mathcal{X}_U}$ is zero. In particular, z_η is rationally equivalent to zero.

For the other direction, since we know that $\text{CH}^*(\mathcal{X}_\eta)_{\text{hom}}$ is torsion, by restriction we can show Conjecture 1.3 for general fibers. Then a standard specialization argument allows us to conclude for all fibers. \square

Thanks to Lemma 2.2, we will focus in this paper on Conjecture 1.3.

2.2. Relation to Beauville–Voisin conjecture. As is mentioned in the introduction, the Generalized Franchetta Conjecture 1.3 is very much related to the following Beauville–Voisin conjecture:

Conjecture 2.3 (Beauville–Voisin [5], [41]). *Let X be a projective hyper-Kähler variety. Let the Beauville–Voisin subring $\langle c_i(X), \text{Pic}(X) \rangle$ be the \mathbf{Q} -subalgebra of $\text{CH}^*(X)$ generated by line bundles and all (Chow theoretic) Chern classes of T_X . Then the restriction of the cycle class map to the Beauville–Voisin subring is injective. In other words, any polynomial of line bundles and Chern classes of X is homologically equivalent to zero if and only if it is rationally equivalent to zero.*

The original version due to Beauville in [5], under the name of *weak splitting property*, contains only line bundles; the Chern classes of the tangent bundle are introduced by Voisin in [41]. Some active progress towards this conjecture has recently been made: see [5], [41], [15], [45], [34], [17, Theorem 1.14] for the known results and more details. More recently, Voisin [44] proposes the following stronger version of Conjecture 2.3 involving the coisotropic subvarieties (in particular lagrangian subvarieties):

Conjecture 2.4 (Voisin’s refinement [44]). *Let X be a projective hyper-Kähler variety. Then the restriction of the cycle class map to the \mathbf{Q} -subalgebra of $\text{CH}(X)$ generated by line bundles, Chern classes of T_X and coisotropic subvarieties, is injective.*

We would like to point out that the generalized Franchetta conjecture implies the part of the Beauville–Voisin conjecture involving only the Chern classes of the tangent bundle and the polarization class. More generally it actually implies part of the refined Conjecture 2.4 once taking into account coisotropic subvarieties which are defined universally over the moduli space (see Corollaries 1.6, 1.7, 1.8, 5.9 and 1.13 for examples):

Proposition 2.5. *Let \mathcal{F} be a moduli space of polarized hyper-Kähler varieties. If Conjecture 1.3 holds true for the universal family over \mathcal{F} , then for any member X of this family, the cycle class map restricted to the \mathbf{Q} -subalgebra generated by the polarization line bundle and the Chern classes of X , is injective.*

More generally, still assuming Conjecture 1.3, for any member X of this family, the cycle class map restricted to the \mathbf{Q} -subalgebra generated by the algebraic cycles of X that exist universally over the moduli space, is injective.

Proof. Let $\mathcal{X}^\circ \rightarrow \mathcal{F}^\circ$ be the universal family, where $\mathcal{F}^\circ \subset \mathcal{F}$ is an open subset. For any member X and any given polynomial of polarization line bundle and Chern classes of the tangent bundle $z := P(h, c_i(T_X)) \in \text{CH}(X)$ such that the cohomology class of z vanishes, we want to show that $z = 0$. Consider $\gamma := P(h, c_i(T_{\mathcal{X}^\circ/\mathcal{F}^\circ})) \in \text{CH}(\mathcal{X}^\circ)$. If X belongs to \mathcal{F}° , then clearly $\gamma|_X = z$ and hence γ has fiber-wise vanishing cohomology class. Then the generalized Franchetta conjecture 1.3 says exactly

that z is rationally equivalent to zero. If X does not belong to \mathcal{F}° , the specialization argument for algebraic cycles allows us to conclude. The last assertion is more or less tautological. \square

2.3. Moduli space *vs.* parameter space.

Remark 2.6. To show the Generalized Franchetta Conjecture 1.3 in some cases, it will be convenient to work over some parameter space which dominates the moduli space, instead of the moduli space itself. More precisely, keep the same notation as in Conjecture 1.3 and let $B \rightarrow U$ be a surjective morphism to some Zariski dense open subset U of the moduli space \mathcal{F}° . Denote by $\mathcal{Y} \rightarrow B$ the pulled-back family of the universal family $\mathcal{X}^\circ \rightarrow \mathcal{F}^\circ$. Then the generalized Franchetta conjecture for $\mathcal{Y} \rightarrow B$ implies the generalized Franchetta conjecture for $\mathcal{X}^\circ \rightarrow \mathcal{F}^\circ$.

$$\begin{array}{ccccc} \mathcal{Y} & \longrightarrow & \mathcal{X}_U & \hookrightarrow & \mathcal{X}^\circ \\ \downarrow & & \downarrow & & \downarrow \\ B & \longrightarrow & U & \hookrightarrow & \mathcal{F}^\circ \end{array}$$

Indeed, for any $z \in \text{CH}(\mathcal{X}^\circ)$, denote by $z' \in \text{CH}(\mathcal{Y})$ its pull-back image in \mathcal{Y} . Obviously, the hypothesis that the restriction of z to a very general fiber of $\mathcal{X}^\circ/\mathcal{F}^\circ$ is homologically trivial implies the same thing for the restriction of z' to the fibers of \mathcal{Y}/B . The generalized Franchetta conjecture for \mathcal{Y}/B then implies that z' restricts to zero on each fiber of \mathcal{Y}/B . Hence so is z for each fiber of $\mathcal{X}_U \rightarrow U$. A specialization argument shows that the same thing holds for each fiber of $\mathcal{X}^\circ \rightarrow \mathcal{F}^\circ$.

3. FANO VARIETIES OF LINES OF CUBIC FOURFOLDS

In this section, we prove Theorem 1.10, which by Remark 2.6 confirms the Generalized Franchetta Conjecture 1.3 for the 20-dimensional complete family of hyper-Kähler fourfolds constructed by Beauville–Donagi in [6]. The key idea of the proof is as in [42] and [33]: the universal family has very simple Chow groups.

We start by setting up some notations. Let V be a 6-dimensional vector space and $\mathbf{P}^5 = \mathbf{P}(V)$ be its projectivization. The parameter space of possibly singular cubic fourfolds is given by the following projective space:

$$B := \mathbf{P}(H^0(\mathbf{P}^5, \mathcal{O}(3))) = \mathbf{P}(\text{Sym}^3 V^\vee) \simeq \mathbf{P}^{55}.$$

Let $B^\circ \subset B$ be the open subset parameterizing smooth cubic fourfolds. We thus have the universal family $\mathcal{X} \rightarrow B$ as well as the smooth family $\mathcal{X}^\circ \rightarrow B^\circ$ by base-change.

Let $G := \text{Gr}(\mathbf{P}^1, \mathbf{P}^5) \simeq \text{Gr}(2, 6)$ be the Grassmannian variety parameterizing all projective lines in \mathbf{P}^5 . Denote by S (*resp.* Q) the tautological rank-2 subbundle (*resp.* rank-4 quotient bundle), fitting into the following short exact sequences of vector bundles over G :

$$0 \rightarrow S \rightarrow \mathcal{O}_G \otimes V \rightarrow Q \rightarrow 0.$$

Note that for any equation $f \in \text{Sym}^3 V^\vee$, the above short exact sequence gives a section s_f of the vector bundle $\text{Sym}^3 S^\vee$, whose zero locus ($s_f = 0$) is exactly the Fano variety of lines of the cubic fourfold defined by f .

Consider the incidence subvariety \mathcal{F} in $B \times G$ defined by

$$\mathcal{F} := \{([f], l) \in B \times G \mid f|_l = 0\},$$

together with the two natural projections :

$$\begin{array}{ccc} & \mathcal{F} & \\ \pi \swarrow & & \searrow p \\ B & & G \end{array}$$

It is easy to see that $\pi : \mathcal{F} \rightarrow B$ is the universal Fano variety of lines of fibers of \mathcal{X}/B and that $p : \mathcal{F} \rightarrow G$ is a projective bundle whose fiber over a line $l \in G$ parametrizes all (possibly singular) cubic fourfolds containing l .

As in [33, Lemma 2.1], we have the following :

Lemma 3.1. *For any $b \in B$, the following two images of restriction maps are the same :*

$$\mathrm{Im}(\mathrm{CH}(\mathcal{F}) \rightarrow \mathrm{CH}(F_b)) = \mathrm{Im}(\mathrm{CH}(G) \rightarrow \mathrm{CH}(F_b)).$$

Proof. The inclusion “ \supseteq ” is trivial (we have the factorization $F_b \hookrightarrow \mathcal{F} \rightarrow G$).

Let us show the inverse inclusion. Given any cycle $z \in \mathrm{CH}(\mathcal{F})$, by the projective bundle formula,

$$z = \sum_{k \geq 0} p^*(z_k) \cdot \xi^k,$$

where $z_k \in \mathrm{CH}(G)$ and $\xi = c_1(\mathcal{O}_p(1))$. As in [33, Lemma 2.1], we easily check that ξ is a linear combination of cycles pulled back from B by π and cycles pulled back from G by p . Hence z is a polynomial of cycles of the form $p^*(\alpha)$ and $\pi^*(\beta)$. The latter type being zero when restricted to any fiber F_b , the restriction of z to F_b is therefore the restriction of some cycle of G . \square

Lemma 3.2. *For any $b \in B^\circ$,*

$$\mathrm{Im}(\mathrm{CH}(G) \rightarrow \mathrm{CH}(F_b)) \subseteq \langle c_i(F_b), \mathrm{Pic}(F_b) \rangle,$$

where the right hand side is the Beauville–Voisin subring of $\mathrm{CH}(F_b)$ generated (as a \mathbf{Q} -algebra) by line bundles and all Chern classes of the tangent bundle of F_b .

Proof. Since $\mathrm{CH}(G)$ is generated (as a \mathbf{Q} -algebra) by $c_1(S^\vee)$ and $c_2(S^\vee)$, it suffices to show that both of their restrictions to F_b lie in the Beauville–Voisin ring. The first one being a line bundle, it remains to show $c_2(S^\vee|_{F_b}) \in \langle c_i(F_b), \mathrm{Pic}(F_b) \rangle$. However, using the short exact sequence

$$0 \rightarrow T_{F_b} \rightarrow T_G|_{F_b} \rightarrow \mathrm{Sym}^3 S^\vee|_{F_b} \rightarrow 0$$

together with $T_G \simeq S^\vee \otimes Q$, one finds that

$$ch(T_{F_b}) = ch(S^\vee|_{F_b}) (6 - ch(S|_{F_b})) - ch(\mathrm{Sym}^3 S^\vee|_{F_b}),$$

and hence $c_2(T_{F_b}) = -ch_2(T_{F_b}) = 5c_1(S^\vee|_{F_b})^2 - 8c_2(S^\vee|_{F_b})$. Therefore, $c_2(S^\vee|_{F_b})$ also belongs to the Beauville–Voisin ring.³ \square

We can now easily conclude :

Proof of Theorem 1.10. For any $z \in \mathrm{CH}(\mathcal{F}^\circ)$, by the surjectivity of $\mathrm{CH}(\mathcal{F}) \rightarrow \mathrm{CH}(\mathcal{F}^\circ)$, we can actually assume that $z \in \mathrm{CH}(\mathcal{F})$. For any $b \in B^\circ$, thanks to Lemma 3.1, $z|_{F_b}$ is the restriction of some cycle from G , which must lie in the Beauville–Voisin ring $\langle c_i(F_b), \mathrm{Pic}(F_b) \rangle$ by Lemma 3.2. Now the equivalence between homological triviality and rational triviality of $z|_{F_b}$ is a consequence of Voisin’s result [41, Theorem 1.4(ii)] saying that the cycle class map restricted to the Beauville–Voisin

³The classes $c_1(S^\vee|_{F_b})$ and $c_2(S^\vee|_{F_b})$ are the classes that Claire Voisin calls g and c respectively in [41].

ring is injective. Finally, numerical equivalence and homological equivalence coincide for Fano varieties of lines of cubic fourfolds by [10]. \square

Remark 3.3. In fact, the above proof shows that the restriction of a cycle $z \in \text{CH}(\mathcal{F}^\circ)$ to a fiber Fano variety of lines F is in the so-called *tautological ring* $R^*(F)$, which is the \mathbf{Q} -subalgebra of $\text{CH}^*(F)$, in general smaller than the Beauville–Voisin ring, generated by the Plücker polarization class g and the Chern classes of F . In particular,

- $R^1(F) = \mathbf{Q} \cdot g$;
- $R^2(F) = \mathbf{Q} \cdot g^2 \oplus \mathbf{Q} \cdot c_2$;
- $R^3(F) = \mathbf{Q} \cdot g^3$ (by [41, Lemma 3.5] gc_2 and g^3 are proportional);
- $R^4(F) = \mathbf{Q} \cdot \nu_F$, where ν_F is the canonical 0-cycle class and c_2^2, c_4, g^4, g^2c_2 are all proportional to it by [41, Lemma 3.2].

4. HILBERT SQUARES OF COMPLETE INTERSECTION K3 SURFACES

In this section, we prove Theorem 1.4 for squares and Hilbert squares. There are three families of complete intersection K3 surfaces, namely, quartic surfaces in \mathbf{P}^3 , complete intersections of quadric and cubic hypersurfaces in \mathbf{P}^4 and complete intersections of three quadric hypersurfaces in \mathbf{P}^5 .

Let us fix some notations: in each of the three cases

- $\mathbf{P} := \mathbf{P}^3, \mathbf{P}^4$ resp. \mathbf{P}^5 is the ambient projective space;
- $E := \mathcal{O}_{\mathbf{P}}(4), \mathcal{O}_{\mathbf{P}}(2) \oplus \mathcal{O}_{\mathbf{P}}(3)$, resp. $\mathcal{O}_{\mathbf{P}}(2)^{\oplus 3}$ is the relevant vector bundle;
- $B := \mathbf{P}H^0(\mathbf{P}, E)$ is the parameter (projective) space and B° is the open subset parameterizing smooth K3 surfaces.
- $\mathcal{S} := \{(x, [s]) \in \mathbf{P} \times B \mid s(x) = 0\}$ is the universal family.

We have therefore the natural projections, where p is clearly a projective bundle;

$$(1) \quad \begin{array}{ccc} \mathcal{S} & \xrightarrow{p} & \mathbf{P} \\ \pi \downarrow & & \\ B & & \end{array}$$

Similarly, the relative square and the open complement of the relative diagonal in it fit into the following diagram

$$(2) \quad \begin{array}{ccc} \mathcal{S} \times_B \mathcal{S} \setminus \Delta_{\mathcal{S}/B} & \xrightarrow{q'} & \mathbf{P} \times \mathbf{P} \setminus \Delta_{\mathbf{P}} \\ \downarrow j & & \downarrow \\ \mathcal{S} \times_B \mathcal{S} & \xrightarrow{q := (p,p)} & \mathbf{P} \times \mathbf{P} \\ \downarrow \pi_2 := (\pi, \pi) & & \\ B & & \end{array}$$

Note that although q itself is not a projective bundle, its restriction q' is. Let ξ be the first Chern class of $\mathcal{O}_{q'}(1)$. The relative diagonal $\Delta_{\mathcal{S}/B}$ being of codimension 2, ξ extends uniquely to the whole $\mathcal{S} \times_B \mathcal{S}$, which we still denote by ξ by abuse of notation.

We can show the analogue of Lemma 3.1 in our case⁴ :

Proposition 4.1. *For any $b \in B^\circ$, we have :*

$$\mathrm{Im}(\mathrm{CH}(\mathcal{S} \times_B \mathcal{S}) \rightarrow \mathrm{CH}(S_b \times S_b)) = \mathrm{Im}(\mathrm{CH}(\mathbf{P} \times \mathbf{P}) \rightarrow \mathrm{CH}(S_b \times S_b)) + \Delta_* \mathrm{Im}(\mathrm{CH}(\mathbf{P}) \rightarrow \mathrm{CH}(S_b)),$$

where $\Delta : S_b \hookrightarrow S_b \times S_b$ is the diagonal embedding.

Proof. Notation is as in Diagrams (1) and (2). By base-change, it is easy to see that the right-hand side is contained in the left-hand side. Concerning the inverse inclusion, the projective bundle formula gives, for any $z \in \mathrm{CH}(\mathcal{S} \times_B \mathcal{S})$,

$$j^*(z) = \sum_{k \geq 0} q'^*(z_k) \cdot \xi^k,$$

for some cycles $z_k \in \mathrm{CH}(\mathbf{P} \times \mathbf{P} \setminus \Delta_{\mathbf{P}})$. As in Lemma 3.1, it is easy to see that $\xi = j^* \pi_2^*(h) + q'^*(\alpha)$, where $h = c_1(\mathcal{O}_B(1))$ and $\alpha \in \mathrm{CH}(\mathbf{P} \times \mathbf{P} \setminus \Delta_{\mathbf{P}})$. For each k , we denote still by $z_k \in \mathrm{CH}(\mathbf{P} \times \mathbf{P})$ its closure and similarly for α . Therefore, we have

$$z - \sum_k q^*(z_k) \cdot (\pi_2^*(h) - q^*(\alpha))^k \in \mathrm{Ker}(j^*).$$

By the localization sequence, there exists $\gamma \in \mathrm{CH}(\mathcal{S})$, such that

$$(3) \quad z - \sum_k q^*(z_k) \cdot (\pi_2^*(h) - q^*(\alpha))^k = \Delta_*(\gamma),$$

where $\Delta : \mathcal{S} \hookrightarrow \mathcal{S} \times_B \mathcal{S}$ is the diagonal embedding.

Since $p : \mathcal{S} \rightarrow \mathbf{P}$ is also a projective bundle with $c_1(\mathcal{O}_p(1)) = \pi^*(h)$, we have

$$\gamma = \sum_l p^*(\gamma_l) \cdot \pi^*(h)^l,$$

for some $\gamma_l \in \mathrm{CH}(\mathbf{P})$. Substituting this into (3), we get

$$(4) \quad z = \sum_k q^*(z_k) \cdot (\pi_2^*(h) - q^*(\alpha))^k + \sum_l \Delta_*(p^*(\gamma_l) \cdot \pi^*(h)^l),$$

Now for any $b \in B^\circ$, the restriction $z|_{S_b \times S_b}$ is of the desired form simply because the restrictions of $\pi_2^*(h)$ and $p^*(h)$ to the fibers vanish. \square

We can now prove the first two parts of Theorem 1.4.

Proof of Theorem 1.4 for relative squares. Keep the same notations as before. Thanks to Proposition 4.1, we only need to show that for any smooth complete intersection K3 surface $S \subset \mathbf{P}$, the cycle class map restricted to $\mathrm{Im}(\mathrm{CH}(\mathbf{P} \times \mathbf{P}) \rightarrow \mathrm{CH}(S \times S)) + \Delta_* \mathrm{Im}(\mathrm{CH}(\mathbf{P}) \rightarrow \mathrm{CH}(S))$ is injective. Denote $H := c_1(\mathcal{O}_{\mathbf{P}}(1))$ and $h := H|_S$. Since $\mathrm{CH}(\mathbf{P} \times \mathbf{P})$ is generated by $\mathrm{pr}_1^*(H)$ and $\mathrm{pr}_2^*(H)$, and $\Delta_*(h) = h \times \mathfrak{o}_S + \mathfrak{o}_S \times h$ (see [7]), it is enough to show that the cycle class map of $S \times S$ restricted to the subalgebra generated by $\mathrm{pr}_1^*(h)$, $\mathrm{pr}_2^*(h)$ and Δ is injective. It is the easiest case of Voisin's [41, Proposition 2.2]. \square

⁴Proposition 4.1 will be generalized for the so-called stratified projective bundle in §5.1.

Proof of Theorem 1.4 for relative Hilbert squares. Consider the blow-up of $\mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ$ along the relative diagonal $\Delta_{\mathcal{S}^\circ/B^\circ}$, the natural involution switching two factors lifts to the blow-up. It is well-known that the Hilbert square is the quotient of this lifted involution and

$$\mathrm{CH}^*(\mathrm{Hilb}_{B^\circ}^2(\mathcal{S}^\circ)) \simeq \mathrm{CH}^*(\mathrm{Bl}_\Delta(\mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ))^{inv} \simeq \mathrm{CH}^*(\mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ)^{inv} \oplus \mathrm{CH}^{*-1}(\mathcal{S}^\circ),$$

where all isomorphisms are compatible with the restriction to the fibers. Therefore for any $b \in B^\circ$ and any $z \in \mathrm{CH}^*(\mathrm{Hilb}_{B^\circ}^2(\mathcal{S}^\circ))$, its restriction to the fiber $z|_{\mathrm{Hilb}^2 S_b}$, viewed as an element in $\mathrm{CH}^*(S_b \times S_b)^{inv} \oplus \mathrm{CH}^{*-1}(S_b)$, lives in $\mathrm{Im}(\mathrm{CH}(\mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ)^{inv} \rightarrow \mathrm{CH}^*(S_b \times S_b)^{inv}) \oplus \mathrm{Im}(\mathrm{CH}^{*-1}(\mathcal{S}^\circ) \rightarrow \mathrm{CH}^{*-1}(S_b))$. We can thus conclude thanks to the established cases of the generalized Franchetta conjecture for the relative squares $\mathcal{S}^\circ \times_{B^\circ} \mathcal{S}^\circ$ and for \mathcal{S}° . \square

5. SOME MORE CASES OF HILBERT SCHEMES OF K3 SURFACES

In this section, we try to push the results of §4 to higher (Hilbert) powers and to K3 surfaces of higher genera. Let us first provide the technical tool.

5.1. Stratified projective bundle. As one can observe the similarity between Lemma 3.1 and Proposition 4.1 (also Proposition 6.1 later), the goal of this technical subsection is to summarize these situations.

Definition 5.1 (Stratified projective bundle). A projective morphism $q : \mathcal{X} \rightarrow Y$ is called a *stratified projective bundle* if there exists a commutative cartesian diagram

$$(5) \quad \begin{array}{ccccccc} \mathcal{X}_r & \hookrightarrow & \dots & \hookrightarrow & \mathcal{X}_1 & \hookrightarrow & \mathcal{X}_0 = \mathcal{X} \\ \downarrow q_r & \square & & \square & \downarrow q_1 & \square & \downarrow q_0 = q \\ Y_r & \hookrightarrow & \dots & \hookrightarrow & Y_1 & \hookrightarrow & Y_0 = Y \end{array}$$

where all horizontal morphisms are closed immersions, such that for any $0 \leq i \leq r$, the restriction of q_i

$$q'_i : \mathcal{X}_i \setminus \mathcal{X}_{i+1} \rightarrow Y_i \setminus Y_{i+1}$$

is a projective bundle ($\mathcal{X}_{r+1} = Y_{r+1} = \emptyset$). The above diagram is called a *stratification* of q .

Now we can state the following generalization of Lemma 3.1 and Proposition 4.1 (see also Proposition 6.1 for an example).

Proposition 5.2. *Let $q : \mathcal{X} \rightarrow Y$ be a stratified projective bundle with a given stratification (5) and $\pi : \mathcal{X} \rightarrow B$ be a surjective morphism. Assume moreover that for any $0 \leq i \leq r$, Y_i is smooth projective, \mathcal{X}_i is flat over a (common) Zariski open subset $B^\circ \subset B$, $\mathrm{codim}_{\mathcal{X}_i}(\mathcal{X}_{i+1}) \geq 2$ and finally there exists a line bundle on B whose restriction to fibers of the projective bundle q'_i is non-trivial. Then for any $b \in B^\circ$*

$$\mathrm{Im}(\mathrm{CH}(\mathcal{X}) \rightarrow \mathrm{CH}(X_b)) = \sum_{i=0}^r \iota_{i*} \mathrm{Im}(q_{i,b}^* : \mathrm{CH}(Y_i) \rightarrow \mathrm{CH}(X_{i,b})),$$

where X_b (resp. $X_{i,b}$) is the fiber of \mathcal{X} (resp. the Zariski closure of $\mathcal{X}_i \setminus \mathcal{X}_{i+1}$) over b , $\iota_i : X_{i,b} \hookrightarrow X_b$ is the natural inclusion and $q_{i,b}$ is the restriction of q_i to $X_{i,b}$.

Proof. Since the \mathcal{X}_i 's are flat over B° , by base-change, the right-hand side is clearly contained in the left-hand side. We use induction on r to prove the other inclusion. For any $z \in \text{CH}(\mathcal{X})$, the projective bundle formula shows that

$$j^*(z) = \sum_{k \geq 0} q_0'^*(z_k) \cdot \xi^k,$$

for some cycles $z_k \in \text{CH}(Y_0 \setminus Y_1)$ where $j : \mathcal{X} \setminus \mathcal{X}_1 \hookrightarrow \mathcal{X}$ is the open immersion and $\xi = c_1(\mathcal{O}_{q_0'}(1))$. By hypothesis, $\xi = j^* \pi^*(h) + q_0'^*(\alpha)$, where h is a divisor on B and $\alpha \in \text{CH}(Y_0 \setminus Y_1)$. We extend z_k and α to Y_0 , keeping the same notation for the classes on Y_0 . Therefore

$$z - \sum_k q^*(z_k) \cdot (\pi^*(h) - q^*(\alpha))^k \in \text{Ker}(j^*).$$

By the localization sequence, there exists $\gamma \in \text{CH}(\mathcal{X}_1)$, such that

$$(6) \quad z = \sum_k q^*(z_k) \cdot (\pi^*(h) - q^*(\alpha))^k + \iota_*(\gamma),$$

where $\iota : \mathcal{X}_1 \hookrightarrow \mathcal{X}$ is the natural inclusion.

Noting that the restriction of $\pi^*(h)$ to X_b vanishes, we have that

$$z|_{X_b} \in \text{Im}(q^* : \text{CH}(Y) \rightarrow \text{CH}(X_b)) + \text{Im}(\iota_* : \text{CH}(\mathcal{X}_1 \rightarrow \mathcal{X}))|_{X_b},$$

where the second term is $\iota_{1,*} \text{Im}(\text{CH}(\mathcal{X}_1) \rightarrow \text{CH}(X_{1,b}))$ by flat base-change. Observing that $q_1 : \mathcal{X}_1 \rightarrow Y_1$ is again a stratified projective bundle verifying all the conditions, the induction hypothesis allows us to conclude. \square

5.2. Cubes and Hilbert cubes of complete intersection K3 surfaces. We prove Theorem 1.4 for cubes and Hilbert cubes in this subsection. Notation is as in §4.

The geometry is quite close⁵ to the one considered in [14], in particular, we will study collinear triples in the projective space \mathbf{P} . For three points in \mathbf{P} there are four types of relative positions: non-collinear, collinear and distinct, two coincide but not with the third, all coincide. As a result, the evaluation map of the relative cube of the universal family

$$q : \mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S} \rightarrow \mathbf{P} \times \mathbf{P} \times \mathbf{P}$$

is not a projective bundle but is a *stratified* projective bundle (Definition 5.1) with the following stratification:

$$(7) \quad \begin{array}{ccccccc} \mathcal{S} = \delta_{\mathcal{S}/B} \hookrightarrow \Delta_{12} \cup \Delta_{13} \cup \Delta_{23} \hookrightarrow \Delta_{12} \cup \Delta_{13} \cup \Delta_{23} \cup \mathcal{I} \hookrightarrow \mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S} \xrightarrow{\pi_3} B \\ \downarrow p \quad \square \quad \downarrow \quad \square \quad \downarrow \quad \square \quad \downarrow q \\ \mathbf{P} = \delta_{\mathbf{P}} \hookrightarrow \Delta_{12} \cup \Delta_{13} \cup \Delta_{23} \hookrightarrow J \hookrightarrow \mathbf{P} \times \mathbf{P} \times \mathbf{P} \end{array}$$

where in the first row, $\Delta_{i,j} : \mathcal{S} \times_B \mathcal{S} \hookrightarrow \mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S}$ are three big (relative) diagonals, \mathcal{I} is the Zariski closure of

$$\mathcal{I}^\circ := \{(x, y, z) \in \mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S} \mid x, y, z \text{ collinear and distinct}\};$$

in the second row, $\Delta_{i,j} : \mathbf{P} \times \mathbf{P} \hookrightarrow \mathbf{P} \times \mathbf{P} \times \mathbf{P}$ are three big diagonals and

$$J := \{(x, y, z) \in \mathbf{P} \times \mathbf{P} \times \mathbf{P} \mid x, y, z \text{ collinear}\}.$$

⁵In fact, complete intersection K3 surfaces are special cases of Calabi–Yau complete intersections considered in [14] and so all results in *loc.cit.* apply.

Proposition 5.3. *We have for any $b \in B^\circ$,*

$$\begin{aligned} & \text{Im}(\text{CH}(\mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S}) \rightarrow \text{CH}(S_b \times S_b \times S_b)) \\ &= \text{Im}(\text{CH}(\mathbf{P} \times \mathbf{P} \times \mathbf{P}) \rightarrow \text{CH}(S_b \times S_b \times S_b)) \\ &+ \sum_{1 \leq i < j \leq 3} \Delta_{i,j*} \text{Im}(\text{CH}(\mathbf{P} \times \mathbf{P}) \rightarrow \text{CH}(S_b \times S_b)) \\ &+ \delta_* \text{Im}(\text{CH}(\mathbf{P}) \rightarrow \text{CH}(S_b)), \end{aligned}$$

where $\Delta_{i,j} : S_b^2 \hookrightarrow S_b^3$ are the inclusions of big diagonals and $\delta : S_b \hookrightarrow S_b^3$ is the inclusion of the small diagonal.

Proof. It is straight-forward to check that (7) indeed stratifies q into projective bundles and the codimension of \mathcal{I} in $\mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S}$ is $\dim(\mathbf{P}) - 1$ (cf. [14, Lemma 1.2]) which is ≥ 2 . Moreover, it is clear that $\pi_3^* \mathcal{O}_B(1)$ restricts to the relative ample tautological line bundle on fibers of all projective bundles. All assumptions of Proposition 5.2 being satisfied, it implies that for any $b \in B^\circ$,

$$\begin{aligned} & \text{Im}(\text{CH}(\mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S}) \rightarrow \text{CH}(S_b \times S_b \times S_b)) \\ &= \text{Im}(\text{CH}(\mathbf{P} \times \mathbf{P} \times \mathbf{P}) \rightarrow \text{CH}(S_b \times S_b \times S_b)) + \iota_* \text{Im}(\text{CH}(J) \rightarrow \text{CH}(I_b)) \\ &+ \sum_{1 \leq i < j \leq 3} \Delta_{i,j*} \text{Im}(\text{CH}(\mathbf{P} \times \mathbf{P}) \rightarrow \text{CH}(S_b \times S_b)) + \delta_* \text{Im}(\text{CH}(\mathbf{P}) \rightarrow \text{CH}(S_b)), \end{aligned}$$

where $\iota : I_b \hookrightarrow S_b^3$ is the inclusion of the Zariski closure of the locus of collinear and distinct triples. We only have to show that the second term on the right hand side is redundant. Indeed, for $b \in B^\circ$ consider the cartesian square

$$\begin{array}{ccc} \Delta_{12} \cup \Delta_{13} \cup \Delta_{23} \cup I_b \hookrightarrow & S_b \times S_b \times S_b & \\ \downarrow & \square & \downarrow \\ J \hookrightarrow & \mathbf{P} \times \mathbf{P} \times \mathbf{P} & \end{array}$$

Here the intersection is transversal along $I_b \setminus \cup \Delta_{i,j}$ (without excess intersection) and $\text{codim}_{S_b^3} I_b = \text{codim}_{\mathbf{P} \times \mathbf{P} \times \mathbf{P}}(J) = \dim P - 1$, while along $\Delta_{i,j}$ the intersection has excess dimension $\dim P - 3$ (cf. [14, Lemma 1.2]) with excess normal bundle $\frac{\text{pr}_1^*(E|_{S_b})}{\mathcal{O}(1) \boxtimes \mathcal{O}(-1)}$ (cf. [14, Lemma 1.5])⁶. The excess intersection class on $\Delta_{i,j} = S_b \times S_b$ is therefore a polynomial in h_1 and h_2 with $h_i := \text{pr}_i^*(c_1(\mathcal{O}(1)|_{S_b}))$, hence is the pull-back of an element in $\text{CH}(\mathbf{P} \times \mathbf{P})$. As a result, by the excess intersection formula (cf. [18, §6.3]) applied to the above cartesian square, any element in the second term $\iota_* \text{Im}(\text{CH}(J) \rightarrow \text{CH}(I_b))$, up to an element in the third term $\sum_{1 \leq i < j \leq 3} \Delta_{i,j*} \text{Im}(\text{CH}(\mathbf{P} \times \mathbf{P}) \rightarrow \text{CH}(S_b \times S_b))$, is an element in the first term $\text{Im}(\text{CH}(\mathbf{P} \times \mathbf{P} \times \mathbf{P}) \rightarrow \text{CH}(S_b \times S_b \times S_b))$, thus is redundant. \square

We are now ready to prove the remaining cases of Theorem 1.4:

Proof of Theorem 1.4 for relative cubes. Denote by $h = c_1(\mathcal{O}_{\mathbf{P}}(1)|_{S_b})$ and $h_i := \text{pr}_i^*(h)$. Thanks to Proposition 5.3, for any $z \in \text{CH}(\mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S})$ and any $b \in B^\circ$, the restriction $z|_{S_b \times S_b \times S_b}$ is a polynomial in $h_1, h_2, h_3, \Delta_{12}, \Delta_{13}, \Delta_{23}$ (and $\delta = \Delta_{12}\Delta_{23}$). We can conclude by the $m = 3$ case of Voisin's [41, Proposition 2.2], where the essential point is the decomposition of the small diagonal due to Beauville–Voisin [7, Proposition 3.2]. \square

⁶So there is no excess intersection in the case of quartic surfaces.

Proof of Theorem 1.4 for relative Hilbert cubes. To simplify the notation, we denote by $S^{[m]} := \text{Hilb}^m(S)$ and similarly $S^{[m]/B} := \text{Hilb}_B^m S$. Let us first recall the result of de Cataldo–Migliorini [11] in the special case of Chow groups of Hilbert cubes of surfaces: for any surface S , denote by $\rho : S^{[3]} \rightarrow S^{(3)}$ the Hilbert–Chow morphism which sends a 0-dimensional subscheme to its support 0-cycle. We have the incidence subvarieties

$$\begin{aligned} U &:= \{(z, x_1, x_2, x_3) \in S^{[3]} \times S^3 \mid \rho(z) = x_1 + x_2 + x_3\}; \\ V &:= \{(z, x_1, x_2) \in S^{[3]} \times S^2 \mid \rho(z) = 2x_1 + x_2\}; \\ W &:= \{(z, x) \in S^{[3]} \times S \mid \rho(z) = 3x\}; \end{aligned}$$

and the main result of [11] says that they together induce an injective morphism

$$(U_*, V_*, W_*) : \text{CH}(S^{[3]}) \hookrightarrow \text{CH}(S^3) \oplus \text{CH}(S^2) \oplus \text{CH}(S).$$

Note that the above correspondences have natural family counterparts, denoted by $\mathcal{U}, \mathcal{V}, \mathcal{W}$.

Let $z \in \text{CH}(S^{[3]/B})$ be such that the cohomology class of $z|_{S_b^{[3]}}$ vanishes. By the above injectivity, it is enough to show that for any $b \in B^\circ$, $U_* \left(z|_{S_b^{[3]}} \right)$, $V_* \left(z|_{S_b^{[3]}} \right)$ and $W_* \left(z|_{S_b^{[3]}} \right)$ are zero. To this end, observe that $U_* \left(z|_{S_b^{[3]}} \right) = \mathcal{U}_*(z)|_{S_b^3}$ is the restriction of a cycle of the total family $\mathcal{S} \times_B \mathcal{S} \times_B \mathcal{S}$ with trivial cohomology class, hence is zero by the relative cube case of Theorem 1.4 just proved. Similarly, the vanishing of $V_* \left(z|_{S_b^{[3]}} \right)$ and $W_* \left(z|_{S_b^{[3]}} \right)$ follow from the relative square case proved in §4 and [33] respectively.

Finally, the proof of the case of $\mathcal{S} \times_B S^{[2]/B}$ is similar (easier). \square

5.3. Beyond complete intersection K3 surfaces. The techniques we utilized above in order to prove Theorem 1.4 for (Hilbert) squares and cubes of complete intersection K3 surfaces can also be employed to attack the Generalized Franchetta Conjecture 1.3 for families of K3 surfaces for which Mukai models are available. In this subsection, we give a sufficient condition for the conjecture to hold for Hilbert schemes of K3 surfaces in a certain range.

Recall that for a natural number g , we say that the *Mukai model* for K3 surfaces of genus g exists, if there exist an ambient homogeneous space $G = G_g$ (often a Grassmannian) and a globally generated homogeneous vector bundle $E = E_g$ on G , such that the zero locus of a general section of E gives a general K3 surface of genus g . For the available constructions of Mukai models and the corresponding G and E , we refer to [33] as well as the original sources [26], [27], [28], [29].

The crucial condition for our techniques to work is the following:

Definition 5.4. For an $r \in \mathbf{N}^*$, we say that the Mukai model (G, E) satisfies the condition (\star_r) if

(\star_r) : for any x_1, \dots, x_r distinct points of G , the following evaluation map is surjective

$$H^0(G, E) \rightarrow \bigoplus_{i=1}^r E_{x_i}.$$

Or equivalently, $H^0(G, E \otimes I_{x_1} \otimes \dots \otimes I_{x_r})$ is of codimension $r \cdot \text{rank}(E)$ in $H^0(G, E)$. Clearly, (\star_r) implies (\star_k) for all $k < r$.

Proposition 5.5. *Notation is as before. Fix a genus g for which the Mukai model exists. Assume Condition (\star_r) is satisfied. Then for any $b \in B^\circ$,*

$$\mathrm{Im}(\mathrm{CH}(\mathcal{S}^{r/B}) \rightarrow \mathrm{CH}(S_b^r)) = R(S_b^r),$$

where the tautological ring $R(S^r)$ is the subring of the (rational) Chow ring $\mathrm{CH}(S^r)$ generated by big diagonals $\Delta_{i,j}$ ($1 \leq i < j \leq r$), polarization classes $g_i := \mathrm{pr}_i^*(g)$ and Beauville–Voisin classes $\mathfrak{o}_i := \mathrm{pr}_i^*(\mathfrak{o}_S)$ ($1 \leq i \leq r$).

Proof. The proof is to rephrase every step of §5.2 in the general setting. We proceed by induction on r . Consider the evaluation map $q : \mathcal{S}^{r/B} \rightarrow G^r$, which is a stratified projective bundle (Definition 5.1) with the stratification on G^r given by the different types of incidence relations for r points of G :

$$(8) \quad \begin{array}{ccccccc} \mathcal{X}_n = \mathcal{S} & \hookrightarrow & \dots & \hookrightarrow & \mathcal{X}_1 & \hookrightarrow & \mathcal{X}_0 = \mathcal{S}^{r/B} \longrightarrow B \\ \downarrow q_n = p \quad \square & & & & \downarrow q_1 \quad \square & & \downarrow q_0 = q \\ Y_n = G & \hookrightarrow & \dots & \hookrightarrow & Y_1 & \hookrightarrow & Y_0 = G^r \end{array}$$

By Proposition 5.2, for any $b \in B^\circ$,

$$(9) \quad \mathrm{Im}(\mathrm{CH}(\mathcal{S}^{r/B}) \rightarrow \mathrm{CH}(S_b^r)) = \sum_{i=0}^n \iota_{i,*} \mathrm{Im}(\mathrm{CH}(Y_i) \rightarrow \mathrm{CH}(\mathcal{X}'_i)),$$

where \mathcal{X}'_i is the Zariski closure of $\mathcal{X}_i \setminus \mathcal{X}_{i+1}$. Let us show that each term of (9) is in the tautological ring $R(S_b)$ by ascending order for $0 \leq i \leq n$:

- If $i = 0$, since the Chow ring of G satisfies the Künneth formula, we only need to show that

$$\mathrm{Im}(\mathrm{CH}(G) \rightarrow \mathrm{CH}(S_b)) \subset R(S_b).$$

However, since $\mathrm{CH}(G)$ is generated by the Chern classes of its tautological subbundle, so it is enough to check that $c_1(G)|_{S_b}$ and $c_2(G)|_{S_b} \in R(S_b)$. The first one is precisely the polarization class and the second one follows from a direct computation using the short exact sequence

$$0 \rightarrow T_{S_b} \rightarrow T_G|_{S_b} \rightarrow E|_{S_b} \rightarrow 0,$$

together with the fact that $c_2(S_b) = 24\mathfrak{o}_{S_b}$ ([7]).

- If a general point of Y_i is parameterizing r points of G where at least two of them coincide, then the contribution of the i -th term of (9) factors through $R(S_b^{r-1})$ by the induction hypothesis, hence is contained⁷ in $R(S_b^r)$.
- If a general point of Y_i is parameterizing r different points of G , then the hypothesis (\star_r) means precisely that any r different points of G impose independent conditions on B , each of codimension $\mathrm{rank}(E)$. Therefore, \mathcal{X}'_i , the Zariski closure of $\mathcal{X}_i \setminus \mathcal{X}_{i+1}$, has the same codimension in \mathcal{X}_{i-1} as $\mathrm{codim}_{Y_{i-1}}(Y_i)$. The excess intersection formula ([18, §6.3]) applied to the cartesian diagram

$$\begin{array}{ccc} \mathcal{X}_i = \mathcal{X}_{i+1} \cup \mathcal{X}'_i & \hookrightarrow & \mathcal{X}_{i-1} \\ \downarrow & \square & \downarrow \\ Y_i & \hookrightarrow & Y_{i-1} \end{array}$$

⁷It is easy to show that tautological rings of different powers of a K3 surface are closed under push-forward by diagonal inclusion.

tells us that modulo the $(i + 1)$ -th term of (9), the contribution of the i -th term is contained in the $(i - 1)$ -th term.

□

Theorem 5.6. *Let $g \in \mathbf{N}^*$ such that the Mukai model exists for K3 surfaces of genus g . Assume the Mukai model satisfies the condition (\star_r) . If moreover, the cycle class map restricted to the tautological ring $R(S^r)$ is injective for any K3 surface S of genus g , then the Generalized Franchetta Conjecture 1.3 holds for $\mathcal{S}^{[r_1]/B} \times_B \cdots \times_B \mathcal{S}^{[r_m]/B}$, for any r_1, \dots, r_m whose sum $\leq r$, where $\mathcal{S} \rightarrow B$ is the universal family of K3 surfaces of genus g .*

Proof. The case of relative powers $\mathcal{S}^{k/B}$, for any $k \leq r$ is a direct consequence of Proposition 5.5 and the hypothesis on the injectivity of the cycle class map on the tautological ring. For other cases, we use de Cataldo–Migliorini’s result [11] for Chow motives of Hilbert schemes of surfaces to reduce to the cases of $\mathcal{S}^{k/B}$ for all $1 \leq k \leq r$. □

We apply Theorem 5.6 to some Mukai models to get concrete unconditional results :

Proof of Theorem 1.5. To use Theorem 5.6, we proceed by a case-by-case analysis of the positivity of the homogeneous bundle in the Mukai model. See Mukai’s series of papers [26], [27], [28], [29] for more information on the geometry of these models.

- K3 surfaces of genus $g = 2$ are⁸ smooth degree 6 hypersurfaces in the weighted projective space $\mathbf{P} := \mathbf{P}(1, 1, 1, 3)$. The Mukai model for this family is thus $(G, E) = (\mathbf{P}, \mathcal{O}(6))$. Note that the K3 surfaces in this family all avoid the singular point $O := [0, 0, 0, 1]$. Let us check the condition (\star_3) , *i.e.*, that the evaluation map

$$H^0(\mathbf{P}, \mathcal{O}(6)) \rightarrow \bigoplus_{i=1}^3 \mathbf{C}_{x_i}$$

is surjective for distinct $x_1, x_2, x_3 \neq O$, where \mathbf{C}_x denotes the fiber of $\mathcal{O}(6)$ at x . It is easy to see that $\mathbf{P}(1, 1, 1, 3)$ is isomorphic to the projective cone over the third Veronese embedding of \mathbf{P}^2 (*cf.* [12]) and O is the vertex. By upper-semicontinuity, it is enough to treat the most degenerate case for three distinct points of $\mathbf{P} \setminus \{O\}$, which is when they lie in the same ruling of the projective cone. In this case, as the restriction of $\mathcal{O}(6)$ to the ruling is $\mathcal{O}(2)$, the condition (\star_3) follows from the surjections :

$$H^0(\mathbf{P}, \mathcal{O}(6)) \twoheadrightarrow H^0(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}(2)) \twoheadrightarrow \bigoplus_{i=1}^3 \mathbf{C}_{x_i},$$

where \mathbf{P}^1 is the ruling which contains x_i ’s. To conclude, by Theorem 5.6, we only need to know the injectivity of the cycle class map restricted to the tautological ring $R(S^3)$, which is covered by Voisin’s [41, Proposition 2.2].

- For quartic surfaces ($g = 3$), let us first show that $(\mathbf{P}^3, \mathcal{O}(4))$ satisfies (\star_5) , *i.e.*, that the evaluation map

$$H^0(\mathbf{P}^3, \mathcal{O}(4)) \rightarrow \bigoplus_{i=1}^5 \mathbf{C}_{x_i}$$

is surjective for distinct x_i ’s. Again, it is enough to treat the most degenerate cases, namely :

⁸Equivalently, these K3 surfaces are also double covers of \mathbf{P}^2 ramified along smooth sextic curves.

- when x_1, \dots, x_5 are collinear, then this follows from the surjectivity of the restriction and the evaluation

$$H^0(\mathbf{P}^3, \mathcal{O}(4)) \twoheadrightarrow H^0(\mathbf{P}^1, \mathcal{O}(4)) \twoheadrightarrow \bigoplus_{i=1}^5 \mathbf{C}_{x_i},$$

where \mathbf{P}^1 is the line containing these points.

- when x_1, \dots, x_5 are in a conic C . Then the Koszul resolution provides an exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbf{P}^3}(-3) \rightarrow \mathcal{O}_{\mathbf{P}^3}(-1) \oplus \mathcal{O}_{\mathbf{P}^3}(-2) \rightarrow \mathcal{O}_{\mathbf{P}^3} \rightarrow \mathcal{O}_C \rightarrow 0,$$

which allows us to see that the restriction map $H^0(\mathbf{P}^3, \mathcal{O}(4)) \rightarrow H^0(C, \mathcal{O}_C(8))$ is surjective.

Since $H^0(C, \mathcal{O}_C(8)) \rightarrow \bigoplus_{i=1}^5 \mathbf{C}_{x_i}$ is clearly surjective, we are done.

The condition (\star_5) is proved. By Voisin [41, Proposition 2.2], the cycle class map restricted to the tautological ring $R(S^5)$ is injective for a quartic K3 surface S . Now both assumptions of Theorem 5.6 being satisfied, we can conclude.

- For $g = 6$, the Mukai model is $(G, E) = (\text{Gr}(2, 5), \mathcal{O}(1)^{\oplus 3} \oplus \mathcal{O}(2))$, where $\mathcal{O}(1)$ is the Plücker line bundle. It is clear that the condition (\star_2) is equivalent to the surjectivity of

$$H^0(G, \mathcal{O}(1)) \rightarrow \mathbf{C}_{x_1} \oplus \mathbf{C}_{x_2}$$

for any two distinct points $x_1, x_2 \in G$. However, the last condition follows from the very ampleness of the Plücker line bundle $\mathcal{O}(1)$.

- For $g = 7$, the Mukai model is $(G, E) = (\text{OGr}(5, 10), U^{\oplus 8})$, where $\text{OGr}(5, 10)$ is the orthogonal Grassmannian parameterizing isotropic subspaces of dimension 5 in a vector space of dimension 10 equipped with a non-degenerate quadratic form and U is a line bundle corresponding to the half spinor representation. The proof is similar to the previous case: one uses the very ampleness of U .
- For $g = 8$, the Mukai model is $(G, E) = (\text{Gr}(2, 6), \mathcal{O}(1)^{\oplus 6})$, where $\mathcal{O}(1)$ is the Plücker line bundle. The proof goes as for $g = 6$ by the very ampleness of the Plücker line bundle.
- For $g = 9$, the Mukai model is $(G, E) = (\text{LGr}(3, 6), \mathcal{O}(1)^{\oplus 4})$, where $\text{LGr}(3, 6)$ is the symplectic Grassmannian parameterizing Lagrangian subspaces in a 6-dimensional vector space equipped with a symplectic form and $\mathcal{O}(1)$ is the restriction of the Plücker line bundle of $\text{Gr}(3, 6)$. The proof goes as before: one uses the very ampleness of $\mathcal{O}(1)$.
- For $g = 10$, the Mukai model is $(G, E) = (G_2/P, \mathcal{O}(1)^{\oplus 3})$, where G is the 5-dimensional quotient of the simply-connected semi-simple algebraic group of type G_2 by a maximal parabolic subgroup P and $\mathcal{O}(1)$ is the line bundle associated to the adjoint representation of G_2 ; in other words, $G = G_2/P \hookrightarrow \mathbf{P}(\mathfrak{g}_2^\vee)$. Again, we can conclude by the very ampleness of $\mathcal{O}(1)$.
- For $g = 12$, we use a slight variant of the above argument. Indeed, the general genus 12 K3 surface can be constructed as an anti-canonical section in a smooth genus 12 prime Fano threefold X (cf. [4], [20, Section 3.1]). The Fano threefold X has very ample anti-canonical bundle, and $H^3(X, \mathbf{Q}) = 0$ ([21, Corollary 4.3.5]) so that X has trivial Chow groups⁹ (this Fano threefold X is the variety denoted by $X_{22} \subset \mathbf{P}^{13}$ in [21, Propositions 4.1.11 and 4.1.12]; actually X is an intersection of quadrics). We now consider a variant of Theorem 5.6, replacing G by X and E by $-K_X$. The very ampleness of $-K_X$ ensures that condition (\star_2) holds. As X has trivial Chow groups, there is a Chow–Künneth formula for products of X , and so one is reduced to the statement for the K3 surface S_b , which is [33].

⁹Following Voisin [42], we say a smooth projective variety has *trivial Chow groups* if the cycle map $\text{cl}^i : \text{CH}^i(X)_{\mathbf{Q}} \rightarrow H^{2i}(X, \mathbf{Q})$ is injective for any i .

□

Remark 5.7 (Limit of our method). Given a Mukai model (G, E) ,

- the global generation of E corresponds to condition (\star_1) , which explains the essential reason why one can prove the Generalized Franchetta Conjecture for K3 surfaces with a Mukai model ([33]).
- For K3 surfaces of genus 2, $G = \mathbf{P}(1, 1, 1, 3)$ and $E = \mathcal{O}(6)$, the condition (\star_4) is not satisfied : it is violated by three distinct points lying on the same ruling, away from the singular point.
- For the quartic K3 surfaces, $G = \mathbf{P}^3$ and $E = \mathcal{O}(4)$, the condition (\star_6) is not satisfied : it is violated by six collinear distinct points. Similarly, for the other two families of complete intersection K3 surfaces (genus 4 and 5), (\star_4) is violated by four collinear distinct points.
- For K3 surfaces of genus 6 and 8, whose Mukai model is $(G, E) = (\text{Gr}(2, 5), \mathcal{O}(1)^{\oplus 3} \oplus \mathcal{O}(2))$ and $(\text{Gr}(2, 6), \mathcal{O}(1)^{\oplus 6})$ respectively, the condition (\star_3) is not satisfied. Indeed, it is equivalent to the surjectivity of $H^0(G, \mathcal{O}(1)) \rightarrow \mathbf{C}_{x_1} \oplus \mathbf{C}_{x_2} \oplus \mathbf{C}_{x_3}$, which is violated by three distinct collinear points of G .
- For K3 surfaces of genus 13 and 20, the Mukai models are respectively

$$(G, E) = \left(\text{Gr}(3, 7), (\wedge^2 S^\vee)^{\oplus 2} \oplus \wedge^3 Q \right) \text{ and } \left(\text{Gr}(4, 9), (\wedge^2 S^\vee)^{\oplus 3} \right).$$

where S is the tautological sub-bundle and Q is the tautological quotient bundle. We claim that none of them verifies the condition (\star_2) . For example, in the genus 13 case, the condition (\star_2) is equivalent to the surjectivities of the following two evaluation maps

$$H^0(G, \wedge^2 S^\vee) \rightarrow \wedge^2 S_x^\vee \oplus \wedge^2 S_y^\vee,$$

$$H^0(G, \wedge^2 Q) \rightarrow \wedge^2 Q_x \oplus \wedge^2 Q_y,$$

for any $x \neq y \in G$, which, by Bott theorem, amount to say that for any two different 3-dimensional subspaces W_1, W_2 in a 7-dimensional vector space V , the natural maps

$$\wedge^2 V^\vee \rightarrow \wedge^2(W_1^\vee) \oplus \wedge^2(W_2^\vee)$$

$$\wedge^2 V \rightarrow \wedge^2(V/W_1) \oplus \wedge^2(V/W_2)$$

are surjective. It is not true when $\dim W_1 \cap W_2 \geq 2$. The case of genus 20 is similar.

- For K3 surfaces of genus 18, the Mukai model is $(G, E) = (\text{OGr}(3, 9), U^{\oplus 5})$, where U is the rank 2 vector bundle associated to the representation V associated to the fourth dominant weight $\omega_4 = \frac{1}{2}(\alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4)$, of the (semi-simple part of the) maximal parabolic group P . We claim that (\star_2) does not hold, *i.e.*, there exist two different points $x, y \in G$ such that $H^0(G, U) \rightarrow U_x \oplus U_y$ is not surjective¹⁰. Let $x = P/P$ and $y = wP/P$ where $w = s_{\alpha_3}$, as an element in the Weyl group W , is the reflection with respect to the third simple root. Clearly, w does not belong to the Weyl group of P , which is generated by $s_{\alpha_1}, s_{\alpha_2}, s_{\alpha_4}$. A direct computation shows that the representation $H^0(G, U)$ has multiplicity one for all weights. On the other hand, ω_4 is a common weight for V and its conjugate by w (since $w \cdot \omega_4 = \omega_4$). Hence $H^0(G, U) \rightarrow U_x \oplus U_y$ cannot be surjective.
- If one wants to follow the same strategy of this paper to attack the Generalized Franchetta Conjecture 1.3 for (Hilbert) powers beyond the range stated in Theorem 1.4 and Theorem 1.5, one has to deal with some essentially new universal cycle, which may not belong to the tautological ring, or rather, the tautological ring should be enlarged to include some more incidence classes from projective geometry than just the polarization class.

¹⁰We thank Nicolas Ressayre for his kind help on the proof.

Nevertheless, the injectivity of the cycle class map restricted to the *enlarged* tautological ring is a consequence of the general Bloch-Beilinson conjecture over number fields.

5.4. Applications towards the Beauville–Voisin conjecture. Let us now turn to the consequences of our results in the direction of the Beauville–Voisin conjecture (and its refined version Conjecture 2.4):

Proof of Corollaries 1.6 and 1.7 and 1.8. The coisotropic subvarieties E_μ , the Lagrangian surfaces T and U , and the Lagrangian threefold I of Corollary 1.8, can all be defined over (suitable relative powers of) the universal family, and so these are just special cases of Proposition 2.5, combined with Theorems 1.4 and 1.5. \square

One can also prove a version of Corollaries 1.6 and 1.7 and 1.8 for product varieties of arbitrarily high dimension, but the statement is now restricted to 0-cycles and 1-cycles:

Corollary 5.8. *Let X be a product*

$$X = X_1 \times X_2 \times \cdots \times X_s, \quad \dim X = 2m,$$

where X_j is either a Hilbert square $S^{[2]}$ where S is a K3 surface of genus $g \in \{2, 3, 4, 5, 6, 7, 8, 9, 10, 12\}$, or a Hilbert cube $S^{[3]}$ where S is a degree 6 complete intersection K3, or a Hilbert scheme $S^{[r]}$ where S is a quartic K3 surface and $r \leq 5$. Let $\widetilde{R}(X) \subset \text{CH}(X)$ denote¹¹ the \mathbf{Q} -subalgebra generated by (pullbacks of) divisors on X_j , the Chern classes $c_i(T_{X_j})$, plus the following coisotropic subvarieties:

- the coisotropic subvarieties E_μ of [44, 4.1 item 1)];
- the Lagrangian surfaces $T \subset X_j$ constructed in [20, Proposition 4] (if $X_j = S^{[2]}$);
- the surface of bitangents $U \subset X_j$ (if $X_j = S^{[2]}$ and S is a quartic K3 surface);
- the locus $I \subset X_j$ as in Corollary 1.8 (if $X_j = S^{[3]}$ and S is a K3 surface of degree 6).

Then $\widetilde{R}^{2m}(X)$ and $\widetilde{R}^{2m-1}(X)$ inject into cohomology via the cycle class map.

Proof of Corollary 5.8. This uses the fact that the X_j have a *multiplicative Chow–Künneth decomposition* $\{\pi_{X_j}^k\}$, in the sense of [35, Chapter 8], [36]; see also Appendix A.2. This induces a bigrading of the Chow ring of X_j , given by

$$\text{CH}^i(X_j)_{(k)} := (\pi_{X_j}^{2i-k})_* \text{CH}^i(X_j).$$

It is readily seen that the projectors $\pi_{X_j}^k$ are universally defined (*i.e.*, they exist as a relative cycle for the family $\mathcal{X}_j^\circ \times_{B^\circ} \mathcal{X}_j^\circ$). Theorem 5.6 applied to the relative cycle $\mathcal{T} - (\pi_{X_j}^2)_*(\mathcal{T})$ (where we use the formalism of relative correspondences as in [30, Section 8.1]), thus implies that

$$T \in \text{CH}^2(X_j)_{(0)}.$$

Similarly $U \in \text{CH}^2(X_j)_{(0)}$ and $I \in \text{CH}^3(X_j)_{(0)}$. The same argument also shows that $E_\mu \in \text{CH}^*(X_j)_{(0)}$, which is actually true for Hilbert schemes of arbitrary K3 surfaces, *cf.* [44, Lemma 4.3].

The product X also has a multiplicative Chow–Künneth decomposition, and hence there is a bigrading of the Chow ring $\text{CH}(X)$, by [35, Theorem 8.6]. Since divisors and Chern classes of

¹¹In this paper, the notation $R(X)$ is reserved for the tautological ring of $X = S^n$ a power of K3 surface, generated by big diagonals and Chern classes.

X_j are also in $\text{CH}(X_j)_{(0)}$, and pullback under any projection $X \rightarrow X_j$ preserves the bigrading [36, Corollary 1.6], we see that there is an inclusion

$$\widetilde{R}(X) \subset \text{CH}(X)_{(0)}.$$

The corollary now follows, since it is known that $\text{CH}^i(X)_{(0)}$ injects into cohomology for $i \geq \dim(X) - 1$, see [39, Introduction]. \square

5.5. Double EPW sextics. The interested reader will have no trouble finding further applications in the flavour of Corollaries 1.6, 1.7 and 1.8. For instance, consider the Hilbert square $X = S^{[2]}$, where S is a generic K3 surface of genus 6. As shown by O’Grady [31, Section 4], X is isomorphic to a small resolution X_A^ϵ of a singular double EPW sextic X_A (notation is as in [31]). Let $\epsilon: X \rightarrow X_A$ denote the small resolution, and let $f_A: X_A \rightarrow Y_A$ denote the double cover to the associated EPW sextic Y_A . The surface S being generic corresponds to the fact that the Lagrangian vector space A is generic (in the precise sense given in [31, §4]) in the divisor $\Delta \subset \text{LGr}(\wedge^3 V)$ studied in [31]. This construction produces Lagrangian surfaces in X : the surface

$$P := \epsilon^{-1}(\text{Sing}(X_A))$$

(which is isomorphic to \mathbf{P}^2 since X_A has only one singular point), and the surface

$$\text{Fix} := \epsilon^{-1}(\text{Fix}(\iota)),$$

where $\text{Fix}(\iota)$ denotes the fixed point locus of the (anti-symplectic) covering involution ι of X_A . These Lagrangian surfaces are easily seen to be universally defined. (Indeed, as shown in [31], there is a stratification

$$Y_A[3] \subset Y_A[2] \subset Y_A[1] = Y_A$$

of the EPW sextic Y_A . Here the surface $Y_A[2]$ is the singular locus of Y_A and the point $Y_A[3]$ is the unique singularity of $Y_A[2]$. One has

$$\text{Fix} = (f_A \circ \epsilon)^{-1}(Y_A[2]), \quad P = (f_A \circ \epsilon)^{-1}(Y_A[3]).$$

On the other hand (as explained in [31, Section 3]), there exist family versions $\mathcal{Y}[i]$ of the subvarieties $Y_A[i]$ over the base Δ . One can do a base change

$$\begin{array}{ccc} \overline{\mathcal{X}}_{B^0} & \rightarrow & \overline{\mathcal{X}} \\ \downarrow & & \downarrow \\ \mathcal{Y}_{B^0} & \rightarrow & \mathcal{Y} \\ \downarrow & & \downarrow \\ B^0 & \rightarrow & \Delta \end{array}$$

where $B^0 \subset B$ is an open such that the rational map $B \dashrightarrow \Delta$ of [31, Section 4] is defined, and $\overline{\mathcal{X}}$ is the tautological family of singular double EPW sextics over Δ . One obtains relative versions of P and Fix by pulling back $\mathcal{Y}[i]$ under the birational morphism $\mathcal{X}_{B^0} \rightarrow \overline{\mathcal{X}}_{B^0}$.) Thus, applying Theorem 1.5 one obtains the following:

Corollary 5.9. *Let $X = S^{[2]}$, where S is a generic K3 surface of genus 6. The \mathbf{Q} -subalgebra*

$$\langle D_1, D_2, c_i(T_X), P, \text{Fix}, T \rangle \subset \text{CH}(X)$$

injects into cohomology via the cycle class map. (Here D_1, D_2 are two divisors generating the Picard group of X , and T is the Lagrangian surface of [20, Proposition 4].)

5.6. An application to Bloch's conjecture. Given a quartic K3 surface S , Beauville [3] constructed an interesting involution ι on $X := S^{[2]}$, which generically sends $\{x_1, x_2\}$ to $\{x_3, x_4\}$, where x_1, \dots, x_4 are the four intersection points of the line $\overline{x_1, x_2}$ with S . The involution ι is anti-symplectic. According to the generalized Bloch conjecture (cf. [40, §11.2]), which roughly says that CH_0 is 'controlled' by the holomorphic forms, the action of ι on $\text{CH}_0(X)$ should be id on $\text{Gr}_F^0 \text{CH}_0(X)$ and $\text{Gr}_F^4 \text{CH}_0(X)$ (just as on $H^0(X)$ and $H^{4,0}(X)$) and should be $-\text{id}$ on $\text{Gr}_F^2 \text{CH}_0(X)$ (just as on $H^{2,0}(X)$), where F is the conjectural Bloch–Beilinson filtration. On the other hand, as conjectured in [5] by Beauville and worked out by Shen–Vial in [35] in the case of Hilbert squares of K3 surfaces, we have a canonical splitting of this filtration for X , giving a direct sum decomposition:

$$\text{CH}^4(X) = \text{CH}^4(X)_{(0)} \oplus \text{CH}^4(X)_{(2)} \oplus \text{CH}^4(X)_{(4)}.$$

Hence the action of ι on the three summands should be id , $-\text{id}$ and id , respectively. Our results allow us to confirm this expectation.

Proof of Corollary 1.9. Let $\mathcal{S}^\circ \rightarrow B^\circ$ be the universal family of smooth quartic K3 surfaces and $\mathcal{X}^\circ \rightarrow B^\circ$ be the relative Hilbert square. As noted above, the bigrading $\text{CH}^*(X)_{(*)}$ is induced by a self-dual multiplicative Chow–Künneth decomposition $\{\pi_X^k\}$ that is universally defined. The anti-symplectic involution ι can also be defined on the level of the universal family; let us denote $\Gamma_\iota \in \text{CH}^4(\mathcal{X}^\circ \times_{B^\circ} \mathcal{X}^\circ)$ the graph of the involution $\iota: \mathcal{X}^\circ \rightarrow \mathcal{X}^\circ$.

The relative correspondence

$$\pi_X^i \circ \Gamma_\iota \circ \pi_X^j \in \text{CH}^4(\mathcal{X}^\circ \times_{B^\circ} \mathcal{X}^\circ)$$

is fiberwise homologically trivial for $i \neq j$. Theorem 1.5(ii) for $\text{Hilb}_B^2 \mathcal{S} \times_B \text{Hilb}_B^2 \mathcal{S}$ implies that

$$(10) \quad (\pi_X^i \circ \Gamma_\iota \circ \pi_X^j)|_{X_b \times X_b} = 0 \quad \text{in } \text{CH}^4(X_b \times X_b), \quad \forall i \neq j \quad \forall b \in B^0,$$

i.e., Γ_{ι_b} belongs to $\text{CH}^4(X_b \times X_b)_{(0)}$, and thus ι_b preserves the bigrading $\text{CH}^*(X_b)_{(*)}$.

Next, the fact that ι_b is anti-symplectic means that for any $b \in B^\circ$ there exists a divisor $D_b \subset X_b$, and a cycle γ_b supported on $D_b \times D_b$, such that

$$((\Delta_X + \Gamma_\iota) \circ \pi_X^2)|_{X_b \times X_b} = \gamma_b \quad \text{in } H^8(X_b \times X_b).$$

Using a Hilbert schemes argument as in [42, Proposition 3.7], the D_b and γ_b can be spread out, i.e., there exists a divisor $\mathcal{D} \subset \mathcal{X}$ and a relative cycle γ supported on $\mathcal{D} \times_{B^\circ} \mathcal{D}$ such that

$$((\Delta_X + \Gamma_\iota) \circ \pi_X^2 - \gamma)|_{X_b \times X_b} = 0 \quad \text{in } H^8(X_b \times X_b) \quad \forall b \in B^0.$$

Applying Theorem 1.5 once more, we find that

$$(11) \quad ((\Delta_X + \Gamma_\iota) \circ \pi_X^2 - \gamma)|_{X_b \times X_b} = 0 \quad \text{in } \text{CH}^4(X_b \times X_b) \quad \forall b \in B^0.$$

For general $b \in B^0$, the restriction $\gamma|_{X_b \times X_b}$ will be supported on (divisor) \times (divisor), and so $\gamma|_{X_b \times X_b}$ will act as 0 on $\text{CH}^2(X_b)_{(2)}$. It follows that

$$(\iota_b)^* = -\text{id}: \quad \text{CH}^2(X_b)_{(2)} \rightarrow \text{CH}^2(X_b)_{(2)} \quad \text{for general } b \in B^0.$$

To extend this to *all* $b \in B^\circ$, one notes that the above construction can be done with a divisor $\mathcal{D} \subset \mathcal{X}$ in general position with respect to X_b .

The statement for $\text{CH}^4(X_b)_{(2)}$ follows upon taking the transpose of relation (11), and using the relation (10). The remaining statements are proven similarly. \square

Remark 5.10. Corollary 1.9 was proven in a more convoluted way in [23].

6. LEHN-LEHN-SORGER-VAN STRATEN HYPER-KÄHLER EIGHTFOLDS

In this section we first show Theorem 1.11 and then deduce from it Theorem 1.12.

Keep the same notation as in §3. We still have a correspondence :

$$\begin{array}{ccc} & \mathcal{F} \times_B \mathcal{F} & \\ \pi_2 := (\pi, \pi) \swarrow & & \searrow q := (p, p) \\ B & & G \times G \end{array}$$

However the problem is that q is no longer a projective bundle: the fiber of q over a pair of lines (l, l') is the subspace of cubic fourfolds containing both l and l' , whose dimension depends therefore on the relative position of (l, l') . To adapt the same strategy to this case, we use similar techniques as in [42], [16] by studying the various strata of the morphism q . There are three possible relative positions between two projective lines in \mathbf{P}^5 : identical, intersect but not identical, not intersect.

On the one hand, for a (general) cubic fourfold X with Fano variety of lines F , let

$$I := \{(l, l') \in F \times F \mid l \cap l' \neq \emptyset\}$$

be the 6-dimensional incidence subvariety of $F \times F$. The incidence subvariety I has two natural projections to F with fiber over $l \in F$ the surface S_l parameterizing lines inside X meeting l . Similarly, we consider the family version of this incidence subvariety inside $\mathcal{F} \times_B \mathcal{F}$:

$$I := \{(b, l, l') \in F \times_B F \mid l \cap l' \neq \emptyset\} = \{(b, l, l') \in B \times G \times G \mid l, l' \subset X_b; l \cap l' \neq \emptyset\}.$$

On the other hand, we define $J := \{(l, l') \in G \times G \mid l \cap l' \neq \emptyset\}$ to be the incidence subvariety of $G \times G$.

These incidence subvarieties, together with the diagonals, give the stratification for q :

$$\begin{array}{ccccc} \mathcal{F} = \Delta_{\mathcal{F}/B} & \hookrightarrow & I & \hookrightarrow & \mathcal{F} \times_B \mathcal{F} & \xrightarrow{\pi_2} & B \\ \downarrow p & & \downarrow q|_I & & \downarrow q & & \\ G = \Delta_G & \hookrightarrow & J & \hookrightarrow & G \times G & & \end{array}$$

where q is a projective bundle outside of I and $q|_I$ is also a projective bundle outside of $\Delta_{\mathcal{F}}$; in other words, q is a *stratified projective bundle* in the sense of Definition 5.1.

Applying Proposition 5.2 to q , we have the following analogue of Lemma 3.1 and Proposition 4.1 in our case :

Proposition 6.1. *For any $b \in B$, we have*

$$\begin{aligned} & \text{Im}(\text{CH}(\mathcal{F} \times_B \mathcal{F}) \rightarrow \text{CH}(F_b \times F_b)) \\ = & \text{Im}(\text{CH}(G \times G) \rightarrow \text{CH}(F_b \times F_b)) + i_* \text{Im}(\text{CH}(J) \rightarrow \text{CH}(I_b)) + \Delta_* \text{Im}(\text{CH}(G) \rightarrow \text{CH}(F_b)), \end{aligned}$$

where $i : I_b \hookrightarrow F_b \times F_b$ and $\Delta : F_b \hookrightarrow F_b \times F_b$ are the inclusions.

As the incidence subvariety J is singular along the smaller stratum Δ_G , it is more convenient to work with a natural resolution of singularities. To this end, we define

$$\begin{aligned}\tilde{I} &:= \{(b, x, l, l') \in B \times \mathbf{P}^5 \times G \times G \mid l, l' \subset X_b; x \in l \cap l'\}; \\ \tilde{J} &:= \{(x, l, l') \in \mathbf{P}^5 \times G \times G \mid x \in l \cap l'\}; \\ \mathcal{P} &:= \{(b, x, l) \in B \times \mathbf{P}^5 \times G \mid l \subset X_b; x \in l\}; \\ Q &:= \{(x, l) \in \mathbf{P}^5 \times G \mid x \in l\},\end{aligned}$$

where \tilde{I} (resp. \tilde{J}) admits a natural birational morphism to I (resp. J), which contracts \mathcal{P} (resp. Q) to \mathcal{F} (resp. G). We summarize the situation in the following diagram whose squares are all cartesian :

$$\begin{array}{ccccccc} \mathcal{F} & \longleftarrow & \mathcal{P} & \longrightarrow & \tilde{I} & \longrightarrow & I & \longrightarrow & \mathcal{F} \times_B \mathcal{F} \\ \downarrow p & \square & \downarrow q' |_{\mathcal{P}} & \square & \downarrow q' & \square & \downarrow q|_I & \square & \downarrow q \\ G & \longleftarrow & Q & \longrightarrow & \tilde{J} & \longrightarrow & J & \longrightarrow & G \times G \end{array}$$

Recall that $G = \text{Gr}(\mathbf{P}^1, \mathbf{P}^5)$, S is the tautological rank 2 sub-bundle, $g := c_1(S^\vee|_F) \in \text{CH}^1(F)$ is the Plücker polarization class, and $c := c_2(S^\vee|_F) \in \text{CH}^2(F)$. We computed in Lemma 3.2 that $c_2(F) = 5g^2 - 8c$. In $\text{CH}(F \times F)$, $g_i := \text{pr}_i^*(g)$ and $c_i := \text{pr}_i^*(c)$ for $i = 1, 2$.

Definition 6.2 (Tautological ring of $F \times F$). Let X be a smooth cubic fourfold and F be its Fano variety of lines. We define the *tautological ring* of $F \times F$, denoted by $R(F \times F)$, to be the \mathbf{Q} -subalgebra of $\text{CH}(F \times F)$ generated by the classes $c_1, c_2, g_1, g_2, \Delta, I$, where Δ and I are the class in $\text{CH}(F \times F)$ of the diagonal Δ_F and the incidence subvariety I respectively.

Proposition 6.3. *For any point $b \in B^\circ$, we have*

$$\text{Im}(\text{CH}(\mathcal{F} \times_B \mathcal{F}) \rightarrow \text{CH}(F_b \times F_b)) = R(F_b \times F_b).$$

Proof. To simplify the notation, let us leave out the subscript b . Thanks to Proposition 6.1, we only need to deal with the following three cases :

- For $\text{Im}(\text{CH}(G \times G) \rightarrow \text{CH}(F \times F))$, it is enough to observe that $\text{CH}(G \times G)$ satisfies the Künneth formula (since the cycle class map $\text{CH}(G \times G) \rightarrow H(G \times G, \mathbf{Q})$ is an isomorphism).
- For $i_* \text{Im}(\text{CH}(J) \rightarrow \text{CH}(I))$, consider

$$\begin{aligned}\tilde{I} &:= \{(x, l, l') \in X \times G \times G \mid x \in l \cap l'\} \text{ and} \\ \tilde{J} &:= \{(x, l, l') \in \mathbf{P}^5 \times G \times G \mid x \in l \cap l'\}\end{aligned}$$

fitting into the diagram

$$\begin{array}{ccccc} \tilde{I} & \xrightarrow{\tau'} & I & \xrightarrow{i} & F \times F \\ \downarrow & \square & \downarrow & \square & \downarrow \\ \tilde{J} & \xrightarrow{\tau} & J & \xrightarrow{j} & G \times G \\ \downarrow \pi & & & & \\ \mathbf{P}^5 & & & & \end{array}$$

Denote by $\widetilde{i} = \tau' \circ i$ and $\widetilde{j} = \tau \circ j$. Then any cycle in J can be written as $\tau_*(\alpha)$ for some $\alpha \in \text{CH}(\widetilde{J})$. Observe that \widetilde{J} is a $\mathbf{P}^4 \times \mathbf{P}^4$ -bundle over \mathbf{P}^5 such that the two relative $\mathcal{O}(1)$ on the fibers are given by $\widetilde{j}^*(g_1)$ and $\widetilde{j}^*(g_2)$, respectively. Therefore α is a linear combination of cycles of the form $\pi^*(h^k)\widetilde{j}^*(g_1^l g_2^m)$ where $k, l, m \in \mathbf{N}$ and $h = \mathcal{O}_{\mathbf{P}^5}(1)$.

$$\begin{aligned} & i_*(\tau_*(\pi^*(h^k)\widetilde{j}^*(g_1^l g_2^m))|_I) \\ &= i_* \circ \tau'_*(\pi^*(h^k)\widetilde{j}^*(g_1^l g_2^m)|_{\widetilde{J}}) \\ &= \widetilde{i}_*(\pi^*(h^k)|_{\widetilde{J}} \cdot \widetilde{i}^*(g_1^l g_2^m)) \\ &= g_1^l g_2^m \cdot i_*(\tau_* \pi^*(h^k)|_I) \\ &= g_1^l g_2^m \cdot \Gamma_{h^k} \end{aligned}$$

where Γ_{h^k} , defined in [35, Appendix A] is the cycle of $F \times F$ represented by the subvariety $\{(l, l') \in F \times F \mid \exists x \in H_1 \cap \dots \cap H_k \text{ such that } x \in l \cap l'\}$, where H_1, \dots, H_k are k general hyperplanes in \mathbf{P}^5 . It is proved in [35, Appendix A] that when $k \geq 1$, Γ_{h^k} is actually a polynomial of c_1, c_2, g_1, g_2 , while $\Gamma_{h^0} = I$.

- For $\Delta_* \text{Im}(\text{CH}(G) \rightarrow \text{CH}(F))$, let us remark that for any $\alpha \in \text{CH}(F)$, we have $\Delta_*(\alpha) = \Delta \cdot \text{pr}_1^*(\alpha)$. Thus it suffices to see that $\text{Im}(\text{CH}(G) \rightarrow \text{CH}(F))$ is generated by g and c .

□

Consequently, in order to prove Theorem 1.11, we need to study the injectivity of the cycle class map restricted to the tautological ring $R(F \times F)$.

Proposition 6.4. *Let X be a smooth cubic fourfold and let F be its Fano variety of lines. Then the cycle class map restricted to the tautological ring $R^*(F \times F)$ is injective.*

Proof. It suffices to show the proposition for general cubic fourfolds, in which case

$$\text{cl} : R(F \times F) \rightarrow \text{Hdg}^{2*}(F \times F)_{\mathbf{Q}}$$

is surjective. Let us show it is injective.

Firstly, it is not hard to count the dimensions of the spaces of Hodge classes :

i	0	1	2	3	4	5	6	7	8
$\dim \text{Hdg}^{2i}$	1	2	6	8	12	8	6	2	1

It is enough to show that the $R^i(F \times F)$ have the same dimensions.

The following relations in $R^*(F \times F)$ are at our disposal.

- (i) $g_1 \cdot \Delta = g_2 \cdot \Delta$; $c_1 \cdot \Delta = c_2 \cdot \Delta$.
- (ii) For $i = 1, 2$, we have $12g_i c_i = 5g_i^3$; $4c_i^2 = g_i^4$.
- (iii) Voisin's relation [41]¹²:

$$I^2 = 2\Delta + I \cdot (g_1^2 + g_1 g_2 + g_2^2) + \Gamma_2(g_1, g_2, c_1, c_2),$$

where Γ_2 is a polynomial of weighted degree 4.

- (iv) In [35, Proposition 17.5], one finds

$$\Delta \cdot I = 6c_1 \Delta - 3g_1^2 \Delta.$$

¹²The coefficients are made precise by [35, Proposition 17.4].

(v) In [35, Lemma 17.6], there is a polynomial P of weighted degree 4 such that

$$c_1 \cdot I = P(g_1, g_2, c_1, c_2);$$

$$c_2 \cdot I = P(g_2, g_1, c_2, c_1).$$

Using these relations, we get easily for each degree a list of generators (as vector-spaces) :

- $R^0 = \langle \mathbb{1} \rangle;$
- $R^1 = \langle g_1, g_2 \rangle;$
- $R^2 = \langle g_1^2, g_1 g_2, g_2^2, c_1, c_2, I \rangle;$
- $R^3 = \langle g_1^3, g_1^2 g_2, g_1 g_2^2, g_2^3, g_1 c_2, g_2 c_1, g_1 I, g_2 I \rangle;$
- $R^4 = \langle g_1^4, g_1^3 g_2, g_1^2 g_2^2, g_1 g_2^3, g_2^4, g_1^2 c_2, g_2^2 c_1, c_1 c_2, g_1^2 I, g_2^2 I, g_1 g_2 I, \Delta \rangle;$
- $R^5 = \langle g_1^4 g_2, g_1^3 g_2^2, g_1^2 g_2^3, g_1 g_2^4, g_1^3 c_2, g_2^3 c_1, g_1^2 g_2 I, g_1 g_2^2 I, g_1 \Delta \rangle;$
- $R^6 = \langle g_1^4 g_2^2, g_1^3 g_2^3, g_1^2 g_2^4, g_1^4 c_2, g_2^4 c_1, g_1^2 g_2^2 I, g_1^2 \Delta \rangle;$
- $R^7 = \langle g_1^4 g_2^3, g_1^3 g_2^4 \rangle;$
- $R^8 = \langle g_1^4 g_2^4 \rangle.$

Observe that we have the same number of generators as the dimension of Hdg^{2i} for $i \neq 5$ or 6. Therefore the cycle class map $R^i(F \times F) \rightarrow H^{2i}(F \times F, \mathbf{Q})$ is injective for $i = 0, 1, 2, 3, 4, 7, 8$.

(vi) As for $i = 5$ (resp. $i = 6$), we use the following (new) tautological relation established in the Appendix Theorem A.1 :

$$6\Delta_*(g) + g_1 g_2 (g_1 + g_2) \cdot I = Q(g_1, g_2, c_1, c_2),$$

where Q is a polynomial.

Therefore the generator $g_1 \Delta = \Delta_*(g)$ (resp. $g_2 \Delta$) is redundant, hence $R^i(F \times F) \rightarrow H^{2i}(F \times F)$ is also injective in these two degrees. \square

Remark 6.5. As a manifestation of the same principle as in §5.3, the extra difficulty encountered here (excess dimension of I , the new tautological relation *etc.*) can be traced back to the lack of positivity of the vector bundle $E = \text{Sym}^3 S^\vee$ on $G = \text{Gr}(\mathbf{P}^1, \mathbf{P}^5)$, namely it satisfies only (\star_1) but not (\star_2) , where S is the tautological subbundle on G .

We can now easily conclude the proof of Theorem 1.11 :

Proof of Theorem 1.11. As the standard conjecture is proved for F_b in [10] (this can also be seen more elementarily by noting that the incidence correspondence I induces an isomorphism from $H^6(F_b, \mathbf{Q})$ to $H^2(F_b, \mathbf{Q})$), numerical equivalence coincides with homological equivalence on powers of F_b . Since any cycle of $\mathcal{F}^\circ \times_{B^\circ} \mathcal{F}^\circ$ is the restriction of a cycle of $\mathcal{F} \times_B \mathcal{F}$, it is enough to show that for any $b \in B^\circ$, the restriction of a cycle $\gamma \in \text{CH}(\mathcal{F} \times_B \mathcal{F})$ to $F_b \times F_b$ is zero if and only if it is homologically trivial, which is proved by combining Proposition 6.3 and Proposition 6.4. \square

With Theorem 1.11 being proved, we proceed to study the zero-cycles and codimension 2 cycles of the LLSvS hyper-Kähler eightfolds. The key input is Voisin's degree 6 dominant rational map [44, Proposition 4.8]

$$F \times F \dashrightarrow Z.$$

Consider the family version of Voisin's construction (over $B^{\circ\circ}$): $\psi : \mathcal{F}^{\circ\circ} \times_{B^{\circ\circ}} \mathcal{F}^{\circ\circ} \dashrightarrow Z$.

Proof of Theorem 1.12. Take a resolution of indeterminacies :

$$\begin{array}{ccc} \mathcal{F}^{\circ\circ} \widetilde{\times}_{B^{\circ\circ}} \mathcal{F}^{\circ\circ} & & \\ \downarrow \tau & \searrow f & \\ \mathcal{F}^{\circ\circ} \times_{B^{\circ\circ}} \mathcal{F}^{\circ\circ} & \xrightarrow[\psi]{} & \mathcal{Z} \end{array}$$

For (i), let $\gamma \in \text{CH}^8(\mathcal{Z})$ be a relative zero-cycle whose degree on fibers is zero. Then, for any $b \in B^{\circ\circ}$,

$$(\tau_* f^*(\gamma))|_{F_b \times F_b} = \tau_{b*} (f^*(\gamma)|_{\widetilde{F_b \times F_b}}) = \tau_{b*} f_b^* (\gamma|_{Z_b}).$$

Thus $\tau_* f^*(\gamma)$ is a relative zero-cycle of fiber degree zero on $\mathcal{F}^{\circ\circ} \times_{B^{\circ\circ}} \mathcal{F}^{\circ\circ}$ and by Theorem 1.11, we know that

$$\tau_{b*} f_b^* (\gamma|_{Z_b}) = 0 \text{ in } \text{CH}^8(F_b \times F_b).$$

For $b \in B^{\circ\circ}$ general, τ_b is birational hence induces an isomorphism on CH_0 , hence $f_b^* (\gamma|_{Z_b}) = 0$. Moreover, since f_b is generically finite of degree 6 (still under the assumption that b is general), we have

$$\gamma|_{Z_b} = \frac{1}{6} f_{b*} f_b^* (\gamma|_{Z_b}) = 0.$$

A specialization argument shows that $\gamma|_{Z_b} = 0$ for all $b \in B^{\circ\circ}$.

For (ii), *i.e.*, codimension 2 cycles: since $H^3(Z_b, \mathbf{Q}) = H^3(F_b \times F_b) = 0$, any cycle in $\text{CH}^2(Z_b)$ or $\text{CH}^2(F_b \times F_b)$ is homologically trivial if and only if its Abel-Jacobi invariant vanishes. Now the same proof as in (i) works because the subspace of Abel-Jacobi kernel for codimension 2 cycles CH_{AJ}^2 , just as CH_0 , is a birational invariant (for smooth projective varieties), hence

$$\tau_{b*} : \text{CH}^2(\widetilde{F_b \times F_b})_{\text{hom}} \rightarrow \text{CH}^2(F_b \times F_b)_{\text{hom}}$$

is an isomorphism. □

Proof of Corollary 1.13. In view of Theorem 1.12, this is just a special case of Proposition 2.5. □

APPENDIX A. ON A NEW TAUTOLOGICAL RELATION

LIE FU, ROBERT LATERVEER, MINGMIN SHEN AND CHARLES VIAL

Let X be a smooth cubic fourfold and F be its Fano variety of lines, which is a hyper-Kähler fourfold by [6]. In this appendix, we establish a new relation (Theorem A.1), up to rational equivalence, among 3-dimensional tautological cycle classes of $F \times F$. Some interesting applications of this tautological relation are also discussed. We try to keep the appendix as self-contained as possible.

Throughout this appendix, let us fix the following notation :

- \mathbf{P}^5 is the ambient space and X is a smooth cubic hypersurface in it.
- $h := c_1(\mathcal{O}_{\mathbf{P}^5}(1))$; $h|_X$ is still denoted by h .
- $G := \text{Gr}(\mathbf{P}^1, \mathbf{P}^5) \simeq \text{Gr}(2, 6)$ is the Grassmannian of projective lines in \mathbf{P}^5 .
- $F := F(X)$ is the Fano variety of lines of X .
- S is the tautological subbundle on G .
- $g := c_1(S^\vee)$ is the Plücker polarization class; $g|_F$ is still denoted by g .
- $c := c_2(S)$; $c|_F$ is still denoted by c .
- $h_i := \text{pr}_i^*(h)$, $g_i := \text{pr}_i^*(g)$ and $c_i := \text{pr}_i^*(c)$ where pr_i is the i -th projection.

- Let $P := \mathbf{P}(S|_F)$ be the incidence variety in $F \times X$. Then the natural projection $p : P \rightarrow F$ is the universal projective line and $q : P \rightarrow X$ is the evaluation map.
- $I \subset F \times F$ is the incidence subvariety parametrizing pairs of intersecting lines contained in X .
- $\tilde{I} := P \times_X P$. Note that I is its image in $F \times F$ via the natural projection.

The main result of this appendix is the following.

Theorem A.1. *There exists a polynomial Q (of weighted degree 5), such that the following equality holds in $\mathrm{CH}^5(F \times F)$:*

$$(12) \quad 6\Delta_*(g) + g_1 g_2 (g_1 + g_2) \cdot I = Q(g_1, g_2, c_1, c_2)$$

where $\Delta : F \hookrightarrow F \times F$ is the diagonal embedding.

Remark A.2. The polynomial Q is not unique. A cohomological computation shows that

$$Q(g_1, g_2, c_1, c_2) = \frac{1}{4}(g_1^4 g_2 + g_1 g_2^4) + \frac{7}{12}(g_1^3 g_2^2 + g_1^2 g_2^3)$$

is one possible choice of Q .

A.1. Proof of the tautological relation. We have the following diagram

$$(13) \quad \begin{array}{ccc} \tilde{I} & \longrightarrow & X \\ \downarrow i & & \downarrow \Delta_X \\ P \times P & \xrightarrow{(q,q)} & X \times X \\ \downarrow (p,p) & & \\ F \times F & & \end{array}$$

Let us first introduce some natural cycles on $F \times F$: for any $i \in \mathbf{N}$, define

$$\Gamma_{h^i} := (p, p)_*(q, q)^*(\Delta_{X*}(h^i)) \in \mathrm{CH}^{i+2}(F \times F).$$

Note that Γ_{h^0} is nothing but the incidence correspondence I . Geometrically, Γ_{h^i} is represented by the locus of pairs of lines contained in X intersecting at a point which lies on the intersection of i general hyperplane sections of X .

Lemma A.3. *For any $i > 0$, the cycle Γ_{h^i} is a polynomial of g_1, g_2, c_1, c_2 . Precisely,*

$$\begin{aligned} \Gamma_h &= \frac{1}{18}(g_1^3 + 6g_1^2 g_2 + 6g_1 g_2^2 + g_2^3 - 6g_1 c_2 - 6g_2 c_1); \\ \Gamma_{h^2} &= \frac{1}{18}(g_1^3 g_2 + 6g_1^2 g_2^2 + g_1 g_2^3 - 6g_1^2 c_2 - 6g_2^2 c_1 + 6c_1 c_2); \\ \Gamma_{h^3} &= \frac{1}{18}(g_1^3 g_2^2 + g_1^2 g_2^3 - g_1^3 c_2 - g_2^3 c_1); \\ \Gamma_{h^4} &= \frac{1}{108} g_1^3 g_2^3. \end{aligned}$$

Proof. A slightly more complicated (but equivalent) form of the first two formulas is proved in [35, Proposition A.6]. For the convenience of the reader, we give a complete proof here. The excess

intersection formula [18, §6.3] applied to the following cartesian diagram

$$\begin{array}{ccc} X & \longrightarrow & \mathbf{P}^5 \\ \downarrow \Delta_X & \square & \downarrow \Delta_P \\ X \times X & \longrightarrow & \mathbf{P}^5 \times \mathbf{P}^5 \end{array}$$

yields that for any $i \in \mathbf{N}$, we have in $\mathrm{CH}(X \times X)$

$$3\Delta_{X^*}(h^{i+1}) = \Delta_{P^*}(h^i)|_{X \times X}.$$

From $\Delta_{P^*}(h^i) = h_1^5 h_2^i + \cdots + h_1^i h_2^5$, we obtain

$$\Delta_{X^*}(h^i) = \frac{1}{3} (h_1^4 h_2^i + \cdots + h_1^i h_2^4).$$

Therefore

$$\begin{aligned} \Gamma_{hi} &= (p, p)_*(q, q)^*(\Delta_{X^*}(h^i)) \\ &= \frac{1}{3} (p, p)_*(q, q)^*(h_1^4 h_2^i + \cdots + h_1^i h_2^4) \\ &= \frac{1}{3} (f_4 \times f_i + \cdots + f_i \times f_4) \end{aligned}$$

where $f_j := p_* q^*(h^j)$ and \times is the exterior product $\mathrm{pr}_1^*(-) \cdot \mathrm{pr}_2^*(-)$. All the formulas in the statement then follow from the facts that $f_1 = 1$, $f_2 = g$, $f_3 = g^2 - c$, $f_4 = \frac{1}{6}g^3$ (cf. [35, Lemma A.4] and [41, Lemma 3.2] and [41, Lemma 3.5]). \square

Define $I_0 := I \setminus \Delta_F$ the subvariety of $F \times F$ parametrizing pairs of distinct intersecting lines in X . We then have a natural morphism

$$q_0 : I_0 \rightarrow X$$

which sends two lines to their intersection point.

Lemma A.4. *The inclusion $I_0 \hookrightarrow F \times F \setminus \Delta_F$ is a local complete intersection and the Chern classes of the normal bundle $N := N_{I_0/F \times F \setminus \Delta_F}$ are given by*

$$\begin{aligned} c_1(N) &= (g_1 + g_2)|_{I_0} - q_0^*(h); \\ c_2(N) &= (g_1^2 + g_1 g_2 + g_2^2)|_{I_0} - 3(g_1 + g_2)|_{I_0} \cdot q_0^*(h) + 6q_0^*(h^2). \end{aligned}$$

Proof. Note that $\tilde{I} \subset P \times P$ is a local complete intersection (since $\tilde{I} \subset P \times P$ is obtained from the local complete intersection $\Delta_X \subset X \times X$ via base change) and that $\tilde{I} \setminus \Delta_P \subset P \times P$ is a section of $P \times P \rightarrow F \times F$ over I_0 . We apply [18, B.7.5] and see that $I_0 \subset F \times F \setminus \Delta_F$ is a local complete intersection. Using the section $\tilde{I} \setminus \Delta_P$, we view I_0 as a subvariety of $P \times P$. Then we get the following short exact sequence

$$0 \longrightarrow \mathrm{pr}_1^* T_{P/F} \oplus \mathrm{pr}_2^* T_{P/F} \longrightarrow N_{I_0/P \times P} \longrightarrow N_{I_0/F \times F} \longrightarrow 0$$

Note that by construction, we have

$$N_{I_0/P \times P} = q_0^* T_X.$$

The Chern classes of $N_{I_0/F \times F}$ are computed as follows

$$\begin{aligned} c(N) &= \frac{q_0^* c(T_X)}{pr_1^* c(T_{P/F}) \cdot pr_2^* c(T_{P/F})} \\ &= \frac{(1+h)^6}{(1+3h)(1+2q_0^* h - g_1|_{I_0})(1+2q_0^* h - g_2|_{I_0})} \end{aligned}$$

The lemma follows from the expansion of the above equation. \square

Remark A.5. The previous lemma implies that

$$I^2|_{F \times F \setminus \Delta_F} = I \cdot (g_1^2 + g_1 g_2 + g_2^2) - 3(g_1 + g_2)\Gamma_h + 6\Gamma_{h^2}.$$

Thus by Lemma A.3 there exists $\alpha \in \mathbf{Q}$ and a polynomial Γ_2 , such that in $\text{CH}^4(F \times F)$ we have

$$I^2 = \alpha \cdot \Delta_F + I \cdot (g_1^2 + g_1 g_2 + g_2^2) + \Gamma_2(g_1, g_2, c_1, c_2),$$

for some $\alpha \in \mathbf{Q}$. This was proven by Voisin [41]. In fact, $\alpha = 2$, as is computed in [35, Proposition 17.4].

Proof of Theorem A.1. Let us first prove the theorem for a general cubic fourfold X . Fix three general hyperplane sections H_1, H_2, H_3 of X . For $i = 1, 2, 3$, let

$$Z_i := \{(l, l') \in F \times F \mid l \cap l' \cap H_1 \cap \cdots \cap H_i \neq \emptyset\}.$$

On the one hand, as is mentioned before, the class of Z_i in $\text{CH}^{2+i}(F \times F)$ is equal to Γ_{h^i} ; on the other hand, denoting by $Z_i^o := Z_i \setminus \Delta_F$ the complement of the diagonal in Z_i , the class of Z_i^o in $\text{CH}^i(I_0)$ is equal to $q_0^*(h^i)$ by definition. We have thus the diagram

$$\begin{array}{ccc} Z_3^o \subset Z_2^o \subset Z_1^o & \xrightarrow{\iota} & I_0 \xrightarrow{\iota} F \times F \setminus \Delta_F \\ & & \downarrow q_0 \\ & & X \end{array}$$

Denoting by N the normal bundle of ι , we obtain

$$\begin{aligned} I \cdot \Gamma_h|_{F \times F \setminus \Delta_F} &= I_0 \cdot \iota_* q_0^*(h) \\ &= \iota_* (q_0^*(h) \cdot c_2(N)) \\ &= \iota_* \left((g_1^2 + g_1 g_2 + g_2^2)|_{I_0} \cdot q_0^*(h) - 3(g_1 + g_2)|_{I_0} \cdot q_0^*(h^2) + 6q_0^*(h^3) \right) \\ &= \left((g_1^2 + g_1 g_2 + g_2^2) \cdot Z_1 - 3(g_1 + g_2) \cdot Z_2 + 6Z_3 \right)|_{F \times F \setminus \Delta_F} \\ &= \left((g_1^2 + g_1 g_2 + g_2^2) \cdot \Gamma_h - 3(g_1 + g_2) \cdot \Gamma_{h^2} + 6\Gamma_{h^3} \right)|_{F \times F \setminus \Delta_F} \end{aligned}$$

where the third equality uses Lemma A.4. By Lemma A.3, there exists a polynomial P_1 such that

$$I \cdot \Gamma_h|_{F \times F \setminus \Delta_F} = P_1(g_1, g_2, c_1, c_2)|_{F \times F \setminus \Delta_F},$$

Here more precisely, one can compute by Lemma A.3 and the relation $12gc = 5g^3$ that

$$P_1(g_1, g_2, c_1, c_2) = \frac{5}{12} (g_1^4 g_2 + 4g_1^3 g_2^2 + 4g_1^2 g_2^3 + g_1 g_2^4 - 3g_1^3 c_2 - 3g_2^3 c_1).$$

By the localization short exact sequence of Chow groups, there exists an element $D \in \text{CH}^1(F)$, such that in $\text{CH}^5(F \times F)$ we have

$$I \cdot \Gamma_h + \Delta_*(D) = P_1(g_1, g_2, c_1, c_2).$$

Since X is assumed (for now) to be general, $\mathrm{CH}^1(F)$ is generated by g , hence $D = \lambda g$ for some $\lambda \in \mathbf{Q}$. It yields that in $\mathrm{CH}^5(F \times F)$,

$$(14) \quad I \cdot \Gamma_h + \lambda \Delta_*(g) = P_1(g_1, g_2, c_1, c_2).$$

However, we know that $I \cdot c_1, I \cdot c_2, I \cdot g_1^3$ and $I \cdot g_2^3$ are polynomials of g_1, g_2, c_1, c_2 by [35, Lemma 17.6] (cf. the known relations collected in the proof of Proposition 6.4). The first formula in Lemma A.3 then yields that

$$(15) \quad I \cdot \Gamma_h = \frac{1}{3} I \cdot (g_1^2 g_2 + g_1 g_2^2) + P_2(g_1, g_2, c_1, c_2)$$

for some polynomial P_2 .

Putting (14) and (15) together, we know that there exists a polynomial Q such that the following equality holds in $\mathrm{CH}^5(F \times F)$:

$$3\lambda \cdot \Delta_*(g) + I \cdot (g_1^2 g_2 + g_1 g_2^2) = Q(g_1, g_2, c_1, c_2).$$

By considering the action of both sides on the cohomology, we easily see that $\lambda = 2$ and that

$$Q(g_1, g_2, c_1, c_2) = \frac{1}{4}(g_1^4 g_2 + g_1 g_2^4) + \frac{7}{12}(g_1^3 g_2^2 + g_1^2 g_2^3).$$

Therefore the desired relation is proven for a general cubic fourfold. As all the cycles appearing are universally defined in the universal Fano variety of lines, a specialization argument shows that this relation must also hold for any smooth cubic fourfold. \square

A.2. Some applications to the Fourier decomposition of F . Our aim is to use Theorem 1.11, which is based on Theorem A.1, to complement the results of [35] concerning the multiplicative structure of the Chow motive of the Fano variety of lines on a smooth cubic fourfold.

A.2.1. An explicit Chow–Künneth decomposition for F . Recall that a Chow–Künneth decomposition for a smooth projective variety X of dimension d is a decomposition of the diagonal $\Delta_X \in \mathrm{CH}^d(X \times X)$ into a sum $\Delta_X = \pi_X^0 + \cdots + \pi_X^{2d}$ of mutually orthogonal idempotent correspondences $\pi_X^i \in \mathrm{CH}^d(X \times X)$ whose action in cohomology is given by $(\pi_X^i)_* H^*(X, \mathbf{Q}) = H^i(X, \mathbf{Q})$. It is a conjecture of Murre that all smooth projective varieties should admit a Chow–Künneth decomposition. In [35], it is shown that the Fano variety of lines on a smooth cubic fourfold admits a Chow–Künneth decomposition; see especially [35, Theorem 3.3]. Such a decomposition is obtained by modifying the following correspondences in $\mathrm{CH}^4(F \times F)$:

$$(16) \quad \pi_F^0 = \frac{1}{23 \cdot 25} l_1^2, \quad \pi_F^2 = \frac{1}{25} L \cdot l_1, \quad \pi_F^4 = \frac{1}{2} (L^2 - \frac{1}{25} l_1 \cdot l_2), \quad \pi_F^6 = \frac{1}{25} L \cdot l_2, \quad \pi_F^8 = \frac{1}{23 \cdot 25} l_2^2.$$

Here, $L := \frac{1}{3}(g_1^2 + \frac{3}{2}g_1 g_2 + g_2^2 - c_1 - c_2) - I \in \mathrm{CH}^2(F \times F)$ is a, and in fact “the” by Proposition 6.4, tautological cycle representing the Beauville–Bogomolov form; see [35, Proposition 19.1]. The cycle $l \in \mathrm{CH}^2(F)$ is the restriction of L to the diagonal, and, as before, a subscript i indicates the pull-back along the projection $F \times F \rightarrow F$ to the i -th factor.

As was expected from [35, Conjecture 3], these correspondences already define a Chow–Künneth decomposition:

Proposition A.6. *The correspondences in (16) define a Chow–Künneth decomposition of F .*

Proof. The correspondences π_F^{2i} of (16) are cycles on $F \times F$ that belong to the image of the restriction map $\text{CH}(\mathcal{F} \times_B \mathcal{F}) \rightarrow \text{CH}(F \times F)$, and they define a Künneth decomposition of the diagonal in cohomology by [35, Corollary 1.7]. (Here $\mathcal{F} \rightarrow B$ is the universal Fano variety of lines as defined in §3). It follows readily from Theorem 1.11 that they define a Chow–Künneth decomposition. \square

A.2.2. *A new multiplicativity statement.* Using the Chow–Künneth decomposition (16) given by Proposition A.6, we can define, for all integers i and j ,

$$\text{CH}^i(F)_{(j)} := (\pi_F^{2i-j})_* \text{CH}^i(F).$$

Concretely, we have (cf. [35])

$$\begin{aligned} \text{CH}^4(F) &= \text{CH}^4(F)_{(0)} \oplus \text{CH}^4(F)_{(2)} \oplus \text{CH}^4(F)_{(4)} \\ \text{CH}^3(F) &= \text{CH}^3(F)_{(0)} \oplus \text{CH}^3(F)_{(2)} \\ \text{CH}^2(F) &= \text{CH}^2(F)_{(0)} \oplus \text{CH}^2(F)_{(2)} \\ \text{CH}^1(F) &= \text{CH}^1(F)_{(0)} \\ \text{CH}^0(F) &= \text{CH}^0(F)_{(0)}. \end{aligned}$$

In [35], it was proved that, for the Fano variety of lines on a *very general* cubic fourfold, the decomposition $\text{CH}^i(F)_{(j)}$ defines a bigrading on the Chow ring $\text{CH}(F)$, in the sense that for all integers i, i', j, j' we have

$$\text{CH}^i(F)_{(j)} \cdot \text{CH}^{i'}(F)_{(j')} \subseteq \text{CH}^{i+i'}(F)_{(j+j')}.$$

In the case of the Fano variety of lines on a non-very general cubic fourfold, the following two relations could not be established (see [35, Remark 22.9]):

$$(17) \quad \text{CH}^1(F) \cdot \text{CH}^2(F)_{(0)} \subseteq \text{CH}^3(F)_{(0)};$$

$$(18) \quad \text{CH}^2(F)_{(0)} \cdot \text{CH}^2(F)_{(0)} \subseteq \text{CH}^4(F)_{(0)} = \mathbf{Q} \cdot \nu_F.$$

Using the new relation (12), we can now prove one of the missing two inclusions:

Proposition A.7. *Let F be the Fano variety of lines on a smooth cubic fourfold. Then*

$$\text{CH}^1(F) \cdot \text{CH}^2(F)_{(0)} = \text{CH}^3(F)_{(0)}.$$

Proof. We first show that $\text{CH}^3(F)_{(0)} \subseteq \text{CH}^1(F) \cdot \text{CH}^2(F)_{(0)}$. On the one hand, the cycle class map $\text{CH}^3(F)_{(0)} \rightarrow H_{alg}^6(F, \mathbf{Q})$ is an isomorphism; on the other hand, the hard Lefschetz isomorphism implies that $H_{alg}^6(F, \mathbf{Q})$ is generated by $g^2 \cdot H_{alg}^2(F, \mathbf{Q}) = g^2 \cdot \text{CH}^1(F)$. Hence $\text{CH}^3(F)_{(0)}$ is generated by intersections of three divisors, which is contained in $\text{CH}^1(F) \cdot \text{CH}^2(F)_{(0)}$ since we know $\text{CH}^1(F) \cdot \text{CH}^1(F) \subset \text{CH}^2(F)_{(0)}$.

For the inverse inclusion, which is (17), by [35, Proposition 22.7], we only need to show that if α is a cycle in $\text{CH}^2(F)_{(0)}$, then $g \cdot \alpha$ belongs to $\text{CH}^3(F)_{(0)}$. To this end, we consider the correspondence

$$\Gamma := \pi_F^4 \circ \Gamma_\iota \circ {}^t\Gamma_\iota \circ \pi_F^4 \in \text{CH}^5(F \times F),$$

where $\iota: H \hookrightarrow F$ denotes the inclusion of a hyperplane with respect to the Plücker embedding. Clearly, Γ is homologically trivial. But Γ is universally defined, and so Theorem 1.11 implies that Γ is rationally trivial. The action of Γ on $\text{CH}^2(F)_{(0)}$ is the same as

$$\text{CH}^2(F)_{(0)} \xrightarrow{g} \text{CH}^3(F) \rightarrow \text{CH}^3(F)_{(2)}$$

(where the second arrow is projection on a direct summand), and so we are done. \square

With notations as in §6, it seems that the final missing inclusion (18) cannot be obtained from considering the subring $\text{Im}(\text{CH}(\mathcal{F} \times_B \mathcal{F}) \rightarrow \text{CH}(F_b \times F_b))$. Rather, a streamlined proof of all inclusions $\text{CH}^i(F)_{(j)} \cdot \text{CH}^j(F)_{(j')} \subseteq \text{CH}^{i+j}(F)_{(j+j')}$ would follow from establishing that the Chow–Künneth decomposition (16) is *multiplicative* in the sense of [35, §8], meaning that

$$\pi_F^k \circ \delta_F \circ (\pi_F^i \otimes \pi_F^j) = 0 \quad \text{in } \text{CH}^8(F \times F \times F), \quad \text{for all } k \neq i + j,$$

where δ_F denotes the class of the small diagonal in $F \times F \times F$ viewed as a correspondence from $F \times F$ to F . This in turn would follow from showing the generalized Franchetta conjecture for the relative cube of the universal Fano variety of lines, *i.e.* one would need to show that the subring

$$\text{Im}(\text{CH}(\mathcal{F} \times_B \mathcal{F} \times_B \mathcal{F}) \rightarrow \text{CH}(F_b \times F_b \times F_b))$$

injects into cohomology via the cycle class map for all b . An approach would consist in first showing that this subring consists of “tautological cycles” and then in establishing enough “tautological relations”, as was done in Propositions 6.3 and 6.4 in the case of the relative square.

REFERENCES

1. Nicolas Addington and Manfred Lehn, *On the symplectic eightfold associated to a pfaffian cubic fourfold*, J. reine angew. Math., to appear (2015).
2. Enrico Arbarello and Maurizio Cornalba, *The Picard groups of the moduli spaces of curves*, Topology **26** (1987), no. 2, 153–171. MR 895568
3. Arnaud Beauville, *Some remarks on Kähler manifolds with $c_1 = 0$* , in: Classification of algebraic and analytic manifolds (Katata, 1982), Birkhäuser Boston, Boston 1983,
4. ———, *Fano threefolds and K3 surfaces*, in: The Fano conference, University Torino, Turin 2004,
5. ———, *On the splitting of the Bloch-Beilinson filtration*, Algebraic cycles and motives. Vol. 2, London Math. Soc. Lecture Note Ser., vol. 344, Cambridge Univ. Press, Cambridge, 2007, pp. 38–53. MR 2187148 (2009c:14007)
6. Arnaud Beauville and Ron Donagi, *La variété des droites d’une hypersurface cubique de dimension 4*, C. R. Acad. Sci. Paris Sér. I Math. **301** (1985), no. 14, 703–706. MR 818549 (87c:14047)
7. Arnaud Beauville and Claire Voisin, *On the Chow ring of a K3 surface*, J. Algebraic Geom. **13** (2004), no. 3, 417–426. MR 2047674 (2005b:14011)
8. Nicolas Bergeron and Zhiyuan Li, *Tautological classes on moduli space of hyperkähler manifolds*, Preprint, arXiv:1703.04733 (2017).
9. Spencer Bloch and Vasudevan Srinivas, *Remarks on correspondences and algebraic cycles*, Amer. J. Math. **105** (1983), no. 5, 1235–1253. MR 714776 (85i:14002)
10. François Charles and Eyal Markman, *The standard conjectures for holomorphic symplectic varieties deformation equivalent to Hilbert schemes of K3 surfaces*, Compos. Math. **149** (2013), no. 3, 481–494. MR 3040747
11. Mark Andrea de Cataldo and Luca Migliorini, *The Chow groups and the motive of the Hilbert scheme of points on a surface*, J. Algebra **251** (2002), no. 2, 824–848. MR 1919155
12. Igor Dolgachev, *Weighted projective varieties*, Group actions and vector fields, Lecture Notes in Math. **956** (1982) 34–71. MR 0704986
13. A. Franchetta, *Sulle serie lineari razionalmente determinate sulla curva a moduli generali di dato genere*, Matematiche, Catania **9** (1954), 126–147. MR 0069538
14. Lie Fu, *Decomposition of small diagonals and Chow rings of hypersurfaces and Calabi-Yau complete intersections*, Adv. Math. **244** (2013), 894–924. MR 3077892
15. ———, *Beauville-Voisin conjecture for generalized Kummer varieties*, Int. Math. Res. Not. IMRN (2015), no. 12, 3878–3898. MR 3356741
16. ———, *On the action of symplectic automorphisms on the CH_0 -groups of some hyper-Kähler fourfolds*, Math. Z. **280** (2015), no. 1-2, 307–334. MR 3343908
17. Lie Fu and Zhiyu Tian, *Motivic hyperkähler resolution conjecture: II. Hilbert schemes of K3 surfaces*, preprint (2017).
18. William Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. no. 2, Springer-Verlag (1998). MR 1644323
19. John Harer, *The second homology group of the mapping class group of an orientable surface*, Invent. Math. **72** (1983), no. 2, 221–239. MR 700769
20. Atanas Iliev and Laurent Manivel, *Prime Fano threefolds and integrable systems*, Math. Annalen **339** no. 4 (2007), 937–955.

21. Vasilii Iskovskikh and Yuri Prokhorov, *Fano varieties*, in Algebraic Geometry V (A. N. Parsin, I. R. Shafarevich, Eds.), Encyclopaedia of Math. Sci. 47, Springer 1998.
22. Martí Lahoz, Manfred Lehn, Emanuele Macrì, and Paolo Stellari, *Generalized twisted cubics on a cubic fourfold as a moduli space of stable objects*, J. Math. Pures Appl, to appear (2017).
23. Robert Laterveer, *Bloch's conjecture for certain hyperkähler fourfolds*, preprint 2016.
24. Christian Lehn, Manfred Lehn, Christoph Sorger, and Duco van Straten, *Twisted cubics on cubic fourfolds*, preprint, arXiv:1305.0178 (2013).
25. Nicole Mestrano, *Conjecture de Franchetta forte*, Invent. Math. **87** (1987), no. 2, 365–376. MR 870734
26. Shigeru Mukai, *Curves, K3 surfaces and Fano 3-folds of genus = 10*. In Algebraic Geometry and Commutative Algebra, edited by H. Hijikata, H. Hironaka, M. Maruyama, H. Matsumura, M. Miyanishi, T. Oda and K. Ueno, Vol. I, 357–77. (1988).
27. ———, *Polarized K3 surfaces of genus 18 and 20*. In Complex Projective Geometry (Trieste, 1989/Bergen, 1989), edited by G. Ellingsrud, C. Peskine, G. Sacchiero and S. A. Strømme, 264–276. London Mathematical Society, Lecture Note Series 179. Cambridge: Cambridge University Press, 1992.
28. ———, *Polarized K3 surfaces of genus thirteen*. In Moduli Spaces and Arithmetic Geometry, edited by S. Mukai, Y. Miyaoka, S. Mori, A. Moriwaki and I. Nakamura, 315–326. Advanced Studies in Pure Mathematics 45. Tokyo: The Mathematical Society of Japan, 2006.
29. ———, *K3 surfaces of genus sixteen*. Preprint, 2012, available at <http://www.kurims.kyoto-u.ac.jp/preprint/file/RIMS1743.pdf> (accessed July 4, 2016).
30. Jaap Murre, Jan Nagel and Chris Peters, *Lectures on the theory of pure motives*, Amer. Math. Soc. University Lecture Series 61, Providence 2013.
31. Kieran G. O'Grady, *Double covers of EPW–sextics*, Michigan Math. J. 62 (2013), 143–184.
32. Kieran G. O'Grady, *Moduli of sheaves and the Chow group of K3 surfaces*, J. Math. Pures Appl. (9) **100** (2013), no. 5, 701–718. MR 3115830
33. Nebojsa Pavic, Junliang Shen, and Qizheng Yin, *On O'Grady's generalized Franchetta conjecture*, International Mathematics Research Notices, to appear (2016).
34. Ulrike Rieß, *On Beauville's conjectural weak splitting property*, Int. Math. Res. Not. IMRN (2016), no. 20, 6133–6150. MR 3579961
35. Mingmin Shen and Charles Vial, *The Fourier transform for certain hyperkähler fourfolds*, Mem. Amer. Math. Soc. **240** (2016), no. 1139, vii+163. MR 3460114
36. ———, *On the motive of the Hilbert cube $X^{[3]}$* , Forum Math. Sigma **4** (2016).
37. Evgeny Shinder and Andrey Soldatenkov, *On the geometry of the Lehn–Lehn–Sorger–van Straten eightfold*, Kyoto Journal of Mathematics, (2017).
38. Charles Vial, *Algebraic cycles and fibrations*, Doc. Math. **18** (2013), 1521–1553. MR 3158241
39. ———, *On the motive of some hyperkähler varieties*, J. für Reine u. Angew. Math. 725 (2017), 235–247,
40. Claire Voisin, *Hodge theory and complex algebraic geometry. II*, Cambridge Studies in Advanced Mathematics, vol. 77, Cambridge University Press, Cambridge, 2003, Translated from the French by Leila Schneps. MR 1997577 (2005c:32024b)
41. ———, *On the Chow ring of certain algebraic hyper-Kähler manifolds*, Pure Appl. Math. Q. **4** (2008), no. 3, part 2, 613–649. MR 2435839 (2009m:14004)
42. ———, *The generalized Hodge and Bloch conjectures are equivalent for general complete intersections*, Ann. Sci. Éc. Norm. Supér. (4) **46** (2013), no. 3, 449–475 (2013). MR 3099982
43. ———, *Chow rings, decomposition of the diagonal, and the topology of families*, Annals of Mathematics Studies, vol. 187, Princeton University Press, Princeton, NJ, 2014. MR 3186044
44. ———, *Remarks and questions on coisotropic subvarieties and 0-cycles of hyper-Kähler varieties*, K3 surfaces and their moduli, Progr. Math., vol. 315, Birkhäuser/Springer, [Cham], 2016, pp. 365–399. MR 3524175
45. Qizheng Yin, *Finite-dimensionality and cycles on powers of K3 surfaces*, Comment. Math. Helv. **90** (2015), no. 2, 503–511. MR 3351754

INSTITUT CAMILLE JORDAN, UNIVERSITÉ CLAUDE BERNARD LYON 1, FRANCE

E-mail address: `fu@math.univ-lyon1.fr`

INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE, CNRS, UNIVERSITÉ DE STRASBOURG, FRANCE

E-mail address: `robert.laterveer@math.unistra.fr`

UNIVERSITÄT BIELEFELD, GERMANY

E-mail address: `vial@math.uni-bielefeld.de`

KdV INSTITUTE FOR MATHEMATICS, UNIVERSITY OF AMSTERDAM, NETHERLANDS

E-mail address: `M.Shen@uva.nl`