

Masters course: Representations of Algebras 2

I want to cover a number of key topics in the representation theory of finite-dimensional associative algebras. Specifically:

- Correspondences given by faithfully balanced modules, and applications to Auslander algebras and homological conjectures. (Originally planned for the previous semester, but carried over, since there was not enough time.)
- Tilting and tau-tilting theory, including equivalences of derived categories.
- Geometric methods for studying representations of algebras, including relevant facts about varieties and schemes without proofs. (There will be less time for this than originally planned.)
- If time, possibly preprojective algebras and Kleinian singularities.

Some relevant books:

- I. Assem and F. U. Coelho, Basic representation theory of algebras, Springer 2020.
- I. Assem, D. Simson and A. Skowroński, Elements of the representation theory of associative algebras. Volume 1, Techniques of representation theory, CUP 2006.
- H. Derksen and J. Weyman, An introduction to quiver representations, American Mathematical Society 2017.
- P. Gabriel and A. V. Roiter, Representations of finite dimensional algebras, Springer 1977.
- A. Kirillov Jr., Quiver Representations and Quiver Varieties, American Mathematical Society 2016.
- A. Skowroński and K. Yamagata, Frobenius algebras 2 Tilted and Hochschild extension algebras, European Mathematical Society 2017.

The section numbering continues from the previous lecture course.

Contents

4	Homological topics	1
4.1	Higher generation and cogeneration	1
4.2	Faithfully balanced modules and endomorphism correspondence . .	4
4.3	Dominant dimension and Auslander correspondence	7
4.4	Homological conjectures for f.d. algebras	13
4.5	No loops conjecture	17
5	Tilting theory	19
5.1	Torsion theories and tau-rigid modules	20
5.2	Tilting modules	26
5.3	The Brenner-Butler Theorem	28
5.4	Derived equivalences	33
5.5	Consequences for Grothendieck groups	35
5.6	Some tau-tilting theory	37
6	Varieties and schemes of algebras and modules	42
6.1	Varieties of algebras	42
6.2	Schemes and varieties of modules	49
	More coming...	

4 Homological topics

In this section I want to discuss

- Some of the many homological conjectures for f.d. algebras
- Auslander's correspondence between algebras A of finite representation type and algebras B with $\text{gl. dim } B \leq 2 \leq \text{dom. dim } B$, and Iyama's generalization of this with cluster tilting objects.

The unifying feature is what I call endomorphism correspondence for faithfully balanced modules.

4.1 Higher generation and cogeneration

We are interested in finite-dimensional algebras A over a field K (but most things generalize easily to Artin algebras).

Except where explicitly stated, all modules are f.d., and we write $A\text{-mod}$ for the category of finite-dimensional left A -modules.

We write D for the duality $\text{Hom}_K(-, K)$ between $A\text{-mod}$ and $A^{op}\text{-mod}$.

Recall that a *module class* in $A\text{-mod}$ is a full subcategory closed under isomorphisms, direct sums and direct summands. Given any module M , $\text{add}(M)$ is the smallest module class containing M . It is given by the modules isomorphic to a direct summand of M^n for some n .

Definition. Given a module M , $\text{gen}(M)$ denotes the module class consisting of quotients of direct sums of copies of M and $\text{cogen}(M)$ the module class of submodules of a direct sum of copies of M .

We say M is a *generator* if $\text{gen}(M) = A\text{-mod}$. It is equivalent that $A \in \text{gen}(A)$, or that $A \in \text{add}(M)$. We say M is a *cogenerator* if $\text{cogen}(M) = A\text{-mod}$. It is equivalent that $DA \in \text{cogen}(M)$, or $DA \in \text{add}(M)$.

There are higher versions as follows. Here

Proposition (1). *Let M be an A -module and $n \geq 0$ and consider M also as a B -module, where $B = \text{End}_A(M)$. For an A -module X , the following are equivalent.*

(a) *There is an exact sequence*

$$M_n \xrightarrow{f_n} M_{n-1} \rightarrow \cdots \rightarrow M_0 \xrightarrow{f_0} X \rightarrow 0$$

with $M_i \in \text{add } M$, such that the sequence

$$\text{Hom}_A(M, M_{n-1}) \rightarrow \cdots \rightarrow \text{Hom}_A(M, M_0) \rightarrow \text{Hom}_A(M, X) \rightarrow 0$$

is exact (note that this is automatic if M is projective).

(b) *The natural map $\text{Hom}_A(M, X) \otimes_B M \rightarrow X$ is surjective (in case $n = 0$) or an isomorphism (in case $n > 0$) and $\text{Tor}_i^B(\text{Hom}_A(M, X), M) = 0$ for $0 < i < n$.*

Definition. We define $\text{gen}_n(M)$ to be the full subcategory of $A\text{-mod}$ given by the modules X satisfying these conditions. Using condition (b) it is easy to see that it is a module class. Clearly

$$\text{add}(M) \subseteq \cdots \subseteq \text{gen}_2(M) \subseteq \text{gen}_1(M) \subseteq \text{gen}_0(M) = \text{gen}(M).$$

Proof. (a) $\Rightarrow$ (b). First note that we may assume that the sequence

$$\text{Hom}_A(M, M_n) \rightarrow \text{Hom}_A(M, M_{n-1}) \rightarrow \cdots \rightarrow \text{Hom}_A(M, M_0) \rightarrow \text{Hom}_A(M, X) \rightarrow 0$$

is exact. By assumption it is exact except possibly at $\text{Hom}_A(M, M_{n-1})$. Recall from section 1.9, that $\text{add } M$ is functorially finite in $A\text{-mod}$. Thus the module $\text{Im}(f_n)$ has a right $\text{add } M$ -approximation, say $f' : M' \rightarrow \text{Im}(f_n)$. Since it is an approximation, we can factorize $f_n = f'g$ for some $g : M_n \rightarrow M'$. Thus the map f' has image $\text{Im}(f_n)$. Thus the sequence

$$M' \xrightarrow{f'} M_{n-1} \xrightarrow{f_{n-1}} \cdots \xrightarrow{f_1} M_0 \xrightarrow{f_0} X \rightarrow 0$$

is exact. Also, the sequence

$$\text{Hom}_A(M, M') \rightarrow \text{Hom}_A(M, M_{n-1}) \rightarrow \cdots \rightarrow \text{Hom}_A(M, M_0) \rightarrow \text{Hom}_A(M, X) \rightarrow 0$$

is exact, since any morphism in $\text{Hom}(M, M_{n-1})$ which is sent to zero in $\text{Hom}(M, M_{n-2})$ has image contained in $\text{Ker}(f_{n-1}) = \text{Im}(f_n)$, and hence factors through the approximation f' . Thus replacing M_n by M' and f_n by f' if necessary, we have the claimed exactness.

Now we have a commutative diagram

$$\begin{array}{ccccccc} \text{Hom}(M, M_n) \otimes M & \longrightarrow & \cdots & \longrightarrow & \text{Hom}(M, M_0) \otimes M & \longrightarrow & \text{Hom}(M, X) \otimes M \longrightarrow 0 \\ \phi_n \downarrow & & & & \phi_0 \downarrow & & \theta \downarrow \\ M_n & \xrightarrow{f_n} & \cdots & \xrightarrow{f_1} & M_0 & \xrightarrow{f_0} & X \longrightarrow 0 \end{array}$$

For any $M' \in \text{add } M$, the natural map $\text{Hom}(M, M') \otimes_B M \rightarrow M'$ is an isomorphism, since it is for $M' = M$. Thus the ϕ_i are isomorphisms.

Since ϕ_0 and f_0 are surjective, so is θ . If $n > 0$, then since tensor products are right exact, the part of the diagram below and to the right of $\text{Hom}(M, M_1) \otimes M$ has exact rows, so implies that θ is an isomorphism.

Since $M_i \in \text{add}(M)$, as a right B -module, we have

$$\text{Hom}_A(M, M_i) \in \text{add}(\text{Hom}_A(M, M)) = \text{add}(B_B),$$

so the exact sequence

$$\text{Hom}_A(M, M_n) \rightarrow \text{Hom}_A(M, M_{n-1}) \rightarrow \cdots \rightarrow \text{Hom}_A(M, M_0) \rightarrow \text{Hom}_A(M, X) \rightarrow 0$$

is part of a projective resolution of $\text{Hom}_A(M, X)$ as a right B -module. We can use it to compute $\text{Tor}_i^B(\text{Hom}_A(M, X), M)$ for $i < n$ as the homology of the complex

$$\text{Hom}(M, M_n) \otimes_B M \rightarrow \cdots \rightarrow \text{Hom}(M, M_0) \otimes_B M \rightarrow 0.$$

But by the commutative diagram above, this is isomorphic to the complex

$$M_n \rightarrow \cdots \rightarrow M_0 \rightarrow 0$$

This is exact at M_i for $0 < i < n$, giving the Tor vanishing.

(b) $\Rightarrow$ (a). Take the start of a projective resolution of $\text{Hom}_A(M, X)$ as a right B -module, say

$$P_n \xrightarrow{g_n} \cdots \rightarrow P_0 \xrightarrow{g_0} \text{Hom}_A(M, X) \rightarrow 0$$

Applying $-\otimes_B M$ gives a complex, which by the hypotheses is exact:

$$M_n \xrightarrow{f_n} \cdots \rightarrow M_0 \xrightarrow{f_0} X \rightarrow 0,$$

where $M_i = P_i \otimes_B M \in \text{add } M$. Applying $\text{Hom}_A(M, -)$ to this, gives a complex

$$\text{Hom}_A(M, M_n) \rightarrow \cdots \rightarrow \text{Hom}_A(M, M_0) \rightarrow \text{Hom}_A(M, X) \rightarrow 0.$$

Identifying $\text{Hom}_A(M, M_i) = \text{Hom}_A(M, P_i \otimes_B M) \cong P_i$, we see that this is the projective resolution we started with, so it is exact. Thus (a) holds.

Remark: if we took the projective resolution to be minimal, then the maps g_i would all be right minimal in the sense of section 1.6. It follows that the maps f_i are right minimal, for otherwise there is a decomposition $M_i = M'_i \oplus M''_i$ with $M''_i \neq 0$ and $f_i(M''_i) = 0$. But then we get

$$P_i \cong \text{Hom}_A(M, M_i) \cong \text{Hom}_A(M, M'_i) \oplus \text{Hom}_A(M, M''_i)$$

and g_i is zero on the summand corresponding to $\text{Hom}_A(M, M''_i)$, contradicting the minimality of g_i . $\square$

Dually we have the following.

Proposition (2). *Let M be an A -module and $n \geq 0$ and consider M also as a B -module, where $B = \text{End}_A(M)$. For an A -module X , the following are equivalent.*

(a') *There is an exact sequence $0 \rightarrow X \rightarrow M^0 \rightarrow \cdots \rightarrow M^n$ with $M^i \in \text{add } M$ such that the sequence*

$$\text{Hom}(M^{n-1}, M) \rightarrow \cdots \rightarrow \text{Hom}(M^0, M) \rightarrow \text{Hom}(X, M) \rightarrow 0$$

is exact (this is automatic if M is injective).

(b') *The natural map $X \rightarrow \text{Hom}_B(\text{Hom}_A(X, M), M)$ is a monomorphism (in case $n = 0$) or an isomorphism (in case $n > 0$) and $\text{Ext}_B^i(\text{Hom}_A(X, M), M) = 0$ for $0 < i < n$.*

Definition. We define $\text{cogen}^n(M)$ to be the full subcategory of $A\text{-mod}$ given by the modules X satisfying these conditions. By the second condition it is a module class. Clearly

$$\text{add}(M) \subseteq \cdots \subseteq \text{cogen}^2(M) \subseteq \text{cogen}^1(M) \subseteq \text{cogen}^0(M) = \text{cogen}(M).$$

It is clear from conditions (a) and (a') that $X \in \text{cogen}^n({}_A M) \Leftrightarrow DX \in \text{gen}_n({}_{A^{op}} DM)$.

4.2 Faithfully balanced modules and endomorphism correspondence

Definition. Let M be an A -module, and let $B = \text{End}_A(M)$. Then M can be considered as a B -module, and there is a natural map

$$A \rightarrow \text{End}_B(M).$$

Clearly M is faithful iff this map is injective. We say that M is a *balanced* A -module or that M has the *double centralizer property* if this map is onto, and that M is *faithfully balanced* (f.b.) if this map is an isomorphism.

Clearly M is a f.b. A -module iff DM is a f.b. A^{op} -module.

Lemma. Let M be an A -module.

- (i) M is f.b. iff $A \in \text{cogen}^1(M)$ iff $DA \in \text{gen}_1(M)$.
- (ii) If M is a generator or cogenerator, it is f.b.

Proof. (i) Apply the second proposition in the last section with $X = A$ and $n = 1$. Now M is f.b. iff DM is f.b. iff $A^{op} \in \text{cogen}^1({}_{A^{op}} DM)$ iff $DA \in \text{gen}_1({}_A M)$.

(ii) If M is a generator, then $A \in \text{add}(M) \subseteq \text{cogen}^1(M)$. If M is a cogenerator, then DM is a generator, so f.b., hence so is M . $\square$

Definition. By an *f.b. pair* we mean a pair (A, M) consisting of an algebra and a f.b. A -module.

Given an f.b. pair, we construct a new f.b. pair (B, M) , its *endomorphism correspondent*, where $B = \text{End}_A(M)$ and M is considered in the natural way as a B -module.

Repeating the construction twice, one recovers essentially the original pair.

We say that f.b. pairs (A, M) and (A', M') are *equivalent* if there is an equivalence $A\text{-mod} \rightarrow A'\text{-mod}$ sending $\text{add}(M)$ to $\text{add}(M')$.

One can show that equivalent pairs have equivalent endomorphism correspondents.

Theorem. *If (A, M) and (B, M) are f.b. pairs which are endomorphism correspondents, then $\text{Hom}_A(-, M)$ and $\text{Hom}_B(-, M)$ give inverse antiequivalences between $\text{cogen}^1({}_A M)$ and $\text{cogen}^1({}_B M)$.*

Proof. In view of (b') in the second proposition of the last section, and the symmetrical role of A and B , it suffices to show that if $X \in \text{cogen}^1({}_A M)$, then $\text{Hom}_A(X, M) \in \text{cogen}^1({}_B M)$. Take a free presentation of ${}_A X$, say $A^m \rightarrow A^n \rightarrow X \rightarrow 0$. Applying $\text{Hom}_A(-, M)$ gives an exact sequence


$$0 \rightarrow \text{Hom}_A(X, M) \rightarrow M^n \rightarrow M^m.$$

Applying $\text{Hom}_B(-, M)$ to this gives

$$A^m \rightarrow A^n \rightarrow \text{Hom}_B(\text{Hom}_A(X, M), M) \rightarrow 0$$

which is isomorphic to the original exact sequence, so exact. Thus $\text{Hom}_A(X, M) \in \text{cogen}^1({}_B M)$. $\square$

Example. Let A be the path algebra of the linear quiver $Q = 1 \rightarrow 2 \rightarrow 3$. We display its AR quiver below. Let ${}_A M$ be the direct sum of the circled indecomposables.




The endomorphism algebra of ${}_A M$ is



Considering M as a B -module, means to consider it as a representation of this quiver. The vector space at each vertex is the corresponding indecomposable A -module. In this example, the indecomposable A -modules are at most one-dimensional at each vertex of Q . In the following diagram we write i for the

natural basis element at vertex i of Q . The arrows in the quiver for B correspond to homomorphisms of the indecomposable A -modules, and act on the basis elements as indicated below.



Thus

$${}_B M \cong \begin{array}{c} \text{ } \\ \text{ } \\ \text{ } \\ \text{ } \end{array} \oplus \begin{array}{c} \text{ } \\ \text{ } \\ \text{ } \\ \text{ } \end{array} \oplus \begin{array}{c} \text{ } \\ \text{ } \\ \text{ } \\ \text{ } \end{array}$$

Observe that ${}_A M$ has all of the projective A -modules as summands, but not all injectives, so ${}_A M$ is a generator but not a cogenerator. On the other hand all of the summands of ${}_B M$ are projective, and one summand is not injective.

Proposition. *Let (A, M) and (B, M) be f.b. pairs which are endomorphism correspondents. Then:*

- (a) ${}_A M$ is a generator iff ${}_B M$ is projective.
- (b) ${}_A M$ is a cogenerator iff ${}_B M$ is injective.
- (c) $A \in \text{cogen}^n({}_A M)$ iff $\text{Ext}_B^i(M, M) = 0$ for $0 < i < n$.

Proof. (a) If ${}_A M$ is a generator, then $A \in \text{add}({}_A M)$, so ${}_B M \cong \text{Hom}_A(A, M) \in \text{add}(\text{Hom}_A(M, M)) = \text{add}(B)$, so ${}_B M$ is projective.

Conversely if ${}_B M$ is projective, then ${}_B M \in \text{add}(B)$, so $A \cong \text{Hom}_B(M, M) \in \text{add}(\text{Hom}_B(B, M)) = \text{add}({}_A M)$.

(b) Apply (a) to DM .

(c) Second proposition in last section with $X = A$. □

For (a), see Azumaya, Completely faithful modules and self-injective rings, Nagoya Math. J. 1966. Also (a) is similar to the Wedderburn correspondence introduced by Auslander, Representation theory of Artin algebras I, Comm. Algebra 1974.

For things similar to (b), see T. Kato, Rings of U-dominant dimension ≥ 1 , Tohoku Math. J. 1969.

(c) is essentially Müller, The classification of algebras by dominant dimension, Canad. J. Math 1968.

See also:

B. Ma and J. Sauter, On faithfully balanced modules, F-cotilting and F-Auslander algebras, Journal of Algebra 2020.

M. Pressland and J. Sauter, On quiver Grassmannians and orbit closures for gen-finite modules, Algebras and Representation Theory 2022

4.3 Dominant dimension and Auslander correspondence

Definition. Given an algebra A , we take the minimal injective resolution

$$0 \rightarrow A \rightarrow I^0 \rightarrow I^1 \rightarrow \dots$$

of the module ${}_A A$. We say that A has *dominant dimension* $\geq n$ if $I^0, \dots, I^{n-1}$ are projective. This defines $\text{dom. dim } A \in \{0, 1, 2, \dots\} \cup \{\infty\}$.

Recall that an algebra A is *QF-3* if it A has a faithful projective-injective module M . If so, then $\text{add}(M) = \mathcal{P}_A \cap \mathcal{I}_A$, since any indecomposable projective-injective module embeds in A , so in some M^n , so is in $\text{add}(M)$. Thus M is unique, up to multiplicities, since it is the direct sum of all indecomposable projective-injective modules, each with some non-zero multiplicity.

Proposition. (i) $\text{dom. dim } A \geq 1$ iff A is QF-3.

(ii) $\text{dom. dim } A \geq 2$ iff A has a f.b. projective-injective M .

Proof. (i) If A is QF-3, with faithful projective-injective module M , then there is an embedding $A \rightarrow M^n$, and then the injective envelope of A is a direct summand of M^n , so it is projective.

(ii) If $\text{dom. dim } A \geq 2$, there is an exact sequence $0 \rightarrow A \rightarrow I^0 \rightarrow I^1$ with I^0, I^1 projective-injective. Let M be the direct sum of the indecomposable projective-injectives, then $A \in \text{cogen}^1(M)$ by condition (a) in the characterization of $\text{cogen}^1(M)$. Thus M is f.b.

Conversely suppose A has a f.b. projective-injective M . Since it is f.b., $A \in \text{cogen}^1(M)$. By the characterization of this means that there is an exact sequence

$$0 \rightarrow A \xrightarrow{\theta} M^0 \rightarrow M^1$$

with $M^0, M^1 \in \text{add}(M)$. Moreover by the dual result to the remark at the end of the proof of (b) $\Rightarrow$ (a) in Proposition (1) in the first subsection, we may suppose that the maps in this exact sequence are left minimal. Now the M^i are projective-injective, so they are injective, so this is the start of the injective resolution of A . Thus $I^0 \cong M^0$ and $I^1 \cong M^1$ are projective. Thus $\text{dom. dim } B \geq 2$. $\square$

For the following, see C. M. Ringel, Artin algebras of dominant dimension at least 2, manuscript 2007, available from his Bielefeld homepage.

Theorem (Morita-Tachikawa correspondence). *Endomorphism correspondence gives a 1:1 correspondence between equivalence classes of pairs (A, M) where ${}_A M$ is a generator-cogenerator and Morita equivalence classes of algebras B with $\text{dom. dim } B \geq 2$.*

The correspondence sends (A, M) to $B = \text{End}_A(M)$, and it sends B to $A = \text{End}_B(M)$ where ${}_B M$ is the faithful projective-injective B -module.

Proof. By endomorphism correspondence, the pairs (A, M) are in 1:1 correspondence with f.b. pairs (B, M) with M projective-injective. By the discussion above, these are in 1:1 correspondence with the Morita equivalence classes of algebras B with $\text{dom. dim } B \geq 2$. $\square$

The following correspondence comes from Auslander, Representation dimension of Artin algebras, Queen Mary College Lecture Notes, 1971. See also Auslander, Representation theory of Artin algebras II, Comm. Algebra 1974.

Theorem (Auslander correspondence). *There is a 1-1 correspondence between algebras A of finite representation type up to Morita equivalence and algebras B with $\text{gl. dim } B \leq 2 \leq \text{dom. dim } B$ up to Morita equivalence.*

The correspondence sends A to $B = \text{End}_A(M)$ where ${}_A M$ is the direct sum of all the indecomposable A -modules, and it sends B to $A = \text{End}_B(M)$ where ${}_B M$ is the faithful projective-injective B -module.

The algebra B is called the Auslander algebra of A .

Proof. We show that under endomorphism correspondence, pairs (A, M) where $\text{add}(M) = A\text{-mod}$ correspond to pairs (B, M) where $\text{gl. dim } B \leq 2 \leq \text{dom. dim } B$ and ${}_B M$ is the faithful projective-injective.

Suppose $\text{add}(M) = A\text{-mod}$. Given a B -module Z , choose a projective presentation

$$P_1 \xrightarrow{f} P_0 \rightarrow Z \rightarrow 0.$$

Applying $\text{Hom}_B(-, M)$ gives an exact sequence

$$0 \rightarrow \text{Hom}_B(Z, M) \rightarrow \text{Hom}_B(P_0, M) \xrightarrow{g} \text{Hom}_B(P_1, M) \rightarrow \text{Coker}(g) \rightarrow 0.$$

Applying $\text{Hom}_A(-, M)$ we get a commutative diagram with bottom row exact

$$\begin{array}{ccccccc} & & P_1 & & \xrightarrow{f} & & P_0 \\ & & \downarrow & & & & \downarrow \\ 0 & \longrightarrow & \text{Hom}_A(\text{Coker}(g), M) & \longrightarrow & \text{Hom}_A(\text{Hom}_B(P_1, M), M) & \longrightarrow & \text{Hom}_A(\text{Hom}_B(P_0, M), M) \end{array}$$

The two vertical maps are isomorphisms, so $\text{Ker}(f) \cong \text{Hom}_A(\text{Coker}(g), M)$. Now $\text{Coker}(g)$ is an A -module, so in $\text{add}(M)$, so as a B -module, we have

$$\text{Hom}_A(\text{Coker}(g), M) \in \text{add}(\text{Hom}_A(M, M)) = \text{add}(B),$$

so it is projective. Thus $\text{proj. dim } Z \leq 2$. Thus $\text{gl. dim } B \leq 2$.

Conversely suppose $\text{gl. dim } B \leq 2 \leq \text{dom. dim } B$. If Y is an A -module, it has a projective resolution starting

$$P_1 \rightarrow P_0 \rightarrow Y \rightarrow 0.$$

Applying $\text{Hom}_A(-, M)$ we get an exact sequence of B -modules

$$0 \rightarrow \text{Hom}_A(Y, M) \rightarrow \text{Hom}_A(P_0, M) \rightarrow \text{Hom}_A(P_1, M).$$

The $\text{Hom}_A(P_i, M)$ are projective B -modules since they are in $\text{add}({}_B M)$, and ${}_B M$ is projective. Thus, since $\text{gl. dim } B \leq 2$, by dimension shifting we see that $\text{Hom}_A(Y, M)$ is a projective B -module. Now ${}_B M$ is injective, so applying $\text{Hom}_B(-, {}_B M)$ gives an exact sequence

$$\text{Hom}_B(\text{Hom}_A(P_1, M), M) \rightarrow \text{Hom}_B(\text{Hom}_A(P_0, M), M) \rightarrow \text{Hom}_B(\text{Hom}_A(Y, M), M) \rightarrow 0.$$

For any A -module X there is a natural transformation from X to $\text{Hom}_B(\text{Hom}_A(X, M), M)$, and this is an isomorphism for X projective. We deduce that

$$Y \cong \text{Hom}_B(\text{Hom}_A(Y, M), M) \in \text{add}(\text{Hom}_B(B, M)) = \text{add}({}_A M),$$

so $\text{add}(M) = A\text{-mod}$. □

Example. We can check $\text{gl. dim } B = 2 = \text{dom. dim } B$ for the Auslander algebra of the linear quiver with three vertices.

$$A = K(1 \rightarrow 2 \rightarrow 3)$$

$${}_A M = \bigoplus_{i=1}^3 P(i) \text{ full module}$$

$$B = \text{End}_A(M) = K \left(\begin{array}{ccc} & & \\ & & \\ & & \end{array} \right)$$

$$\begin{aligned} p[a] &= I[c] \\ p[b] &= I[e] \\ p[c] &= I[f] \\ p[d] &= d^2 \\ p[e] &= e^2 \\ p[f] &= f \end{aligned}$$

$$\begin{aligned} 0 &\rightarrow p[d] \rightarrow I[c] \rightarrow I[c] \rightarrow I[a] \rightarrow 0 \\ 0 &\rightarrow p[e] \rightarrow I[f] \rightarrow I[c] \rightarrow I[b] \rightarrow 0 \\ 0 &\rightarrow p[f] \rightarrow I[f] \rightarrow I[e] \rightarrow I[d] \rightarrow 0 \end{aligned}$$

(projecting)

$$\begin{aligned} 0 &\rightarrow p[d] \rightarrow p[b] \rightarrow p[a] \rightarrow S[a] \rightarrow 0 \\ 0 &\rightarrow p[e] \rightarrow p[c] \oplus p[d] \rightarrow p[b] \rightarrow S[b] \rightarrow 0 \\ 0 &\rightarrow p[e] \rightarrow p[c] \rightarrow S[c] \rightarrow 0 \\ 0 &\rightarrow p[f] \rightarrow p[e] \rightarrow p[d] \rightarrow S[d] \rightarrow 0 \\ 0 &\rightarrow p[f] \rightarrow p[e] \rightarrow S[e] \rightarrow 0 \\ 0 &\rightarrow p[f] \rightarrow p[c] \rightarrow S[f] \rightarrow 0 \end{aligned}$$

Definition. Let $n \geq 1$. A module ${}_A M$ is an n -cluster tilting object if

- (i) $\text{Ext}_A^i(M, M) = 0$ for $0 < i < n$
- (ii) $\text{Ext}_A^i(U, M) = 0$ for $0 < i < n$ implies $U \in \text{add } M$
- (iii) $\text{Ext}_A^i(M, U) = 0$ for $0 < i < n$ implies $U \in \text{add } M$

Clearly (ii) implies $A \in \text{add } M$ and (iii) implies $DA \in \text{add } M$, so any n -cto is a generator-cogenerator.

Observe that M is a 1-cto iff $\text{add}(M) = A\text{-mod}$.

Example. For the algebra with quiver

$$0 \rightarrow 1 \rightarrow 2 \rightarrow \cdots \rightarrow n$$

with all paths of length 2 zero, the module $S[0]$ has projective resolution

$$0 \rightarrow P[n] \rightarrow P[n-1] \rightarrow \cdots \rightarrow P[1] \rightarrow P[0] \rightarrow S[0] \rightarrow 0$$

so $\dim \text{Ext}^i(S[0], S[j]) = \delta_{ij}$. It follows that

$$M = S[0] \oplus P[0] \oplus \cdots \oplus P[n-1] \oplus P[n] \cong I[0] \oplus I[1] \oplus \cdots \oplus I[n] \oplus S[n]$$

is an n -cto. Its endomorphism algebra B is the path algebra of the quiver

$$n \rightarrow \cdots \rightarrow 1 \rightarrow 0 \rightarrow *$$

with all paths of length 2 zero. It has global dimension $n+1$. The projectives $P[n], \dots, P[0]$ are injective, and $P[*]$ has injective resolution

$$0 \rightarrow P[*] \rightarrow I[*] \rightarrow I[0] \rightarrow \cdots \rightarrow I[n-1] \rightarrow I[n] \rightarrow 0.$$

Now $I[*] \cong P[0]$, $I[0] \cong P[1]$, $\dots$, $I[n-1] \cong P[n]$ and $I[n] \cong S[n]$ is not projective, so $\text{dom. dim } B = n+1$.

The following generalization of Auslander correspondence is due to Iyama, Auslander correspondence, *Advances in Math.* 2007.

Theorem (Iyama). *There is a 1:1 correspondence between equivalence classes of pairs (A, M) where ${}_A M$ is an n -cto and Morita equivalence classes of algebras B with $\text{gl. dim } B \leq n+1 \leq \text{dom. dim } B$.*

Proof. (To be omitted.) We are in the setting of Morita-Tachikawa correspondence.

Now $\text{Ext}_A^i(M, M) = 0$ for $1 < i < n$ corresponds to $B \in \text{cogen}^n({}_B M)$, and since ${}_B M$ is the faithful projective-injective, this corresponds to $\text{dom. dim } B \geq n+1$.

Suppose $\text{gl. dim } B \leq n+1$.

We show that if $\text{Ext}_A^i(U, M) = 0$ for $0 < i < n$ then $U \in \text{add } M$. Take the start of a projective resolution of U , say

$$P_n \rightarrow \cdots \rightarrow P_0 \rightarrow U \rightarrow 0.$$

Applying $\text{Hom}_A(-, M)$ gives a complex

$$0 \rightarrow \text{Hom}_A(U, M) \rightarrow \text{Hom}_A(P_0, M) \rightarrow \cdots \rightarrow \text{Hom}_A(P_n, M)$$

which is exact because the Exts vanish. Since ${}_B M$ is injective, applying $\text{Hom}_B(-, M)$ gives an exact sequence

$$\text{Hom}_B(\text{Hom}_A(P_n, M), M) \rightarrow \cdots \rightarrow \text{Hom}_B(\text{Hom}_A(P_0, M), M) \rightarrow \text{Hom}_B(\text{Hom}_A(U, M), M) \rightarrow 0.$$

Now the maps $P_i \rightarrow \text{Hom}_B(\text{Hom}_A(P_i, M), M)$ are isomorphisms since $P_i \in \text{add } M$. Thus the map $U \rightarrow \text{Hom}_B(\text{Hom}_A(U, M), M)$ is an iso (so $U \in \text{cogen}^1({}_A M)$). Also $\text{Hom}_A(P_i, M) \in \text{add}(\text{Hom}_A(A, M)) = \text{add}({}_B M)$. Thus, since $\text{gl. dim } B \leq n + 1$, the B -module $\text{Hom}_A(U, M)$ must be projective, so it is in $\text{add}({}_B B)$, and then $U \cong \text{Hom}_B(\text{Hom}_A(U, M), M) \in \text{add}(\text{Hom}_B(B, M)) = \text{add}({}_A M)$.

Next we show that if $\text{Ext}_A^i(M, U) = 0$ for $0 < i < n$ then $U \in \text{add } M$. Take the start of an injective resolution of U , say

$$0 \rightarrow U \rightarrow I^0 \rightarrow \cdots \rightarrow I^n.$$

Applying $\text{Hom}_A(M, -)$ gives a complex

$$0 \rightarrow \text{Hom}_A(M, U) \rightarrow \text{Hom}_A(M, I^0) \rightarrow \cdots \rightarrow \text{Hom}_A(M, I^n)$$

which is exact because the Exts vanish. Since ${}_B M$ is projective, applying $- \otimes_B M$ gives an exact sequence

$$0 \rightarrow \text{Hom}_A(M, U) \otimes_B M \rightarrow \text{Hom}_A(M, I^0) \otimes_B M \rightarrow \cdots \rightarrow \text{Hom}_A(M, I^n) \otimes_B M.$$

Now the maps $I^i \rightarrow \text{Hom}_A(M, I^i) \otimes_B M$ are isomorphisms since $I^i \in \text{add } M$. Thus the map $U \rightarrow \text{Hom}_A(M, U) \otimes_B M$ is an iso. Also $\text{Hom}_A(M, I^i) \in \text{add}(\text{Hom}_A(M, M)) = \text{add}({}_B B)$. Thus, since $\text{gl. dim } B \leq n + 1$, the right B -module $\text{Hom}_A(M, U)$ must be projective, so it is in $\text{add}({}_B B)$, and then $U \cong \text{Hom}_A(M, U) \otimes_B M \in \text{add}({}_B B \otimes_B M) = \text{add}({}_A M)$.

Now suppose that M is an n -cto. Given a B -module Z , choose a projective presentation

$$P_1 \xrightarrow{f} P_0 \rightarrow Z \rightarrow 0.$$

Applying $\text{Hom}_B(-, M)$ gives an exact sequence

$$0 \rightarrow \text{Hom}_B(Z, M) \rightarrow \text{Hom}_B(P_0, M) \xrightarrow{g} \text{Hom}_B(P_1, M) \rightarrow \text{Coker}(g) \rightarrow 0.$$

Let $C^0 = \text{Coker}(g)$. Applying $\text{Hom}_A(-, M)$ we get a commutative diagram with bottom row exact

$$\begin{array}{ccccccc} & & P_1 & & \xrightarrow{f} & & P_0 \\ & & \downarrow & & & & \downarrow \\ 0 & \longrightarrow & \text{Hom}_A(C^0, M) & \longrightarrow & \text{Hom}_A(\text{Hom}_B(P_1, M), M) & \longrightarrow & \text{Hom}_A(\text{Hom}_B(P_0, M), M) \end{array}$$

The two vertical maps are isomorphisms, so $\text{Ker}(f) \cong \text{Hom}_A(C^0, M)$.

Now since M is a cogenerator, by repeatedly taking left M -approximations we can get an exact sequence

$$0 \rightarrow C^0 \rightarrow M^0 \rightarrow \dots \rightarrow M^{n-2}$$

such that the sequence

$$\text{Hom}_A(M^{n-2}, M) \rightarrow \dots \rightarrow \text{Hom}_A(M^0, M) \rightarrow \text{Hom}_A(C^0, M) \rightarrow 0$$

is exact. Let C^i be the cosyzygies for this sequence, so

$$0 \rightarrow C^i \rightarrow M^i \rightarrow C^{i+1} \rightarrow 0.$$

Then

$$\text{Hom}(M^i, M) \twoheadrightarrow \text{Hom}(C^i, M) \rightarrow \text{Ext}^1(C^{i+1}, M) \rightarrow \text{Ext}^1(M^i, M) = 0 \rightarrow \dots,$$

so by dimension shifting

$$\text{Ext}^{n-1}(C^{n-1}, M) \cong \text{Ext}^{n-2}(C^{n-1}, M) \cong \dots \cong \text{Ext}^1(C^1, M) = 0$$

and similarly $\text{Ext}^i(C^{n-1}, M) = 0$ for $0 < i < n$. Thus $C^{n-1} \in \text{add } M$. Thus Z has projective resolution

$$0 \rightarrow \text{Hom}_A(C^{n-1}, M) \rightarrow \text{Hom}_A(M^{n-2}, M) \rightarrow \dots \rightarrow \text{Hom}_A(M^0, M) \rightarrow P_1 \rightarrow P_0 \rightarrow Z \rightarrow 0.$$

Thus $\text{proj. dim } Z \leq n + 1$. Thus $\text{gl. dim } B \leq n + 1$. □

4.4 Homological conjectures for f.d. algebras

Let $0 \rightarrow A \rightarrow I^0 \rightarrow I^1 \rightarrow \dots$ be the minimal injective resolution of a f.d. algebra A . Recall that A has dominant dimension $\geq n$ if $I^0, \dots, I^{n-1}$ are all projective.

Conjecture (Nakayama conjecture 1958). *If all I^n are projective, i.e. $\text{dom. dim } A = \infty$, then A is self-injective.*

Proposition. *The following are equivalent.*

- (i) *The Nakayama conjecture (if $\text{dom. dim } B = \infty$ then B is self-injective).*
- (ii) *If ${}_A M$ is a generator-cogenerator and $\text{Ext}_A^i(M, M) = 0$ for all $i > 0$ then M is projective.*

Proof. (i) implies (ii). Say ${}_A M$ satisfies the hypotheses. Let (B, M) be the endomorphism correspondent. Then ${}_B M$ is projective-injective and $B \in \text{cogen}^n(M)$ for all n . Thus for all n there is an exact sequence

$$0 \rightarrow B \rightarrow I^0 \rightarrow \cdots \rightarrow I^n$$

with the I^i projective-injective. Thus $\text{dom.dim } B = \infty$. Thus B is self-injective, so $\text{add}(M) = \text{add}(B)$, so ${}_B M$ is a generator, so ${}_A M$ is projective.

(ii) implies (i). Say $\text{dom.dim } B = \infty$. Thus B is QF-3 and let ${}_B M$ be the faithful projective-injective module. Let ${}_A M$ be the endomorphism correspondent. It is a generator-cogenerator. Then $B \in \text{cogen}^n(M)$ for all n , so $\text{Ext}_A^i(M, M) = 0$ for all $i > 0$. Thus by (ii), ${}_A M$ is projective, so ${}_B M$ is a generator. Thus $B \in \text{add}(M)$ is injective. $\square$

Conjecture (Generalized Nakayama conjecture, Auslander and Reiten 1975). *For any f.d. algebra A , every indecomposable injective occur as a summand of some I^n .*

It clearly implies the Nakayama conjecture, for if the I^n are projective, and each indecomposable injective occurs as a summand of some I^n , then the indecomposable injectives are projective.

Example. For the commutative square, vertices 1(source), 2, 3, 4(sink). There are injective resolutions

$$\begin{aligned} 0 \rightarrow P[1] \rightarrow I[4] \rightarrow 0, \\ 0 \rightarrow P[2] \rightarrow I[4] \rightarrow I[3] \rightarrow 0, \\ 0 \rightarrow P[3] \rightarrow I[4] \rightarrow I[2] \rightarrow 0, \\ 0 \rightarrow P[4] \rightarrow I[4] \rightarrow I[2] \oplus I[3] \rightarrow I[1] \rightarrow 0, \end{aligned}$$

so

$$0 \rightarrow A \rightarrow I[4]^4 \rightarrow I[2]^2 \oplus I[3]^2 \rightarrow I[1] \rightarrow 0,$$

so all indecomposable injectives occur.

Proposition. *The following are equivalent.*

- (i) *The generalized Nakayama conjecture (every indecomposable injective occurs as a summand of some I^i in the minimal injective resolution of B).*
- (ii) *If ${}_A M$ is a cogenerator and $\text{Ext}_A^i(M, M) = 0$ for all $i > 0$ then M is injective.*

Proof. (i) implies (ii). Suppose ${}_A M$ satisfies the conditions. Then there is corresponding ${}_B M$ which is injective, and $B \in \text{cogen}^n(M)$ for all n . Thus by (i) every indecomposable injective is a summand of ${}_B M$. Thus ${}_B M$ is a cogenerator. Thus ${}_A M$ is injective.

(ii) implies (i). Let ${}_B M$ be the sum of all indecomposable injectives occurring in the I^i . Then $B \in \text{cogen}^n(M)$ for all n . Let ${}_A M$ be the endomorphism correspondent. Then ${}_A M$ is a cogenerator and $\text{Ext}_A^i(M, M) = 0$ for all $i > 0$. Thus by (ii) ${}_A M$ is injective. Thus ${}_B M$ is a cogenerator. Thus all indecomposable injectives occur as a summand of ${}_B M$. $\square$

For the next conjecture, see Happel, Selforthogonal modules, 1995.

Conjecture (Boundedness Conjecture). *If M is an A -module with $\text{Ext}_A^i(M, M) = 0$ for all $i > 0$ then $\#M \leq \#A$, where $\#M$ denotes the number of non-isomorphic indecomposable summands of M .*

Since a cogenerator has all indecomposable injectives as summands, the boundedness conjecture implies the generalized Nakayama conjecture.

Definition. An algebra A is (Iwanaga) Gorenstein if both $\text{inj. dim } {}_A A < \infty$ and $\text{inj. dim } A_A < \infty$.

In Auslander and Reiten, Applications of contravariantly finite subcategories, Adv. Math 1991, one finds:

Conjecture (Gorenstein Symmetry Conjecture). *If one of $\text{inj. dim } {}_A A$ and $\text{inj. dim } A_A$ is finite, so is the other.*

Lemma. (i) *If $\text{inj. dim } {}_A A = n < \infty$, any A -module has $\text{proj. dim } M \leq n$ or ∞ .*
(ii) *If $\text{inj. dim } {}_A A = n$ and $\text{inj. dim } A_A = m$ are both finite, they are equal.*

For example, by (i) every non-projective module for a self-injective algebra has infinite projective dimension.

Proof. (i) Say $\text{proj. dim } M = i < \infty$. There is some N with $\text{Ext}^i(M, N) \neq 0$. Choose $0 \rightarrow L \rightarrow P \rightarrow N \rightarrow 0$ with P projective. The long exact sequence for $\text{Hom}(M, -)$ gives

$$\dots \rightarrow \text{Ext}^i(M, P) \rightarrow \text{Ext}^i(M, N) \rightarrow \text{Ext}^{i+1}(M, L) \rightarrow \dots$$

Now $\text{Ext}^{i+1}(M, L) = 0$, so $\text{Ext}^i(M, P) \neq 0$, so $\text{Ext}^i(M, A) \neq 0$, so $i \leq n$.

(ii) $\text{proj. dim } {}_A A = \text{inj. dim } A_A = m$, so $m \leq n$ by (i). Dually $m \geq n$. $\square$

This also holds for noetherian rings, see Zaks, Injective dimension of semi-primary rings, J. Alg. 1969.

For the following, see H. Bass, Finitistic dimension and a homological generalization of semiprimary rings, Trans. Amer. Math. Soc. 1960.

Conjecture (Finitistic Dimension Conjecture). *For any f.d. algebra A ,*

$$\text{fin. dim } A = \sup\{\text{proj. dim } M \mid \text{proj. dim } M < \infty\}$$

is finite.

For example if A is Gorenstein, with $\text{inj. dim } {}_A A = n = \text{inj. dim } A_A$, then $\text{fin. dim } A = n$. For the lemma implies that any A -module M has $\text{proj. dim } M \leq n$ or ∞ , and $\text{proj. dim } D(A_A) = n$.

Note that $\text{fin. dim } A$ is not necessarily the same as the maximum of the projective dimensions of the simple modules of finite projective dimension.

There is also a big finitistic dimension, where the modules need not be finite-dimensional, and this may also always be finite.

Proposition. *The finitistic dimension conjecture implies the Gorenstein symmetry conjecture.*

Proof. Assuming $\text{inj. dim } A_A = n < \infty$, we want to prove that $\text{inj. dim } {}_A A < \infty$. We have $\text{proj. dim } {}_A D A = n < \infty$. Thus any injective module has projective dimension $< \infty$. Take a minimal injective resolution $0 \rightarrow {}_A A \rightarrow I^0 \rightarrow \dots$. We show by induction on i that $\text{proj. dim } \Omega^i A < \infty$. There is an exact sequence

$$0 \rightarrow \Omega^{i-1} A \rightarrow I^{i-1} \rightarrow \Omega^i A \rightarrow 0.$$

Applying $\text{Hom}_A(-, X)$ for a module X gives a long exact sequence

$$\dots \rightarrow \text{Ext}^m(\Omega^{i-1} A, X) \rightarrow \text{Ext}^{m+1}(\Omega^i A, X) \rightarrow \text{Ext}^{m+1}(I^{i-1}, X) \rightarrow \dots$$

For m sufficiently large, independent of X , the outside terms are zero, hence so is the middle.

Let $i > 0$. If $\Omega^i A = 0$, or is injective, then $\text{inj. dim } {}_A A < \infty$, as desired, so suppose otherwise. Let $f : \Omega^i A \rightarrow I^i$ be the inclusion. Then f belongs to the middle term in the complex

$$\text{Hom}(\Omega^i A, I^{i-1}) \rightarrow \text{Hom}(\Omega^i A, I^i) \rightarrow \text{Hom}(\Omega^i A, I^{i+1})$$

and it is sent to zero in the third term. Now f is not in the image of the map from the first term, for otherwise the map $I^{i-1} \rightarrow \Omega^i A$ is a split epimorphism, so $\Omega^i A$ is injective. Thus the homology of this complex at the middle term is non-zero. Thus $\text{Ext}^i(\Omega^i A, A) \neq 0$. Thus $\text{proj. dim } \Omega^i A \geq i$. This contradicts that $\text{fin. dim } A < \infty$. $\square$

Proposition. *The finitistic dimension conjecture implies the generalized Nakayama conjecture.*

Proof. Assume the FDC. We show that if ${}_A M$ is a module and $\text{Ext}^n(M, A) = 0$ for all $n \geq 0$ then $M = 0$ (the *strong Nakayama conjecture*).

If $S[i]$ is a simple A -module and $I[i]$ is its injective envelope, recall from section 1.10, that $\dim \text{Ext}^n(S[i], A)$ is $\dim \text{End}(S[i])$ times the multiplicity of $I[i]$ as a direct summand of I^n . Thus taking $M = S[i]$, the strong Nakayama conjecture gives the generalized Nakayama conjecture.

Take a minimal projective resolution $\cdots \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$. By assumption the sequence

$$0 \rightarrow \text{Hom}_A(P_0, A) \xrightarrow{f_0} \text{Hom}_A(P_1, A) \xrightarrow{f_1} \text{Hom}_A(P_2, A) \rightarrow \cdots$$

of right A -modules is exact. Let $\text{fin. dim } A^{op} = n < \infty$. Then $\text{Coker}(f_n)$ has projective resolution

$$0 \rightarrow \text{Hom}_A(P_0, A) \xrightarrow{f_0} \text{Hom}_A(P_1, A) \rightarrow \cdots \rightarrow \text{Hom}_A(P_{n+1}, A) \rightarrow \text{Coker}(f_n) \rightarrow 0$$

so it has finite projective dimension, so projective dimension $\leq n$, so by dimension shifting $\text{Im } f_1$ is projective, so f_0 must be a split mono. But $\text{Hom}_A(-, A)$ is an antiequivalence from $\mathcal{P}_A$ to $\mathcal{P}_{A^{op}}$. Thus the map $P_1 \rightarrow P_0$ must be a split epi, so $M = 0$. $\square$

4.5 No loops conjecture

It is nice to see that some homological conjecture has been proved. In this section we do not assume that K is algebraically closed, but we do assume that $A = KQ/I$ with I admissible. The following conjecture was proved by Igusa, Notes on the no loops conjecture, J. Pure Appl. Algebra 1990.

Theorem (No loops conjecture). *If $\text{gl. dim } A < \infty$ then Q has no loops (that is, $\text{Ext}^1(S[i], S[i]) = 0$ for all i).*

Proof. We use the trace function of Hattori and Stallings. I only sketch the proof of its properties.

(1) For any matrix $\theta \in M_n(A)$ we consider its trace $\text{tr}(\theta) \in A/[A, A]$, where $[A, A]$ is the subspace of A spanned by the commutators $ab - ba$. This ensures that $\text{tr}(\theta\phi) = \text{tr}(\phi\theta)$. This equality holds also for $\theta \in M_{m \times n}(A)$ and $\phi \in M_{n \times m}(A)$.

(2) If P is a f.g. projective A -module it is a direct summand of a f.g. free module $F = A^n$. Let $p : F \rightarrow P$ and $i : P \rightarrow F$ be the projection and inclusion. One defines $\text{tr}(\theta)$ for $\theta \in \text{End}(P)$ to be $\text{tr}(i\theta p)$. This is well defined, for if

$$A^n = F \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{i} \end{array} P \begin{array}{c} \xrightarrow{i'} \\ \xleftarrow{p'} \end{array} F' = A^m$$

with $pi = 1_P = p'i'$, then $\text{tr}(i\theta p) = \text{tr}((ip')(i'\theta p)) = \text{tr}((i'\theta p)(ip')) = \text{tr}(i'\theta p')$.

(3) Any module M has a finite projective resolution $P_* \rightarrow M$, and an endomorphism θ of M lifts to a map between the projective resolutions

$$\begin{array}{ccccccccc} 0 & \longrightarrow & P_n & \longrightarrow & \dots & \longrightarrow & P_1 & \longrightarrow & P_0 & \longrightarrow & M & \longrightarrow & 0 \\ & & \theta_n \downarrow & & & & \theta_1 \downarrow & & \theta_0 \downarrow & & \theta \downarrow & & \\ 0 & \longrightarrow & P_n & \longrightarrow & \dots & \longrightarrow & P_1 & \longrightarrow & P_0 & \longrightarrow & M & \longrightarrow & 0. \end{array}$$

Define $\text{tr}(\theta) = \sum_i (-1)^i \text{tr}(\theta_i)$. One can show that does not depend on the projective resolution or the lift of θ , see section 4 of Lenzing, Nilpotente Elemente in Ringen von endlicher globaler Dimension, Math. Z. 1969.

(4) One can show that given a commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' & \longrightarrow & 0 \\ & & \theta' \downarrow & & \theta \downarrow & & \theta'' \downarrow & & \\ 0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' & \longrightarrow & 0 \end{array}$$

one has $\text{tr}(\theta) = \text{tr}(\theta') + \text{tr}(\theta'')$.

(5) It follows that any nilpotent endomorphism has trace 0, since

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \text{Im } \theta & \longrightarrow & M & \longrightarrow & M/\text{Im } \theta & \longrightarrow & 0 \\ & & \theta|_{\text{Im } \theta} \downarrow & & \theta \downarrow & & 0 \downarrow & & \\ 0 & \longrightarrow & \text{Im } \theta & \longrightarrow & M & \longrightarrow & M/\text{Im } \theta & \longrightarrow & 0 \end{array}$$

so $\text{tr}(\theta) = \text{tr}(\theta|_{\text{Im } \theta}) = \text{tr}(\theta|_{\text{Im } (\theta^2)}) = \dots = 0$.

(6) Thus any element of $J(A)$ as a map $A \rightarrow A$ has trace 0, so $J(A) \subseteq [A, A]$. Thus $(KQ)_+ \subseteq I + [KQ, KQ]$.

(7) Any loop of Q gives an element of $(KQ)_+$. But it is easy to see that

$$I + [KQ, KQ] \subseteq \text{span of arrows which are not loops} + (KQ)_+^2,$$


for example if p, q are paths then $[p, q] \in (KQ)_+^2$ unless they are trivial paths or one is trivial and the other is an arrow. Thus there are no loops. $\square$

A strengthening (proved by Igusa, Liu and Paquette, A proof of the strong no loop conjecture, Adv. Math. 2011). If S is a 1-dimensional simple module for a f.d. algebra and S has finite injective or projective dimension, then $\text{Ext}^1(S, S) = 0$.

An open problem (stated by Liu and Morin, The strong no loop conjecture for special biserial algebras, Proc. Amer. Math. Soc. 2004). The extension conjecture: if S is simple module for a f.d. algebra and $\text{Ext}^1(S, S) \neq 0$ then $\text{Ext}^n(S, S) \neq 0$ for infinitely many n .

- $\#T = \#A$, the number of non-isomorphic summands of T is the number of simple A -modules.

Let $B = \text{End}_A(T)^{op}$, so T becomes an A - B -bimodule. The Brenner-Butler theorem gives equivalences between the following parts of the module categories.



5.1 Torsion theories and tau-rigid modules

The notion of a torsion theory comes from Dickson, A torsion theory for abelian categories, Trans. Amer. Math. Soc. 1966.

Definition. A *torsion theory* in an abelian category $\mathcal{A}$ is a pair of full subcategories $(\mathcal{T}, \mathcal{F})$, the *torsion* and *torsion-free* classes, such that

- (i) $\text{Hom}(\mathcal{T}, \mathcal{F}) = 0$.
- (ii) Any object X has a subobject $t_{\mathcal{T}}X \in \mathcal{T}$ with $X/t_{\mathcal{T}}X \in \mathcal{F}$ (so it fits in an exact sequence $0 \rightarrow t_{\mathcal{T}}X \rightarrow X \rightarrow X/t_{\mathcal{T}}X \rightarrow 0$ with first term in $\mathcal{T}$ and last term in $\mathcal{F}$).

Examples. (1) The torsion and torsion-free modules give a torsion theory in the category of $\mathbb{Z}$ -modules.

(2) For A the path algebra of the quiver $1 \rightarrow 2$, $A\text{-mod}$ has torsion theory (add $S[2]$, add $S[1]$).

Notation. For an a set $\mathcal{C}$ of modules in $A\text{-mod}$ or more generally of objects in an abelian category

$$\mathcal{C}^{\perp i, j, \dots} = \{X : \text{Ext}^n(M, X) = 0 \text{ for all } M \in \mathcal{C} \text{ and } n = i, j, \dots\},$$

$${}^{\perp i, j, \dots} \mathcal{C} = \{X : \text{Ext}^n(X, M) = 0 \text{ for all } M \in \mathcal{C} \text{ and } n = i, j, \dots\}.$$

Recall that $\text{Ext}^0 = \text{Hom}$.

Properties. Let $(\mathcal{T}, \mathcal{F})$ be a torsion theory.

- (i) $\mathcal{T} = {}^{\perp 0}\mathcal{F}$ and $\mathcal{F} = \mathcal{T}^{\perp 0}$ so either of the classes determines the other.
- (ii) $\mathcal{T}$ is closed under quotients and extensions; $\mathcal{F}$ is closed under subobjects and extensions.
- (iii) The subobject $t_{\mathcal{T}}X$ is uniquely determined, and the assignment sending X to $t_{\mathcal{T}}X$ defines a functor $\mathcal{A} \rightarrow \mathcal{T}$ which is a right adjoint to the inclusion $\mathcal{T}$ in $\mathcal{A}$. The assignment sending X to $X/t_{\mathcal{T}}X$ defines a functor $\mathcal{A} \rightarrow \mathcal{F}$ which is a left adjoint to the inclusion $\mathcal{F}$ in $\mathcal{A}$.

Proof. (i) If $X \in \mathcal{T}^{\perp 0}$, then $\text{Hom}(\mathcal{T}, X) = 0$, so we must have $t_{\mathcal{T}}X = 0$, so $X \cong X/t_{\mathcal{T}}X \in \mathcal{F}$. If $X \in {}^{\perp 0}\mathcal{F}$, then $\text{Hom}(X, \mathcal{F}) = 0$, so we must have $X = t_{\mathcal{T}}X \in \mathcal{T}$.

For (ii), for $\mathcal{T}$ given an exact sequence $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$, apply $\text{Hom}(-, F)$ for $F \in \mathcal{F}$ to get an exact sequence

$$0 \rightarrow \text{Hom}(Z, F) \rightarrow \text{Hom}(Y, F) \rightarrow \text{Hom}(X, F).$$

Now if $X, Z \in \mathcal{T}$, then $\text{Hom}(X, F) = \text{Hom}(Z, F) = 0$, so $\text{Hom}(Y, F) = 0$, so $Y \in \mathcal{T}$. Also, if $Y \in \mathcal{T}$, then $\text{Hom}(Y, F) = 0$, so $\text{Hom}(Z, F) = 0$, so $Z \in \mathcal{T}$.

For (iii) observe that any map $\theta : X \rightarrow Y$ induces a map $t_{\mathcal{T}}X \rightarrow t_{\mathcal{T}}Y$ since the composition $t_{\mathcal{T}}X \rightarrow X \rightarrow Y \rightarrow Y/t_{\mathcal{T}}Y$ must be zero. $\square$

Remark. A *splitting torsion theory* is one in which the sequence $0 \rightarrow t_{\mathcal{T}}X \rightarrow X \rightarrow X/t_{\mathcal{T}}X \rightarrow 0$ is always split exact.

If A is a f.d. algebra, a torsion theory in $A\text{-mod}$ is splitting if and only if every indecomposable module is either torsion or torsion-free.

A splitting torsion theory is thus given by a partition of the indecomposable modules into two sets T, F with $\text{Hom}(T, F) = 0$. Then $(\text{add } T, \text{add } F)$ is a splitting torsion theory in $A\text{-mod}$.

This is very easy to do if A is an algebra whose AR quiver is obtained by knitting, so A is of finite representation type and all of its indecomposable modules are directing. We want there to be no irreducible maps from T to F .

Proposition. *If A is a f.d. algebra, for a module class $\mathcal{T}$ in $A\text{-mod}$ the following are equivalent.*

- (i) $\mathcal{T}$ is a torsion class for some torsion theory in $A\text{-mod}$.
- (ii) $\mathcal{T} = {}^{\perp 0}(\mathcal{T}^{\perp 0})$.
- (iii) $\mathcal{T} = {}^{\perp 0}\mathcal{C}$ for some module class $\mathcal{C}$.
- (iv) $\mathcal{T}$ is closed under quotients and extensions.

Proof. (i) implies (ii) implies (iii) implies (iv). Straightforward.

(iv) implies (i). Define $\mathcal{F} = \mathcal{T}^{\perp 0}$. Given any module X , let T be a submodule of X in $\mathcal{T}$ of maximal dimension. Then $\text{Hom}(\mathcal{T}, X/T) = 0$, for if T'/T is the image of such a map, then T'/T is in $\mathcal{T}$, hence so is T' , contradicting maximality. Thus $X/T \in \mathcal{F}$. $\square$

Thus a pair of module classes $(\mathcal{T}, \mathcal{F})$ is a torsion theory in $A\text{-mod}$ if and only if $\mathcal{T} = {}^{\perp 0}\mathcal{F}$ and $\mathcal{F} = \mathcal{T}^{\perp 0}$.

Lemma (Auslander-Smalø, 1981). *For modules M, N , the following are equivalent:*

- (i) $\text{Hom}(N, \tau M) = 0$.
- (ii) $\text{Ext}^1(M, \text{gen } N) = 0$ (that is, $\text{Ext}^1(M, G) = 0$ for all $G \in \text{gen } N$).

Proof. (i) $\Rightarrow$ (ii). If $\text{Hom}(N, \tau M) = 0$, then $\text{Hom}(G, \tau M) = 0$ for all $G \in \text{gen } N$, so $\overline{\text{Hom}}(G, \tau M) = 0$, so $\text{Ext}^1(M, G) = 0$ by the Auslander-Reiten formula.

(ii) $\Rightarrow$ (i). Say $f : N \rightarrow \tau M$ is a non-zero map. Factorize it as a surjection $g : N \rightarrow G$ followed by a mono $h : G \rightarrow \tau M$. Suppose that h factors through an injective. Then it factors through the injective envelope $E(G)$ of G . Since τM has no injective summand, the induced map $E(G) \rightarrow \tau M$ cannot be injective, so its kernel is non-zero. Since G is essential in $E(G)$, the kernel meets G . Thus $G \rightarrow \tau M$ has non-zero kernel. Contradiction. Thus $\overline{\text{Hom}}(G, \tau M) \neq 0$, so $\text{Ext}^1(M, G) \neq 0$. $\square$

Definition. Given a module class $\mathcal{C}$ in $A\text{-mod}$ and $X \in \mathcal{C}$, we say that

- (i) X is *Ext-projective* in $\mathcal{C}$ if $\text{Ext}^1(X, \mathcal{C}) = 0$.
- (ii) X is *Ext-injective* in $\mathcal{C}$ if $\text{Ext}^1(\mathcal{C}, X) = 0$.

Lemma. *If $(\mathcal{T}, \mathcal{F})$ is a torsion theory in $A\text{-mod}$, then*

- (i) $X \in \mathcal{T}$ is *Ext-projective* for $\mathcal{T}$ iff $\tau X \in \mathcal{F}$.
- (ii) $X \in \mathcal{F}$ is *Ext-injective* for $\mathcal{F}$ iff $\tau^- X \in \mathcal{T}$.
- (iii) *There are bijections*

$\text{Non-proj indec Ext-projs in } \mathcal{T} \text{ up to iso} \xrightleftharpoons[\tau^-]{\tau} \text{Non-inj indec Ext-injs in } \mathcal{F} \text{ up to iso}$

Proof. (i) Say $X \in \mathcal{T}$. Then $\tau X \in \mathcal{F} \Leftrightarrow \text{Hom}(T, \tau X) = 0$ for all $T \in \mathcal{T} \Leftrightarrow \text{Ext}^1(X, \text{gen } T) = 0$ for all $T \in \mathcal{T} \Leftrightarrow X$ is Ext-projective.

(ii) is dual.

(iii) follows. $\square$

Lemma. *If $(\mathcal{T}, \mathcal{F})$ is a torsion theory in $A\text{-mod}$, then*

(i) *The Ext-injectives for $\mathcal{T}$ are the modules $t_{\mathcal{T}}I$ with I injective. The indecomposable Ext-injectives are the modules $t_{\mathcal{T}}I[i]$ with $I[i] \notin \mathcal{F}$.*

(ii) *The Ext-projectives for $\mathcal{F}$ are the modules $P/t_{\mathcal{T}}P$ with P projective. The indecomposable Ext-projectives are the modules $P[i]/t_{\mathcal{T}}P[i]$ with $P[i] \notin \mathcal{T}$.*

Proof. (i) $t_{\mathcal{T}}I$ is in $\mathcal{T}$, and it is Ext-injective since if $T \in \mathcal{T}$ and $0 \rightarrow t_{\mathcal{T}}I \rightarrow E \rightarrow T \rightarrow 0$ is an exact sequence, then the pushout along $t_{\mathcal{T}}I \rightarrow I$ splits, giving a map $E \rightarrow I$. But $E \in \mathcal{T}$, so it gives a map $E \rightarrow t_{\mathcal{T}}I$, which is a retraction for the given sequence.

Conversely suppose X is Ext-injective in $\mathcal{T}$ and $X \rightarrow I$ is its injective envelope. Then we have an injection $X \rightarrow t_{\mathcal{T}}I$. Since $\mathcal{T}$ is closed under quotients, all terms in the exact sequence $0 \rightarrow X \rightarrow t_{\mathcal{T}}I \rightarrow t_{\mathcal{T}}I/X \rightarrow 0$ are in $\mathcal{T}$. Thus this sequence splits, so X is a direct summand of $t_{\mathcal{T}}I$, and we have equality since X is essential in I .

Also, if $I[i] \notin \mathcal{F}$, then $t_{\mathcal{T}}I[i]$ is non-zero and contained in $I[i]$, so it has simple socle, so it is indecomposable.

(ii) is dual. $\square$

The following definition comes from Adachi, Iyama and Reiten, τ -tilting theory, 2014.

Definition. A module M is τ -rigid if $\text{Hom}(M, \tau M) = 0$. Dually, it is τ^- -rigid if $\text{Hom}(\tau^- M, M) = 0$

Note that M is τ -rigid iff DM is τ^- -rigid, since

$$\begin{aligned} \text{Hom}(M, \tau M) &= \text{Hom}(M, D \text{Tr } M) \cong \text{Hom}(\text{Tr } M, DM) \\ &\cong \text{Hom}(\text{Tr } DDM, DM) = \text{Hom}(\tau^- DM, DM). \end{aligned}$$

Proposition. *The following are equivalent*

- (i) M is τ -rigid.
- (ii) $\text{Ext}^1(M, \text{gen } M) = 0$.
- (iii) $\text{gen } M$ is a torsion class and M is Ext-projective in $\text{gen } M$.
- (iv) M is Ext-projective in some torsion class.

Proof. (i) $\Leftrightarrow$ (ii). The lemma of Auslander and Smalø.

(ii) $\Rightarrow$ (iii). Suppose M is τ -rigid. To show that $\text{gen } M$ is a torsion class, it suffices to show that if $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ is exact and $X, Z \in \text{gen } M$, then so is Y . Choose a surjection $M^n \rightarrow Z$. By (ii) The pullback sequence splits, so the middle term of it is in $\text{gen } M$, and hence so is Y . Now $\text{Ext}^1(M, \text{gen } M) = 0$, so M is Ext-projective.

(iii) $\Rightarrow$ (iv). Trivial.

(iv) $\Rightarrow$ (ii). If M is Ext-projective in $\mathcal{T}$, then $\text{Ext}^1(M, \text{gen } M) = 0$ since $\text{gen } M \subseteq \mathcal{T}$. $\square$

Note that the torsion theory given by a τ -rigid module M is $(\text{gen } M, M^{\perp 0})$.

Example. Let A be the path algebra of $1 \rightarrow 2 \rightarrow 3$. Let $M = 2 \oplus 123$. It is τ -rigid. Then $\mathcal{T} = \text{gen } M$ contains $123, 12, 2, 1$. The torsion-free class is $\mathcal{F} = \mathcal{T}^{\perp 0} = M^{\perp 0}$. It contains 3 and 23 .

The Ext-projectives in $\mathcal{T}$ are $2, 12, 123$.

The Ext-injectives in $\mathcal{T}$ are $1, 12, 123$.

The Ext-projectives in $\mathcal{F}$ are $3, 23$.

The Ext-injectives in $\mathcal{F}$ are $3, 23$.

The next result is dual to Theorem 4.1(c) of Auslander and Smalø, Almost split sequences in subcategories, J. Algebra 1981.

Theorem. *Let $\mathcal{T}$ be a torsion class which is functorially finite and let*

$$A \xrightarrow{f} M^0 \xrightarrow{c} M^1 \rightarrow 0$$

be an exact sequence with f be a minimal left $\mathcal{T}$ -approximation of A . Then

- (i) $\mathcal{T} = \text{gen } M^0 = \text{gen}(M^0 \oplus M^1)$.
- (ii) M^0 is a splitting projective for $\mathcal{T}$, meaning that any epimorphism $\theta : T \twoheadrightarrow M^0$ with $T \in \mathcal{T}$ must be a split epi.
- (iii) M^0 and $M^0 \oplus M^1$ are Ext-projective in $\mathcal{T}$, so they are τ -rigid.
- (iv) Any module $T \in \mathcal{T}$ is a quotient of a module in $\text{add}(M^0 \oplus M^1)$ by a submodule in $\mathcal{T}$.
- (v) Any Ext-projective in $\mathcal{T}$ is in $\text{add}(M^0 \oplus M^1)$, so there are only finitely many indecomposable Ext-projectives in $\mathcal{T}$.

Proof. (i) Clearly $\text{gen } M^0 = \text{gen}(M^0 \oplus M^1) \subseteq \mathcal{T}$. If $T \in \mathcal{T}$, then there is a map $A^n \twoheadrightarrow T$, and each component factors through M , giving an epimorphism $M^n \twoheadrightarrow T$.

(ii) Since A is projective, the map $f : A \rightarrow M^0$ lifts to a map $A \rightarrow T$. By the approximation property, this factors as $A \rightarrow M^0 \rightarrow T$. Now the composition $M^0 \rightarrow T \rightarrow M^0$ must be an isomorphism by minimality.

(iii) Let $T \in \mathcal{T}$. Any exact sequence $0 \rightarrow T \rightarrow E \rightarrow M^0 \rightarrow 0$ splits by (ii). Thus M^0 is Ext-projective.

Since f is a $\mathcal{T}$ -approximation, the induced map $\text{Hom}(M^0, T) \rightarrow \text{Hom}(A, T)$ is surjective. This is a composition $\text{Hom}(M^0, T) \rightarrow \text{Hom}(\text{Im } f, T) \rightarrow \text{Hom}(A, T)$ and the second map is injective, so actually the second map is a bijection and the first map $\text{Hom}(M^0, T) \rightarrow \text{Hom}(\text{Im } f, T)$ is surjective.

Now the exact sequence $0 \rightarrow \text{Im } f \xrightarrow{i} M^0 \xrightarrow{c} M^1 \rightarrow 0$ gives

$$\text{Hom}(M^0, T) \rightarrow \text{Hom}(\text{Im } f, T) \rightarrow \text{Ext}^1(M^1, T) \rightarrow \text{Ext}^1(M^0, T) = 0.$$

so $\text{Ext}^1(M^1, T) = 0$.

(iv) (My thanks to Andrew Hubery for this argument). Take a right $\text{add}(M^0 \oplus M^1)$ -approximation $\phi : W \rightarrow T$ for T . Since $T \in \text{gen } M^0$, the map ϕ is surjective, so it gives an exact sequence

$$0 \rightarrow U \xrightarrow{\theta} W \xrightarrow{\phi} T \rightarrow 0.$$

Given $u \in U$ there is a map $r : A \rightarrow U$, $a \mapsto au$. Since $A \rightarrow M^0$ is a $\mathcal{T}$ -approximation and $W \in \mathcal{T}$, there is a map p , and hence a map q giving a commu-

tative diagram

$$\begin{array}{ccccccc}
A & \xrightarrow{f} & M^0 & \xrightarrow{c} & M^1 & \longrightarrow & 0 \\
r \downarrow & & p \downarrow & & q \downarrow & & \\
0 & \longrightarrow & U & \xrightarrow{\theta} & W & \xrightarrow{\phi} & T \longrightarrow 0.
\end{array}$$

Since ϕ is an approximation, $q = \phi h$ for some $h : M^1 \rightarrow W$. Then $\phi(p - hc) = 0$. Thus $p - hc = \theta \ell$ for some $\ell : M^0 \rightarrow U$. Then $\theta(r - \ell f) = 0$, so since θ is mono, $r = \ell f$. Thus $u \in \text{Im}(\ell)$. Repeating for a basis of U , we get a map from a direct sum of copies of M^0 onto U , so $U \in \mathcal{T}$.

(v) Follows. $\square$

Corollary. *If M is a τ -rigid module, then $\text{gen } M$ is a functorially finite torsion class. Conversely, any functorially finite torsion class $\mathcal{T}$ is of the form $\text{gen } M$ for some τ -rigid module M , which we can take to be the direct sum of the indecomposable Ext-projectives in $\mathcal{T}$.*

Proof. Any torsion class in $A\text{-mod}$ is contravariantly finite, since the inclusion has a right adjoint. Recall also that if M is a module, then $\text{gen } M$ is always covariantly finite by the proposition at the end of section 1.9. In particular, if M is τ -rigid, then $\text{gen } M$ is a functorially finite torsion class.

The last part follows from the theorem, since up to multiplicities, $M^0 \oplus M^1$ is the direct sum of the indecomposable Ext-projectives in $\mathcal{T}$. $\square$

There is a better description of the Ext-injectives in a torsion class.

Remark. If $\mathcal{C}$ is a module class in $A\text{-mod}$, we write $I = \text{ann}(\mathcal{C})$ for the ideal of all $a \in A$ annihilating all modules in $\mathcal{C}$. Then we can consider $\mathcal{C}$ as a module class in $A/I\text{-mod}$. Since A is finite-dimensional, some module in $\mathcal{C}$ is a faithful module for A/I . Thus A/I embeds in some module in $\mathcal{C}$.

Lemma. *If $(\mathcal{T}, \mathcal{F})$ is a torsion theory in $A\text{-mod}$, then*

- (i) *The Ext-injectives for $\mathcal{T}$ are the injective $A/\text{ann}(\mathcal{T})$ -modules.*
- (ii) *The Ext-projectives for $\mathcal{F}$ are the projective $A/\text{ann}(\mathcal{F})$ -modules.*

Proof. (i) Let $I = \text{ann}(\mathcal{T})$. Any injective A/I -module E has an epi $(A/I)^n \rightarrow E$. Now A/I embeds in some module $T \in \mathcal{T}$, and by the injective property the epi extends to an epi $T^n \rightarrow E$. Thus $E \in \mathcal{T}$.

Now if U is an Ext-injective, it embeds in an injective A/I -module, say $0 \rightarrow U \rightarrow E \rightarrow E/U \rightarrow 0$. Then $E/U \in \mathcal{T}$, so this sequence splits, so U is injective as an A/I -module.

(ii) is dual. $\square$

5.2 Tilting modules

Definition. Let M be an A -module.

M is a *partial tilting module* if $\text{proj. dim } M \leq 1$ and $\text{Ext}^1(M, M) = 0$.

A partial tilting module M is a *tilting module* if there is an exact sequence $0 \rightarrow A \rightarrow M^0 \rightarrow M^1 \rightarrow 0$ with $M^i \in \text{add } M$. (Later we will see that it is equivalent that $\#M = \#A$.)

M is a *partial cotilting module* if $\text{inj. dim } M \leq 1$ and $\text{Ext}^1(M, M) = 0$.

A partial cotilting module is a *cotilting module* if there is an exact sequence $0 \rightarrow M_1 \rightarrow M_0 \rightarrow DA \rightarrow 0$ with $M_i \in \text{add } M$. (Again, it is equivalent that $\#M = \#A$.)

Clearly M is a (partial) tilting A -module iff DM is a (partial) cotilting A^{op} -module.

Note that we deal only with *classical* tilting theory. There is a version allowing higher projective dimension.

Lemma. *If M is a partial tilting module, then M is τ -rigid. Conversely if M is τ -rigid, then it is a partial tilting module for $A/\text{ann}(M)$.*

Proof. Use the AR formula $D\text{Ext}^1(M, N) \cong \overline{\text{Hom}}(N, \tau M)$. If $\text{proj. dim } M \leq 1$ then $\text{Hom}(DA, \tau M) = 0$ by Lemma (2) in §2.2, so the AR formula takes the form $D\text{Ext}^1(M, N) \cong \text{Hom}(N, \tau M)$. The converse is the special case $\mathcal{T} = \text{gen } M$ of (i) in the next lemma. $\square$

Lemma. (i) *If $\mathcal{T}$ is a torsion class in $A\text{-mod}$, then any Ext-projective M in $\mathcal{T}$ is a partial tilting module for $A/\text{ann}(\mathcal{T})$.*

(ii) *If furthermore $\mathcal{T}$ is functorially finite, then the direct sum of all indecomposable Ext-projectives is a tilting module for $A/\text{ann}(\mathcal{T})$.*

Proof. (i) Consider $\mathcal{T}$ as a module class in $A/I\text{-mod}$, where $I = \text{ann}(\mathcal{T})$. Clearly $\text{Ext}_{A/I}^1(M, M) = 0$. Also the injective A/I -modules are in $\mathcal{T}$, and $\tau_{A/I}M$ is in the corresponding torsion-free class, so $\text{Hom}(D(A/I), \tau_{A/I}M) = 0$. Thus by Lemma (2) in §2.2, $\text{proj. dim}_{A/I} M \leq 1$.

(ii) If $\mathcal{T}$ is functorially finite, in the theorem of Auslander-Smalø, the map $f : A \rightarrow M^0$ induces an injection $A/I \rightarrow M^0$, so $M^0 \oplus M^1$ is a tilting module for A/I . $\square$

Proposition (Bongartz). *Let M be a partial tilting module. Take a basis of $\xi_1, \dots, \xi_n$ of $\text{Ext}^1(M, A)$, consider the tuple $(\xi_1, \dots, \xi_n)$ as an element of $\text{Ext}^1(M^n, A)$, and let*

$$0 \rightarrow A \rightarrow E \rightarrow M^n \rightarrow 0.$$

be the corresponding universal extension. Then $T = E \oplus M$ is a tilting module. Thus every partial tilting module is a direct summand of a tilting module, and by duality every partial cotilting module is a direct summand of a cotilting module.

Proof. The long exact sequence for $\text{Hom}(M, -)$ gives

$$\text{Hom}(M, M^n) \xrightarrow{\xi} \text{Ext}^1(M, A) \rightarrow \text{Ext}^1(M, E) \rightarrow \text{Ext}^1(M, M^n),$$

the map ξ is onto, and $\text{Ext}^1(M, M^n) = 0$, so $\text{Ext}^1(M, E) = 0$. From the long exact sequence for $\text{Hom}(-, M)$ one gets $\text{Ext}^1(E, M) = 0$, from the long exact sequence for $\text{Hom}(-, E)$ one gets $\text{Ext}^1(E, E) = 0$. Also A and M^n have projective dimension ≤ 1 , hence so does E . $\square$

A partial tilting module M is τ -rigid, so gives a torsion theory $(\text{gen } M, M^{\perp 0})$. Moreover $\text{gen}_1 M \subseteq \text{gen } M \subseteq M^{\perp 1}$.

Proposition (1). *For a partial tilting module M , the following are equivalent:*

- (i) M is a tilting module.
- (ii) $M^{\perp 0,1} = 0$.
- (iii) $\text{gen } M = M^{\perp 1}$.
- (iv) $\text{gen}_1 M = M^{\perp 1}$.
- (v) X is Ext-projective in $M^{\perp 1} \Leftrightarrow X \in \text{add } M$.

Proof. (i) $\Rightarrow$ (ii). If $X \in M^{\perp 0,1}$, apply $\text{Hom}(-, X)$ to the exact sequence $0 \rightarrow A \rightarrow M^0 \rightarrow M^1 \rightarrow 0$, to deduce that $\text{Hom}(A, X) = 0$.

(ii) $\Rightarrow$ (iii). Suppose $X \in M^{\perp 1}$. Take a basis of $\text{Hom}(M, X)$ and use it to form the universal map $f : M^n \rightarrow X$. Then $\text{Im } f \in \text{gen } M$. Consider the exact sequence $0 \rightarrow \text{Im } f \rightarrow X \rightarrow X/\text{Im } f \rightarrow 0$. Apply $\text{Hom}(M, -)$ giving an exact sequence

$$0 \rightarrow \text{Hom}(M, \text{Im } f) \rightarrow \text{Hom}(M, X) \rightarrow \text{Hom}(M, X/\text{Im } f) \rightarrow \text{Ext}^1(M, \text{Im } f).$$

By construction the map $\text{Hom}(M, M^n) \rightarrow \text{Hom}(M, X)$ is onto, hence so is the map $\text{Hom}(M, \text{Im } f) \rightarrow \text{Hom}(M, X)$. Also $\text{Ext}^1(M, \text{Im } f) = 0$ since M is τ -rigid. Thus $\text{Hom}(M, X/\text{Im } f) = 0$. Also $\text{Ext}^1(M, X/\text{Im } f) = 0$. Thus $X/\text{Im } f \in M^{\perp 0,1}$. Thus $X/\text{Im } f = 0$, so f is onto, so $X \in \text{gen } M$.

(iii) $\Rightarrow$ (iv). Suppose $X \in M^{\perp 1}$. Then it is in $\text{gen } M$. Let L be the kernel of the universal map $M^n \rightarrow X$. Then applying $\text{Hom}(M, -)$ we see that $L \in M^{\perp 1}$, so $L \in \text{gen } M$. Say $M'' \twoheadrightarrow L$. Now the sequence $M'' \rightarrow M^n \rightarrow X \rightarrow 0$ shows that $X \in \text{gen}_1 M$.

(iv) $\Rightarrow$ (v). Clearly M and so any $X \in \text{add}(M)$ is in $M^{\perp 1}$ and Ext-projective. Conversely if X is in $M^{\perp 1}$ and Ext-projective, then by (iv) there is an exact sequence $M'' \xrightarrow{f} M' \rightarrow X \rightarrow 0$. This gives an exact sequence $0 \rightarrow \text{Im } f \rightarrow M' \rightarrow X \rightarrow 0$ with $\text{Im } f \in \text{gen } M \subseteq M^{\perp 1}$. By assumption this sequence splits, so $X \in \text{add } M$.

(v) $\Rightarrow$ (i). It suffices to show that E in Bongartz's sequence is in $\text{add } M$, and for this it suffices to show it is Ext-projective in $M^{\perp 1}$. We know it is in $M^{\perp 1}$. If $Y \in M^{\perp 1}$, apply $\text{Hom}(-, Y)$ to the Bongartz sequence to get $\text{Ext}^1(M^n, Y) \rightarrow \text{Ext}^1(E, Y) \rightarrow \text{Ext}^1(A, Y)$, so $\text{Ext}^1(E, Y) = 0$. $\square$

Dually, a partial cotilting module M is τ^- -rigid, so gives a torsion theory $({}^{\perp 0}M, \text{cogen } M)$. Moreover $\text{cogen}^1 M \subseteq \text{cogen } M \subseteq {}^{\perp 1}M$. The following is dual to the last proposition.

Proposition (2). *For a partial cotilting module M , the following are equivalent:*

- (i') M is a cotilting module.
- (ii') ${}^{\perp 0,1}M = 0$.
- (iii') $\text{cogen } M = {}^{\perp 1}M$.
- (iv') $\text{cogen}^1 M = {}^{\perp 1}M$.
- (v') X is Ext-injective in ${}^{\perp 1}M \Leftrightarrow X \in \text{add } M$.

Proposition (3). *If ${}_A M$ is a (co)tilting module, then it is f.b. and if $B = \text{End}_A(M)$, then ${}_B M$ is also a (co)tilting module.*

Proof. If ${}_A M$ is tilting, then $\text{gen}_1 M = M^{\perp 1}$, which contains DA , so ${}_A M$ is f.b.

(i) Applying $\text{Hom}_A(-, M)$ to the exact sequence $0 \rightarrow A \rightarrow M^0 \rightarrow M^1 \rightarrow 0$ gives

$$0 \rightarrow \text{Hom}_A(M^1, M) \rightarrow \text{Hom}_A(M^0, M) \rightarrow M \rightarrow 0$$

and $\text{Hom}_A(M^i, M) \in \text{add}(\text{Hom}_A(M, M)) = \text{add}({}_B B)$, so $\text{proj. dim } {}_B M \leq 1$.

(ii) The tilting sequence $0 \rightarrow A \rightarrow M^0 \rightarrow M^1 \rightarrow 0$ stays exact on applying $\text{Hom}(-, M)$. Thus $A \in \text{cogen}^2({}_A M)$. Thus $\text{Ext}_B^1(M, M) = 0$ by the proposition about endomorphism correspondents.

(iii) Applying $\text{Hom}_A(-, M)$ to a projective resolution $0 \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$ of M gives an exact sequence

$$0 \rightarrow B \rightarrow M^0 \rightarrow M^1 \rightarrow 0$$

where $M^i = \text{Hom}_A(P_i, M) \in \text{add}({}_B M)$. Thus ${}_B M$ is a tilting module.

Dually for cotilting. □

5.3 The Brenner-Butler Theorem

Let ${}_A M$ be a cotilting module and $B = \text{End}_A(M)$, so ${}_B M$ is also cotilting.

In $A\text{-mod}$ we have a torsion theory $(\mathcal{T}_A, \mathcal{F}_A) = ({}^{\perp 0}{}_A M, \text{cogen } {}_A M)$. Since ${}_A M$ is cotilting we have

$$\mathcal{F}_A = \text{cogen}({}_A M) = \text{cogen}^1({}_A M) = {}^{\perp 1}{}_A M = \{X \in A\text{-mod} : \text{Ext}_A^1(X, M) = 0\}.$$

In $B\text{-mod}$ we have a torsion theory $(\mathcal{T}_B, \mathcal{F}_B) = ({}^{\perp 0}{}_B M, \text{cogen } {}_B M)$. Since ${}_B M$ is cotilting we have the equivalent alternative descriptions of $\mathcal{F}_B$.

Theorem (Brenner-Butler Theorem, 1st version). *There are antiequivalences*

$$\mathcal{F}_A \begin{array}{c} \xrightarrow{\text{Hom}_A(-, M)} \\ \xleftarrow{\text{Hom}_B(-, M)} \end{array} \mathcal{F}_B \quad \text{and} \quad \mathcal{T}_A \begin{array}{c} \xrightarrow{\text{Ext}_A^1(-, M)} \\ \xleftarrow{\text{Ext}_B^1(-, M)} \end{array} \mathcal{T}_B.$$

Proof. Since $\mathcal{F}_A = \text{cogen}^1({}_A M)$ and $\mathcal{F}_B = \text{cogen}^1({}_B M)$, the first antiequivalence is given by endomorphism correspondence.

Given a module ${}_A X$ in $\mathcal{T}_A$, so with $\text{Hom}_A(X, M) = 0$, we show that

$$\text{Hom}_B(\text{Ext}_A^1(X, M), M) = 0$$

and construct a natural isomorphism

$$X \rightarrow \text{Ext}_B^1(\text{Ext}_A^1(X, M), M).$$

Indeed, take a projective cover of X to get a sequence $0 \rightarrow L \rightarrow P \rightarrow X \rightarrow 0$. It gives an exact sequence of B -modules

$$0 \rightarrow \text{Hom}_A(P, M) \rightarrow \text{Hom}_A(L, M) \rightarrow \text{Ext}_A^1(X, M) \rightarrow 0$$

Now $P, L \in \text{cogen } M = \text{cogen}^1 M$, so the natural maps $P \rightarrow \text{Hom}_B(\text{Hom}_A(P, M), M)$ and $L \rightarrow \text{Hom}_B(\text{Hom}_A(L, M), M)$ are isomorphisms. Also


$$\text{Hom}_A(L, M) \in \text{cogen}^1({}_B M) = {}^{\perp 1}({}_B M),$$

so $\text{Ext}_B^1(\text{Hom}_A(L, M), M) = 0$. Thus we get a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & L & \longrightarrow & P & \longrightarrow & X & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & & & \\ 0 & \longrightarrow & ({}^1(X, M), M) & \longrightarrow & ((L, M), M) & \longrightarrow & ((P, M), M) & \longrightarrow & {}^1({}^1(X, M), M) \longrightarrow 0 \end{array}$$

(where we omit the words Hom and Ext) with exact rows and in which the vertical maps are isomorphisms. Thus $\text{Hom}_B(\text{Ext}_A^1(X, M), M) = 0$ and there is an induced isomorphism $X \rightarrow \text{Ext}_B^1(\text{Ext}_A^1(X, M), M)$. One also needs to show that this is a natural isomorphism, but we omit the proof of this. $\square$

Example.



Theorem. If B is hereditary, then the torsion theory $(\mathcal{T}_A, \mathcal{F}_A)$ is split (and by symmetry, if A is hereditary, then $(\mathcal{T}_B, \mathcal{F}_B)$ is split).

Proof. We want to show that $\text{Ext}_A^1(U, V) = 0$ for all $U \in \mathcal{F}_A$ and $V \in \mathcal{T}_A$. Now we have $V = \text{Ext}_B^1(Y, M)$ for some $Y \in \mathcal{T}_B$. Taking a projective A -module Q mapping

onto U , gives an exact sequence

$$0 \rightarrow \Omega_1 U \rightarrow Q \rightarrow U \rightarrow 0$$

and applying $\text{Hom}_A(-, M)$, we get an isomorphism $\text{Ext}^2(U, M) \cong \text{Ext}^1(\Omega_1 U, M)$ (dimension shifting). Also $Q \in \text{cogen } M$ since M is faithful, so $\Omega_1 U \in \text{cogen } M = \mathcal{F}_A$, so $\text{Ext}^1(\Omega_1 U, M) = 0$, so $\text{Ext}^2(U, M) = 0$. We also have $\text{Ext}^1(U, M) = 0$.

Now take a projective resolution

$$0 \rightarrow P_1 \rightarrow P_0 \rightarrow Y \rightarrow 0.$$

Since $\text{Hom}_B(Y, M) = 0$, we get an exact sequence

$$0 \rightarrow \text{Hom}_B(P_0, M) \rightarrow \text{Hom}_B(P_1, M) \rightarrow \text{Ext}_B^1(Y, M) \rightarrow 0.$$

Thus

$$\cdots \rightarrow \text{Ext}_A^1(U, \text{Hom}_B(P_1, M)) \rightarrow \text{Ext}_A^1(U, \text{Ext}_B^1(Y, M)) \rightarrow \text{Ext}_A^2(U, \text{Hom}_B(P_0, M)) \rightarrow \cdots$$

Now $\text{Hom}_B(P_i, M) \in \text{add}({}_A M)$, so the outer terms are zero, giving the result. $\square$

We now give another version of the Brenner-Butler theorem. Let A be an algebra and ${}_A T$ a tilting module. Let $B = \text{End}(T)^{op}$, so T becomes an A - B -bimodule, and T_B is right B -module which is a tilting module. Thus DT is a left B -module which is cotilting.

The tilting module ${}_A T$ gives a torsion theory $(\mathcal{T}, \mathcal{F})$ in $A\text{-mod}$ via

$$\begin{aligned} \mathcal{T} &= \text{gen } {}_A T = ({}_A T)^{\perp 1} \\ \mathcal{F} &= ({}_A T)^{\perp 0}. \end{aligned}$$

The cotilting left B -module DT gives a torsion theory $(\mathcal{X}, \mathcal{Y})$ in $B\text{-mod}$ where

$$\mathcal{X} = {}^{\perp 0}({}_B DT)$$

and

$$\mathcal{Y} = \text{cogen } {}_B DT = {}^{\perp 1}({}_B DT).$$

Note that if

$$\cdots \rightarrow P_1 \rightarrow P_0 \rightarrow T \rightarrow 0$$

is a projective resolution of T as a right B -module, then

$$0 \rightarrow DT \rightarrow DP_0 \rightarrow DP_1 \rightarrow \cdots$$

is an injective resolution of DT as a left B -module. Now if Y is a left B -module, then $\text{Tor}_n^B(T, Y)$ is the homology of the complex $P_* \otimes_B Y$, so $D(\text{Tor}_n^B(T, Y))$ is the cohomology of the complex $D(P_* \otimes_B Y) \cong \text{Hom}_B(Y, DP_*)$, so

$$D(\text{Tor}_n^B(T, Y)) \cong \text{Ext}_B^n(Y, DT).$$


Thus $\mathcal{X} = \{Y : T \otimes_B Y = 0\}$ and $\mathcal{Y} = \{Y : \text{Tor}_1^B(T, Y) = 0\}$.

Theorem (Brenner-Butler theorem, 2nd version). *We have inverse equivalences*

$$\mathcal{T} \begin{array}{c} \xrightarrow{\text{Hom}_A(T, -)} \\ \xleftarrow{T \otimes_B -} \end{array} \mathcal{Y}$$

and

$$\mathcal{F} \begin{array}{c} \xrightarrow{\text{Ext}_A^1(T, -)} \\ \xleftarrow{\text{Tor}_1^B(T, -)} \end{array} \mathcal{X}.$$



For the proof, we consider DT as a cotilting right A -module, so as a cotilting left A^{op} -module, and $B = \text{End}_{A^{op}}(DT)$. Use this in 1st version, and compose with duality.

Examples. (1) The Bernstein-Gelfand-Ponomarev reflection functors fit this picture. If i is a sink in Q , the tilting module is

$$T = \tau^{-1}P[i] \oplus \bigoplus_{j \neq i} P[j].$$

In fact, for any algebra A , if $P[i]$ is a simple projective (and not injective), this construction gives a tilting module, called an APR tilting module after Auslander, Platzeck and Reiten, Coxeter functors without diagrams, 1979.

(2) A *tilted algebra* is one of the form $B = \text{End}_A(T)$ where A is hereditary and ${}_A T$ is a tilting module. Then the torsion theory $(\mathcal{X}, \mathcal{Y})$ is split.

(3) A *concealed algebra* is a tilted algebra of the form $B = \text{End}_A(T)$ where A is representation-infinite connected hereditary and ${}_A T$ is a preprojective tilting module.

There is some $n > 0$ with $\tau^n T = 0$. If X is a module with $X \cong \tau^{-(n-1)} \tau^{n-1} X$, for example if X is indecomposable and not preprojective, or not near the start of the preprojective component, then since A is hereditary,

$$\text{Ext}^1(T, X) \cong D \text{Hom}(X, \tau T) \cong D \text{Hom}(\tau^{-(n-1)} \tau^{n-1} X, \tau T)$$

$$\cong D \operatorname{Hom}(\tau^{n-1}X, \tau^n T) = 0$$

so $X \in \mathcal{T}$. Thus $\mathcal{T}$ contains all but finitely many indecomposables and $\mathcal{F}$ contains only finitely many indecomposables.

Then $B\text{-mod}$ is obtained by reassembling these two pieces as $\mathcal{X}$ and $\mathcal{Y}$.

There is an example worked out in detail on p336 of Assem, Simson and Skowronski, Elements of the representation theory of associative algebras I.

A theorem of Happel and Vossieck, Minimal algebras of infinite representation type with preprojective component, Manuscripta Math. 1983: If B is an algebra with a preprojective component and B is minimal of infinite representation type, meaning that B/BeB of finite representation type for all nonzero idempotents e , then either B is Morita equivalent to the path algebra of an r -arrow Kronecker quiver with $r \geq 2$, or B is tame concealed, and there is a classification of all such algebras.

5.4 Derived equivalences

I promised to talk about how tilting theory is related to derived categories, but to do this properly would be too much of a digression. So I will only sketch things briefly.

Definition. An A -module T is a *generalized (or Miyashita) tilting module* if

- (i) $\operatorname{proj. dim} T < \infty$, so there is a projective resolution $0 \rightarrow P_r \rightarrow \cdots \rightarrow P_0 \rightarrow T \rightarrow 0$
- (ii) $\operatorname{Ext}^i(T, T) = 0$ for all $i > 0$
- (iii) There is an exact sequence $0 \rightarrow A \rightarrow T^0 \rightarrow \cdots \rightarrow T^r \rightarrow 0$ with $T^i \in \operatorname{add}(T)$.

The following was proved by Happel for $\operatorname{gl. dim} A < \infty$ and in general by Cline, Parshall and Scott, Derived categories and Morita theory, J. Algebra 1986.

Theorem. *Let T be a generalized tilting A -module. Then T is faithfully balanced, and letting $B = \operatorname{End}(T)^{op}$, the module T_B is a generalized tilting right B -module. Moreover T induces inverse equivalences of triangulated categories*

$$D^b(A\text{-mod}) \xrightleftharpoons{\quad} D^b(B\text{-mod})$$

The functor to the right is $\mathbf{R} \operatorname{Hom}(T, -)$, the right derived functor of $\operatorname{Hom}(T, -)$. This can be defined abstractly, but to show it exists and compute it, one uses the isomorphisms

$$D^b(A\text{-mod}) \cong D^{+,b}(A\text{-mod}) \cong K^{+,b}(A\text{-inj}).$$

Then $\operatorname{Hom}(T, -)$ can be applied to a complex of injectives I , giving a complex $\operatorname{Hom}(T, I)$ in $D^+(B\text{-mod})$.

Now if X is an A -module in degree 0, then $\mathbf{R}\mathrm{Hom}(T, X)$ is computed by taking an injective resolution of X , so its n -th cohomology is $\mathrm{Ext}^n(T, X)$. This is nonzero only for finitely many n , so $\mathbf{R}\mathrm{Hom}(T, X) \in D^{+,b}(B\text{-mod})$. Now any complex X in $D^b(A\text{-mod})$ can be built from modules in a finite number of degrees. Thus $\mathbf{R}\mathrm{Hom}(T, X) \in D^{+,b}(B\text{-mod}) \cong D^b(B\text{-mod})$.

Similarly the functor to the left is $\mathbf{L}T \otimes_B -$, constructed using

$$D^b(A\text{-mod}) \cong D^{-,b}(A\text{-mod}) \cong K^{-,b}(A\text{-proj}).$$

Using that $T \otimes_B -$ is left adjoint to $\mathrm{Hom}_A(T, -)$ one can show that $\mathbf{L}T \otimes_B -$ is left adjoint to $\mathbf{R}\mathrm{Hom}_A(T, -)$. Then one can show that they are inverse equivalences.

Now suppose A is hereditary. Then every object in $D^b(A\text{-mod})$ is a direct sum of stalk complexes - living in only one degree.

The shift $X[n]$ of a complex X is given by $X[n]^i = X^{i+n}$ and it multiplies the differential by $(-1)^i$.

Thus if X is an A -module considered as a complex in degree 0, then $X[n]$ is a module in degree $-n$.

Also $\mathrm{Hom}(X[i], Y[j]) \cong \mathrm{Ext}^{j-i}(X, Y)$ which is zero for $j < i$.

Thus we can picture $D^b(A\text{-mod})$ as below.

Now suppose in addition that T is a classical tilting module, so B is tilted.

If X is an A -module in degree 0, then it is isomorphic in the derived category to its injective resolution, and $\mathbf{R}\mathrm{Hom}(T, X)$ is the complex

$$\cdots \rightarrow 0 \rightarrow \mathrm{Hom}(T, I^0) \rightarrow \mathrm{Hom}(T, I^1) \rightarrow 0 \rightarrow \cdots$$


The cohomology in degree i is $\mathrm{Ext}^i(T, X)$.

If $X \in \mathcal{T} = ({}_A T)^\perp$ then $\mathbf{R}\mathrm{Hom}(T, X)$ is in $B\text{-mod}$. It is in the class $\mathcal{X} = \{Y : \mathrm{Tor}_1^B(T, Y) = 0\}$.

If $X \in \mathcal{F} = ({}_A T)^\perp$ then $\mathbf{R}\mathrm{Hom}(T, X)$ is a module in degree 1, so it is in $B\text{-mod}[-1]$. It is in the shift of $\mathcal{Y} = \{Y : T \otimes_B Y = 0\}$.

Since the torsion theory $(\mathcal{X}, \mathcal{Y})$ is splitting, we can picture $D^b(B\text{-mod})$ as fol-

lows.



5.5 Consequences for Grothendieck groups

Definition. We consider two types of *Grothendieck groups*.

If $\mathcal{A}$ is an abelian category, the Grothendieck group $G_0(\mathcal{A})$ is the additive group generated by symbols $[X]$ for each object X in $\mathcal{A}$, modulo the relations $[Y] = [X] + [Z]$ for any short exact sequence $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$.

If $\mathcal{C}$ is an additive category, the Grothendieck group $K_0(\mathcal{C})$ is the additive group generated by symbols $[X]$ for each object X in $\mathcal{C}$, modulo the relations $[Y] = [X] + [Z]$ whenever $Y \cong X \oplus Z$.

Lemma. If A is a f.d. algebra with simples $S[i]$ ($i = 1, \dots, n$) and indecomposable projectives $P[i]$, then:

(i) The map sending a module X to its dimension vector gives an isomorphism $G_0(A\text{-mod}) \cong \mathbb{Z}^n$, $[X] \mapsto \underline{\dim} X$, so $G_0(A\text{-mod})$ is the free $\mathbb{Z}$ -module on the symbols $[S[i]]$.

(ii) $K_0(A\text{-proj})$ is also isomorphic to $\mathbb{Z}^n$ since it is the free $\mathbb{Z}$ -module on the symbols $[P[i]]$.

Proof. (i) is the Jordan-Hölder theorem and (ii) is Krull-Remak-Schmidt. $\square$

Theorem. If ${}_A M$ is a cotilting module and $B = \text{End}_A(M)$, then there is an isomorphism

$$\theta : G_0(A\text{-mod}) \rightarrow G_0(B\text{-mod}), \quad [X] \mapsto [\text{Hom}_A(X, M)] - [\text{Ext}_A^1(X, M)].$$

Thus the canonical basis of $G_0(A\text{-mod})$ gives a new basis of $G_0(B\text{-mod})$, hence the name “tilting”.

Proof. If we apply $\text{Hom}_A(-, M)$ to a short exact sequence of A -modules, say $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ we get a long exact sequence of B -modules

$$0 \rightarrow \text{Hom}(Z, M) \rightarrow \text{Hom}(Y, M) \rightarrow \text{Hom}(X, M) \rightarrow$$

$$\text{Ext}^1(Z, M) \rightarrow \text{Ext}^1(Y, M) \rightarrow \text{Ext}^1(X, M) \rightarrow 0.$$

Now the relations for $G_0(B\text{-mod})$ imply that

$$\begin{aligned} \theta([Y]) &= [\text{Hom}_A(Y, M)] - [\text{Ext}_A^1(Y, M)] \\ &= [\text{Hom}_A(X, M)] - [\text{Ext}_A^1(X, M)] + [\text{Hom}_A(Z, M)] - [\text{Ext}_A^1(Z, M)] \\ &= \theta([X]) + \theta([Z]) \end{aligned}$$

so that θ is well-defined.

Swapping the roles of A and B there is a map ϕ in the reverse direction.

If $X \in \text{cogen } M$ or $X \in {}^{\perp 0}M$, then $\phi(\theta([X])) = [X]$. Because any X belongs to a short exact sequence whose ends are torsion and torsion-free, it follows that $\phi\theta = 1$. Similarly $\theta\phi = 1$. $\square$

Recall that we write $\#M$ for the number of isomorphism classes of indecomposable summands of M . Thus $\#A$ is the number of isomorphism classes of indecomposable projective A -modules, so the number of isomorphism classes of simple A -modules.

Corollary. *Any partial (co)tilting module M has $\#M \leq \#A$, with equality if and only if M is (co)tilting.*

Proof. If ${}_A M$ is a cotilting module and $B = \text{End}_A(M)$, then $\text{Hom}(-, M)$ gives an antiequivalence between $\text{add } M$ and $B\text{-proj}$, so $\#M$ is the rank of $G_0(B\text{-mod})$, which is the rank of $G_0(B\text{-mod})$, which is $\#A$.

By duality any tilting module has $\#A$ summands. By Bongartz, any partial tilting module is a summand of a tilting module. $\square$

Theorem (Smalø, 1984). *If $(\mathcal{T}, \mathcal{F})$ is a torsion theory, then $\mathcal{T}$ is functorially finite iff $\mathcal{F}$ is functorially finite.*

Proof. By symmetry, it suffices to prove that if $\mathcal{T}$ is functorially finite, then so is $\mathcal{F}$.

The number of indec Ext-injectives in $\mathcal{F}$
 $=$ number of indec injectives in $\mathcal{F}$ + number of non-injective indec Ext-injectives in $\mathcal{F}$
 $=$ number of indec injectives in $\mathcal{F}$ + number of non-projective indec Ext-projectives in $\mathcal{T}$

= number of indec injectives in $\mathcal{F}$ + number of indec Ext-projectives in $\mathcal{T}$ - number of indec projectives in $\mathcal{T}$.

Since $\mathcal{T}$ is functorially finite, the direct sum of the indecomposable Ext-projectives in $\mathcal{T}$ is a tilting module for $A/\text{ann}(\mathcal{T})$. Thus we get

= number of indec injectives in $\mathcal{F}$ + $\#A/\text{ann}(\mathcal{T})$ - number of indec projectives in $\mathcal{T}$.

Now the number of indecomposable injectives $I[i]$ not in $\mathcal{F}$ is the number of indecomposable Ext-injectives in $\mathcal{T}$, which is the number of indecomposable injective $A/\text{ann}(\mathcal{T})$ -modules, so it is $\#A/\text{ann}(\mathcal{T})$.

Similarly the number of indecomposable projectives not in $\mathcal{T}$ is $\#A/\text{ann}(\mathcal{F})$.

So we get

$$= (\#A - \#A/\text{ann}(\mathcal{T})) + \#A/\text{ann}(\mathcal{T}) - (\#A - \#A/\text{ann}(\mathcal{F})).$$

$$= \#A/\text{ann}(\mathcal{F}).$$

Now by the dual of an earlier result, any Ext-injective in $\mathcal{F}$ is a partial cotilting module for $A/\text{ann}(\mathcal{F})$. Thus the direct sum M of all indecomposable Ext-injectives in $\mathcal{F}$ is a cotilting module for $A/\text{ann}(\mathcal{F})$.

Thus working in $A/\text{ann}(\mathcal{F})\text{-mod}$, we have $\text{cogen } M = {}^{\perp 1}M$. Now since M is Ext-injective in $\mathcal{F}$, $\text{Ext}^1(\mathcal{F}, M) = 0$, so $\mathcal{F} \subseteq {}^{\perp 1}M = \text{cogen } M \subseteq \mathcal{F}$.

Thus also $\mathcal{F} = \text{cogen } M$ as a module class in $A\text{-mod}$. Thus $\mathcal{F}$ is contravariantly finite by the proposition at the end of §1.9, and it is covariantly finite since the inclusion has a left adjoint. $\square$

5.6 Some tau-tilting theory

It was started by Adachi, Iyama and Reiten, τ -tilting theory, 2014, although there was earlier work, see Derksen and Fei, General Presentations of Algebras and Foundations of tau-tilting Theory, arxiv 2409.12743. It has led to a lot of other work. We shall only do a little.

We have done all the necessary prerequisites in our theorems about functorially finite torsion and torsion-free classes.

Lemma. (i) If M is an A -module, then $\#A/\text{ann}(M)$ is the number of different simple composition factors involved in M .

(ii) If M is τ -rigid, then the number of indecomposable Ext-projectives in $\text{gen } M$ is $\#A/\text{ann}(M)$ and $\#M \leq \#A/\text{ann}(M)$.

Proof. (i) If S is involved in M , then S must be an $A/\text{ann}(M)$ -module. On the other hand, M is faithful as an $A/\text{ann}(M)$ -module, so $A/\text{ann}(M)$ embeds in a direct sum of copies of M , so if S is a simple for $A/\text{ann}(M)$, then it must be a composition factor of M .

(ii) M is a partial tilting module for $A/\text{ann}(M)$, and $\text{gen } M$ is a functorially finite torsion class, so the direct sum of the indecomposable Ext-projectives is a tilting module for $A/\text{ann}(M)$. $\square$

Definition. Let M be a τ -rigid A -module.

(i) M is a *support τ -tilting module* if $\#M = \#A/\text{ann}(M)$, or equivalently M is the direct sum of the indecomposable Ext-projectives in $\text{gen } M$, each with non-zero multiplicity.

(ii) M is a *τ -tilting module* if it is a sincere support τ -tilting module, or equivalently $\#M = \#A$. (Recall that *sincere* means that every simple module occurs as a composition factor.)

Lemma. *If M is τ -rigid, then $\mathcal{T} = {}^{\perp 0}(\tau M)$ is a sincere functorially finite torsion class. If T is the direct sum of the indecomposable Ext-projectives in $\mathcal{T}$, then T is a τ -tilting module, $M \in \text{add}(T)$ and ${}^{\perp 0}(\tau T) = \text{gen } T$.*

The module T is called the *Bongartz completion* of M .

Proof. τM is a τ^- -rigid module, so we get a torsion theory $({}^{\perp 0}(\tau M), \text{cogen } \tau M)$. The torsion class is functorially finite by Smalø's theorem. The torsion class is sincere, since no injective $I[i]$ embeds in τM , so $I[i]$ is not in the torsion-free class, so its torsion submodule is non-zero, and this has $S[i]$ as a submodule. Clearly $M \in \mathcal{T}$ and it is Ext-projective since if $X \in \mathcal{T}$, then

$$\text{Ext}^1(M, X) \cong D\overline{\text{Hom}}(X, \tau M)$$

and $\text{Hom}(X, \tau M) = 0$. Now

$${}^{\perp 0}(\tau M) = \mathcal{T} = \text{gen } T \subseteq {}^{\perp 0}(\tau T) \subseteq {}^{\perp 0}(\tau M)$$

where the second equality holds by the Auslander-Smalø theorem of functorially finite torsion classes, the first inclusion since T is τ -rigid, and the second since $M \in \text{add}(T)$. $\square$

The following is an analogue of a result known as Wakamatsu's lemma.

Lemma. *If M is τ -rigid and $f : M' \rightarrow X$ is a right $\text{add}(M)$ -approximation of a module X , then $\text{Hom}(\text{Ker}(f), \tau M) = 0$.*

Proof. Replacing X by $\text{Im}(f)$, we may suppose that f is surjective. Applying $\text{Hom}(-, \tau M)$ gives an exact sequence

$$\text{Hom}(M', \tau M) \rightarrow \text{Hom}(\text{Ker}(f), \tau M) \rightarrow \text{Ext}^1(X, \tau M) \rightarrow \text{Ext}^1(M', \tau M).$$

The first hom space is zero since M is τ -rigid. Now the map $\text{Hom}(M, M') \rightarrow \text{Hom}(M, X)$ induced by f is surjective, hence so is the map on $\overline{\text{Hom}}$, hence by the Auslander-Reiten formula, the map on Ext^1 is injective. $\square$

Theorem. *A τ -rigid module M is τ -tilting iff $\text{gen } M = {}^{\perp 0}(\tau M)$.*

Proof. If M is τ -tilting, then its Bongartz completion T can have no new indecomposable summands, so $\text{add}(M) = \text{add}(T)$, and we get the result from the lemma.

Suppose $\text{gen } M = {}^{\perp 0}(\tau M)$. Let T be the Bongartz completion of M . Then

$$\text{gen } M \subseteq \text{gen } T = {}^{\perp 0}\tau T \subseteq {}^{\perp 0}\tau M = \text{gen } M$$

so all are equal. Take a minimal right $\text{add}(M)$ -approximation of T , say $f : M' \rightarrow T$. It is surjective since $T \in \text{gen } M$, so we get an exact sequence

$$0 \rightarrow \text{Ker}(f) \rightarrow M' \rightarrow T \rightarrow 0.$$

By the Wakamatsu-type lemma $\text{Hom}(\text{Ker}(f), \tau M) = 0$. Since ${}^{\perp 0}(\tau M) = {}^{\perp 0}(\tau T)$ we get $\text{Hom}(\text{Ker}(f), \tau T) = 0$. Thus $\text{Ext}^1(T, \text{Ker}(f)) = 0$. Thus the sequence $0 \rightarrow \text{Ker}(f) \rightarrow M' \rightarrow T \rightarrow 0$ splits. Thus $T \in \text{add}(M)$, so M is τ -tilting. $\square$

Corollary. *Any basic τ -rigid module M which is not τ -tilting, is a direct summand of at least two basic support τ -tilting modules.*

Proof. $\text{gen } M$ and ${}^{\perp 0}(\tau M)$ are different functorially finite torsion classes containing M , and we can take the direct sum of the indecomposable Ext-projectives in either. $\square$

Remark. It is useful to consider pairs (M, P) where M is a module, P is a projective module and $\text{Hom}(P, M) = 0$, so that P is a direct sum of $P[i]$ such that $S[i]$ is not a composition factor of M .

We call it a τ -rigid pair if M is τ -rigid.

We call it a support τ -tilting pair if $\#M + \#P = \#A$. Note that we always have $\leq$ for a τ -rigid pair. Also M is necessarily support τ -tilting.

We call a pair basic if M and P are basic.

One can show that any basic τ -rigid pair (M, P) , can be extended to a basic support τ -tilting pair $(M \oplus M', P \oplus P')$, and if $\#M + \#P = \#A - 1$, there are exactly two ways to do it.

Thus we get mutations of support τ -tilting pairs where we remove any one indecomposable summand, and replace it by the other possible extension of that pair.

Such mutations are related to the mutations in cluster algebras.

Remark. There is a natural homomorphism

$$\theta : K_0(A\text{-proj}) \rightarrow G_0(A\text{-mod}), \quad \theta([X]_K) = [X]_G.$$

It is an isomorphism if $\text{gl. dim } A < \infty$. For this, it suffices to see that it is surjective. Now a projective resolution

$$0 \rightarrow P_n \rightarrow \cdots \rightarrow P_0 \rightarrow X \rightarrow 0$$

gives

$$[X]_G = \sum_{i=0}^n (-1)^i [P_i]_G \in \text{Im } \theta.$$

In general, however, it is not an isomorphism.

Instead there is a bilinear map

$$\langle -, - \rangle : K_0(A\text{-proj}) \times G_0(A\text{-mod}) \rightarrow \mathbb{Z}, \quad ([P], [X]) \mapsto \dim \text{Hom}(P, X).$$

and

$$\langle [P[i]], [S[j]] \rangle = \dim \text{Hom}(P[i], S[j]) = \delta_{ij} \dim D_i.$$

where $D_i = \text{End}(S[i])^{op}$. The matrix is invertible over $\mathbb{Q}$, so gives a perfect pairing between $K_0(A\text{-proj}) \otimes_{\mathbb{Z}} \mathbb{Q}$ and $G_0(A\text{-mod}) \otimes_{\mathbb{Z}} \mathbb{Q}$.

Definition. The *g-vector* of a module M is

$$g(M) = [P_0] - [P_1] \in K_0(A\text{-proj})$$

where $P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$ is the minimal projective presentation.

Lemma. If M and X are modules, then

$$\langle g(M), [X] \rangle = \dim \text{Hom}(M, X) - \dim \text{Hom}(X, \tau M).$$

Proof. We have exact sequences

$$0 \rightarrow \tau M \rightarrow \nu P_1 \rightarrow \nu P_0$$

and

$$0 \rightarrow \text{Hom}(M, X) \rightarrow \text{Hom}(P_0, X) \rightarrow \text{Hom}(P_1, X)$$

so we get a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}(X, \tau M) & \longrightarrow & \text{Hom}(X, \nu P_1) & \longrightarrow & \text{Hom}(X, \nu P_0) \\ & & & & \parallel & & \parallel \\ & & & & D \text{Hom}(P_1, X) & \longrightarrow & D \text{Hom}(P_0, X) \longrightarrow D \text{Hom}(M, X) \longrightarrow 0 \end{array}$$

□

Lemma. *If M is τ -rigid, and $P_1 \xrightarrow{\theta} P_0 \xrightarrow{\phi} M \rightarrow 0$ is a minimal projective presentation, then P_0 and P_1 have no direct summand in common.*

Proof. The map $\text{Hom}(P_0, M) \rightarrow \text{Hom}(P_1, M)$ is surjective since its dual can be identified with the map $\text{Hom}(M, \nu P_1) \rightarrow \text{Hom}(M, \nu P_0)$ and the kernel of this map is $\text{Hom}(M, \tau M) = 0$.

It suffices to show that any map $s : P_1 \rightarrow P_0$ is in the radical. The composition $\phi s \in \text{Hom}(P_1, M)$, so by the surjectivity, $\phi s = t\theta$ for some $t : P_0 \rightarrow X$.

Since ϕ is surjective and P_0 is projective, we have $t = \phi u$ for some $u : P_0 \rightarrow P_0$.

Then $\phi(s - u\theta) = 0$. Thus since P_1 is projective, $s - u\theta = \theta v$ for some $v : P_1 \rightarrow P_1$. Now θ is in the radical, hence so is s . $\square$

Theorem. *Two τ -rigid modules with the same g -vector must be isomorphic.*

By the lemma, the two modules have the same projectives in their minimal projective presentations. Thus we are dealing with two homomorphisms in $\text{Hom}(P_1, P_0)$. Can reduce to the case of an algebra over an algebraically closed field. Then it is a simple geometric argument. Hopefully we will do it later.

Remark. There are nice connections with semibricks. See Asai, Semibricks, IMRN 2020 and Ringel, Brick chain filtrations, arxiv 2411.18427

Also Demonet, Iyama and Jasso, tau-tilting finite algebras, bricks, and g -vectors, IMRN 2019. For example the following are equivalent.

- A has only finitely many τ -tilting modules.
- Every torsion class in $A\text{-mod}$ is functorially finite.
- A has only finitely many bricks.

6 Varieties and schemes of algebras and modules

6.1 Varieties of algebras

First we need to discuss varieties. We work over an algebraically closed field K , and follow Kempf, Algebraic varieties, 1993.

Definition. A *space with functions* consists of a topological space X and the assignment of a set $\mathcal{O}(U)$ of *regular functions* for each open set $U \subseteq X$, satisfying:

(a) $\mathcal{O}(U)$ is a K -subalgebra of the algebra of all functions $U \rightarrow K$, with point-wise operations.

(b) If U is a union of open sets, $U = \bigcup U_\alpha$, then $f \in \mathcal{O}(U)$ iff $f|_{U_\alpha} \in \mathcal{O}(U_\alpha)$ for all α .

(c) If $f \in \mathcal{O}(U)$, then $D(f) = \{u \in U \mid f(u) \neq 0\}$ is open in U and $1/f \in \mathcal{O}(D(f))$.

A *morphism* of spaces with functions is a continuous map $\theta : X \rightarrow Y$ with the property that for any open subset U of Y , and any $f \in \mathcal{O}(U)$, the composition

$$\theta^{-1}(U) \xrightarrow{\theta} U \xrightarrow{f} K$$

is in $\mathcal{O}(\theta^{-1}(U))$. In this way one gets a category of spaces with functions.

Properties. (1) If X is a space with functions and $V \subseteq X$ is any subset, one defines $\mathcal{O}(V)$ to be the set of functions $f : V \rightarrow K$ such that each $v \in V$ has an open neighbourhood U in X such that $f|_{V \cap U} = g|_{V \cap U}$ for some $g \in \mathcal{O}(U)$.

(2) Any subset V of a space with functions X becomes a space with functions with the induced topology and induced functions, and the inclusion $V \rightarrow X$ is a morphism. (Kempf, Exercise 1.5.3.)

(3) An *embedding* is a morphism $\theta : X \rightarrow Y$ which induces an isomorphism $X \rightarrow \text{Im}(\theta)$. If so, then for any Z is a space with functions, a mapping $\phi : Z \rightarrow X$ is a morphism if and only if $\theta\phi : Z \rightarrow Y$ is a morphism.

(4) If X and Y are spaces with functions, then the set $X \times Y$ can be given the structure of a space with functions, so that it becomes a product of X and Y in the category of spaces with functions. See Kempf, Lemma 3.1.1. The topology is not the usual product topology. Instead a basis of open sets is given by the sets

$$\{(u, v) \in U \times V : f(u, v) \neq 0\}$$

where U is open in X , V is open in Y and $f(x, y) = \sum_{i=1}^n g_i(x)h_i(y)$ with $g_i \in \mathcal{O}(U)$ and $h_i \in \mathcal{O}(V)$.

(5) The diagonal map $X \rightarrow X \times X$ is an embedding, since if Δ_X is its image, then there is a morphism $\Delta_X \rightarrow X$ given by the composition $\Delta_X \rightarrow X \times X \xrightarrow{p_1} X$.

(6) The projection morphism $p : X \times Y \rightarrow X$ is an *open morphism*, that is, the image of any open set U is open. Namely, for $y \in Y$, the identity morphism $X \rightarrow X$ and the morphism $X \rightarrow Y$ sending every element to y induce a morphism $i_y : X \rightarrow X \times Y$ with $i_y(x) = (x, y)$. Now if $U \subseteq X \times Y$, then $p(U) = \bigcup_{y \in Y} i_y^{-1}(U)$, which is open.

Definition. *Affine n -space* is $\mathbb{A}^n = K^n$ considered as a space with functions with:

- The topology is the *Zariski topology*, so closed sets are of the form

$$V(S) = \{(x_1, \dots, x_n) \in K^n \mid f(x_1, \dots, x_n) = 0 \text{ for all } f \in S\}$$

where S is a subset of the polynomial ring $K[X_1, \dots, X_n]$. Equivalently, the sets

$$D(f) = \{(x_1, \dots, x_n) \in K^n \mid f(x_1, \dots, x_n) \neq 0\}$$

with $f \in K[X_1, \dots, X_n]$ are a base of open subsets, and by noetherianness, any open set is a finite union of $D(f)$.

- If U is an open subset of $\mathbb{A}^n$, then the set of regular functions $\mathcal{O}(U)$ consists of the functions $f : U \rightarrow K$ such that each point $u \in U$ has an open neighbourhood $W \subseteq U$ such that $f|_W = p/q$ with $p, q \in K[X_1, \dots, X_n]$ and $q(x_1, \dots, x_n) \neq 0$ for all $(x_1, \dots, x_n) \in W$.

Properties. (a) If X is a space with functions, then a mapping

$$\theta : X \rightarrow \mathbb{A}^n, \quad \theta(x) = (\theta_1(x), \dots, \theta_n(x))$$

is a morphism of spaces with functions iff the θ_i are regular functions on X . If θ is a morphism, then since the i th projection $\pi_i : \mathbb{A}^n \rightarrow K$ is regular, so is $\theta_i = \pi_i \theta$. Conversely suppose that $\theta_1, \dots, \theta_n$ are regular. Let U be an open subset of $\mathbb{A}^n$ and $f = p/q \in \mathcal{O}(U)$ with $q(u) \neq 0$ for $u \in U$. We need to show that $f\theta$ is regular on $\theta^{-1}(U)$. Now by assumption $p\theta = p(\theta_1(x), \dots, \theta_n(x))$ and $q\theta$ are regular on U . Also $q\theta$ is non-vanishing on $\theta^{-1}(U)$. Thus $p\theta/q\theta$ is regular on $\theta^{-1}(U)$.

(b) It follows that $\mathbb{A}^n \times \mathbb{A}^m \cong \mathbb{A}^{n+m}$.

(c) An n -dimensional vector space V can be considered as a space with functions isomorphic to $\mathbb{A}^n$ by choosing any basis. Any linear map $\mathbb{A}^n \rightarrow \mathbb{A}^m$ is a morphism of spaces with functions, and an invertible linear map is an isomorphism, so a different basis gives the same space with functions.

(d) $X = \mathbb{A}^n$ is *separated*, meaning that the diagonal

$$\Delta_X = \{(x, x) : x \in X\}$$

is closed in $X \times X$, since

$$\Delta_{\mathbb{A}^n} = \{(x_1, \dots, x_n, y_1, \dots, y_n) \in \mathbb{A}^{2n} : x_1 = y_1, \dots, x_n = y_n\},$$

so it is closed. Note that if X is a topological space and $X \times X$ is considered with the product topology, then Δ_X is closed if and only if X is Hausdorff.

Definition. An *affine variety* is a space with functions which is, or is isomorphic to, a closed subset of $\mathbb{A}^n$. If X is an affine variety, its *coordinate ring* is $K[X] := \mathcal{O}(X)$. An (abstract) *variety* is a space with functions X which is *separated* and with a finite open covering by affine varieties.

Note that an affine variety is a variety, since separatedness passes to subsets of a space with functions equipped with the induced structure, for if Y is a subset of X , then $\Delta_Y = (Y \times Y) \cap \Delta_X$ in $X \times X$.

Example. *Determinantal varieties.* If V and W are f.d. vector spaces then the space $\text{Hom}(V, W)_{\leq r}$ of linear maps of rank $\leq r$ is closed in $\text{Hom}(V, W)$, so an affine variety. Namely, identifying this with $M_{n \times m}(K)$, it is defined by the vanishing of all minors of size $r + 1$.

Recall that the *radical* of an ideal I in a commutative ring A is

$$\sqrt{I} = \{a \in A : a^n \in I \text{ for some } n > 0\}$$

It is an ideal. The ideal I is *radical* if $I = \sqrt{I}$. Equivalently, if the factor ring A/I is *reduced*, that is, it has no nonzero nilpotent elements. Since $K[X_1, \dots, X_n]$ is a UFD, if f is an irreducible polynomial in $K[X_1, \dots, X_n]$, then (f) is a prime ideal, so $K[X_1, \dots, X_n]/(f)$ is a domain, so (f) is a radical ideal.

Theorem. Let X be a closed subset of $\mathbb{A}^n$, say $X = V(S)$ with S is a subset of $K[X_1, \dots, X_n]$. Then there is a canonical isomorphism

$$K[X] \cong K[X_1, \dots, X_n]/\sqrt{I}$$

where I is the ideal generated by S and $\sqrt{I}$ is its radical.

The kernel of the canonical map $K[X_1, \dots, X_n] \rightarrow K[X]$ is $\sqrt{I}$ by Hilbert's Nullstellensatz. The fact that it is surjective is proved in Kempf §1.5.

Corollary. The assignment $X \mapsto K[X]$ gives an anti-equivalence between the categories of affine varieties and finitely generated reduced commutative K -algebras. Moreover if Z is any space with functions, we get a bijection

$$\text{Hom}_{\text{spaces with functions}}(Z, X) \rightarrow \text{Hom}_{K\text{-algebras}}(K[X], \mathcal{O}(Z))$$

sending $\theta : Z \rightarrow X$ to the composition map $f \mapsto f\theta$.

Proof. The theorem shows that $K[X]$ is a f.g. reduced commutative algebra, and any such occurs. The statement about morphisms follows from our observation about morphisms $Z \rightarrow \mathbb{A}^n$. $\square$

Theorem. A product $X \times Y$ of varieties is a variety. If X and Y are affine varieties, so is $X \times Y$, and $K[X \times Y] \cong K[X] \otimes_K K[Y]$.

Proof. Recall that the product $X \times Y$ exists for any two spaces with functions.

If X is closed in $\mathbb{A}^n$ and Y is closed in $\mathbb{A}^m$ then $X \times Y$ is closed in $\mathbb{A}^n \times \mathbb{A}^m \cong \mathbb{A}^{n+m}$, so affine. Clearly $K[X] \otimes_K K[Y]$ is a f.g. commutative algebra, and with commutative algebra (using that K is algebraically closed) one can show it is reduced. Now the categorical property shows that $X \times Y$ has coordinate ring $K[X] \otimes_K K[Y]$.

In general, it is straightforward that if $U \subseteq X$ and $V \subseteq Y$ are open (resp. closed) subsets, then $U \times V$ is open (resp. closed) in $X \times Y$. Moreover with the induced structure as a space with functions it is a categorical product.

Assuming that X and Y are separated, $\Delta_{X \times Y}$ is identified with $\Delta_X \times \Delta_Y$ which is closed in $(X \times X) \times (Y \times Y)$. $\square$

Properties. (i) If X is a variety and $x \in X$, then the singleton set $\{x\}$ is closed in X . This is easy to see for affine space, it follows immediately for X an affine variety, and then for X an arbitrary variety.

(ii) Any variety is a *noetherian* topological space, that is it has the ascending chain condition on open subsets. The noetherian property of polynomial rings proves this for affine space, and then it follows for affine varieties and then for arbitrary varieties.

(iii) In particular, any variety is *quasi-compact*, meaning that any open covering has a finite subcovering. (Usually this is just called compactness, but in this context it is called quasi-compactness, apparently to make clear that the topological spaces needn't be Hausdorff.)

(iv) For a subset Y of a topological space, the following are equivalent, and then Y is called *locally closed*.

- (1) Y is an open subset of a closed subset of X
- (2) Y is open in its closure
- (3) Y is the intersection of an open and a closed subset of X .

Definition. A *subvariety* Y of a variety X is a locally closed subset equipped with the induced structure as a space with functions.

Clearly a closed subvariety of an affine variety is affine.

Proposition. If X is an affine variety and $f \in K[X]$, then the open subset $D(f) = \{x \in X : f(x) \neq 0\}$ is an affine variety and $K[D(f)] \cong K[X]_f$ (the localization, inverting f).

Proof. We have an isomorphism $D(f) \xrightarrow{\sim} \{(x, t) \in X \times \mathbb{A}^1 : f(x)t = 1\}$, where the map to the left sends (x, t) to x and the map to the right sends x to $(x, 1/f(x))$. It is a morphism since $1/f \in \mathcal{O}(D(f))$. $\square$

Corollary. *Any subvariety of a variety is a variety.*

Proof. Suppose $Y \subseteq X$. We need to show that Y is a finite union of affine open subsets. Since X is a finite union of affine opens, we may reduce to the case when X is affine. We may also assume that Y is open in X . But then $Y = X \cap U$ with $U = D(f_1) \cup \dots \cup D(f_m)$, and then $Y = V_1 \cup \dots \cup V_m$ with $V_i = X \cap D(f_i)$ a closed subset of the affine variety $D(f_i)$, hence affine. $\square$

Example. If V and W are vector spaces, the set of injective linear maps $\text{Inj}(V, W)$ is an open subvariety in $\text{Hom}(V, W)$, since the complement is $\text{Hom}_{\leq r}(V, W)$ where $r = \dim V - 1$.

Remark. The example of $D(f)$ shows that some open subvarieties of affine varieties quasi-affine varieties are again affine. But this is not always true. For example $U = \mathbb{A}^2 \setminus \{0\} = D(X_1) \cup D(X_2)$ is not affine.

To see this, we show first that $\mathcal{O}(U) = K[X_1, X_2]$. A function $f \in \mathcal{O}(U)$ is determined by its restrictions f_i to $D(X_i)$ ($i = 1, 2$). Now $f_i \in \mathcal{O}(D(X_i)) = K[X_1, X_2, X_i^{-1}]$. Moreover the restrictions of f_1 and f_2 to $D(X_1) \cap D(X_2) = D(X_1 X_2)$ are equal, so f_1 and f_2 are equal as elements of $K[X_1, X_2, 1/X_1 X_2]$. But this is only possible if they are both in $K[X_1, X_2]$, and equal. Thus $f \in K[X_1, X_2]$.

Now the inclusion morphism $\theta : U \rightarrow \mathbb{A}^2$ induces a homomorphism $\mathcal{O}(\mathbb{A}^2) \rightarrow \mathcal{O}(U)$ which is actually an isomorphism. Now the category of affine varieties is anti-equivalent to the category of finitely generated reduced K -algebras. If U were affine, then since the map on coordinate rings is an isomorphism, θ would have to be an isomorphism. But it isn't.

Definition. A (non-empty) topological space X is *irreducible* if it cannot be written as a union of two proper closed subsets.

Properties. (1) X is irreducible iff every non-empty open subset U is dense in X . Thus any non-empty open subset of an irreducible space is irreducible.

(2) An affine variety X is irreducible iff $K[X]$ is a domain. (Kempf, Lemma 2.3.1.) In particular $\mathbb{A}^n$ is irreducible.

(3) Any variety is a finite union of maximal irreducible closed subvarieties, its *irreducible components*.

(4) A product of irreducible varieties is irreducible. Indeed if $X \times Y = Z_1 \cup Z_2$ with the Z_i closed, then for all $x \in X$ we have

$$Y = i_x^{-1}(Z_1) \cup i_x^{-1}(Z_2),$$

so by irreducibility, one of the sets on the right is Y . Thus $\{x\} \times Y$ is contained in Z_1 or Z_2 . Thus $X = X_1 \cup X_2$ where

$$X_i = \bigcap_{y \in Y} i_y^{-1}(Z_i)$$

Thus by irreducibility, we have $X = X_i$ for some i , so $Z_i = X \times Y$.

Definition. An *algebraic group* is a group which is also a variety, such that multiplication $G \times G \rightarrow G$ and inversion $G \rightarrow G$ are morphisms of varieties.

A morphism of algebraic groups is a map which is a group homomorphism and a morphism of varieties.

When considering an action of an algebraic group on a variety X we shall suppose that the map $G \times X \rightarrow X$ is a morphism of varieties.

The *general linear group* $\mathrm{GL}_n(K)$ is the open subset $D(\det)$ of $M_n(K)$, so an affine variety. It is an algebraic group thanks to the formula $g^{-1} = \mathrm{adj} g / \det g$. It acts by left multiplication or by conjugation on $M_n(K)$.

A *linear algebraic group* is an algebraic group which is isomorphic to a closed subgroup of $\mathrm{GL}_n(K)$. For example the special linear group, orthogonal group or any finite group. The additive and multiplicative groups of the field are

$$G_a = (K, +) \cong \left\{ \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} : a \in K \right\}, \quad G_m = (K \setminus \{0\}, \times) = \mathrm{GL}_1(K).$$

Any finite product of linear algebraic groups is a linear algebraic group, using that $\mathrm{GL}_n(K) \times \mathrm{GL}_m(K)$ embeds in $\mathrm{GL}_{n+m}(K)$.

Remark. Any linear algebraic group is an affine variety, and conversely one can show that any affine algebraic group is linear, see for example Humphreys, Linear algebraic groups, section 8.6. An elliptic curve is an example of an algebraic group which is a projective variety, so not linear.

Lemma. *A connected algebraic group is an irreducible variety.*

Proof. Write the group as a union of irreducible components $G = G_1 \cup \cdots \cup G_n$. Since G_1 is not a subset of the union of the other components, some element $g \in G_1$ does not lie in any other component. Now any two elements of an algebraic group look the same, since multiplication by any $h \in G$ defines an isomorphism $G \rightarrow G$. It follows that every element of G lies in only one irreducible component. Thus G is the disjoint union of its irreducible components. But then the components are open and closed, and since G is connected, there is only one component. $\square$

Let V be a vector space of dimension n , with basis $e_1, \dots, e_n$. We write $\text{Bil}(n)$ for the set of bilinear maps $V \times V \rightarrow V$. A map $\mu \in \text{Bil}(n)$ is given by its structure constants $(c_{ij}^k) \in K^{n^3}$ with

$$\mu(e_i, e_j) = \sum_k c_{ij}^k e_k.$$

Equivalently $\text{Bil}(n) \cong \text{Hom}(V \otimes V, V)$, Thus it is affine space $\mathbb{A}^{n^3}$.

We write $\text{Ass}(n)$ for the subset consisting of associative multiplications. This is a closed subset of $\text{Bil}(n)$, hence an affine variety, since it is defined by the equations

$$\mu(\mu(e_i, e_j), e_k) = \mu(e_i, \mu(e_j, e_k)),$$

that is

$$\sum_p c_{ij}^p c_{pk}^s = \sum_q c_{iq}^s c_{jk}^q$$

for all s .

We write $\text{Alg}(n)$ for the subset of associative unital multiplications, so algebra structures on V .

Theorem. *$\text{Alg}(n)$ is an affine open subset of $\text{Ass}(n)$, hence an affine variety. The algebraic group $\text{GL}(V)$ acts by basis change, and the orbits correspond to isomorphism classes of n -dimensional algebras.*

Proof. (i) We use that a vector space A with an associative multiplication has a 1 if and only if there is some $a \in A$ for which the maps $\ell_a, r_a : A \rightarrow A$ of left and right multiplication by a are invertible.

Namely, if $u = \ell_a^{-1}(a)$, then $au = a$. Thus $aub = ab$ for all b , so since ℓ_a is invertible, $ub = b$. Thus u is a left 1. Similarly there is a right 1, and they must be equal.

(ii) For the algebra V with multiplication μ , write ℓ_a^μ and r_a^μ for left and right multiplication by $a \in V$. Then $\text{Alg}(n) = \bigcup_{a \in V} D(f_a)$ where $f_a(\mu) = \det(\ell_a^\mu) \det(r_a^\mu)$. Thus $\text{Alg}(n)$ is open in $\text{Ass}(n)$.

(iii) The map

$$\text{Alg}(n) \rightarrow V, \quad \mu \mapsto \text{the } 1 \text{ for } \mu$$

is a morphism of varieties, since on $D(f_a)$ it is given by $(\ell_a^\mu)^{-1}(a)$, whose components are rational functions, with $\det(\ell_a^\mu)$ in the denominator.

(iv) $\text{Alg}(n)$ is affine. In fact

$$\text{Alg}(n) \cong \{(\mu, u) \in \text{Ass}(n) \times V \mid u \text{ is a } 1 \text{ for } \mu\}.$$

The right hand side is a closed subset, hence it is affine. Certainly there is a bijection, and the maps both ways are morphisms.

(v) Last statement is clear. □

Example. The structure of $\text{Alg}(n)$ is known for small n . For example $\text{Alg}(4)$ has 5 irreducible components, of dimensions 15, 13, 12, 12, 9. See P. Gabriel, Finite representation type is open, 1974.

END OF LECTURE ON 2025-11-20. PROVISIONAL SCRIPT FOR THE NEXT LECTURE FOLLOWS (SUBJECT TO CHANGE).

6.2 Schemes and varieties of modules

More general than varieties are schemes. I only discuss affine schemes, using representable functors rather than sheaves. See:

- M. Demazure and P. Gabriel, Groupes Algébriques, 1970. Partial English translation, Introduction to Algebraic Geometry and Algebraic Groups, 1980.
- W. C. Waterhouse, Affine group schemes, 1979.
- D. Eisenbud and J. Harris, The geometry of schemes, 2000. (Chapter VI)

Let K be a commutative ring. We write $K\text{-comm}$ for the category of commutative K -algebras, or equivalently commutative rings R equipped with a homomorphism $K \rightarrow R$.

Definition. The category of *affine (K -)schemes* is the category of representable (covariant) functors

$$F : K\text{-comm} \rightarrow \text{Sets}$$

with morphisms given by natural transformations. (These are not additive categories.)

Recall that a functor F is said to be *representable* if there is an object A in the category (a commutative K -algebra) such that

$$F(-) \cong \text{Hom}_{K\text{-comm}}(A, -).$$

By Yoneda's lemma, the functor $A \mapsto \text{Hom}_{K\text{-comm}}(A, -)$ defines an anti-equivalence from $K\text{-comm}$ to the category of affine schemes.

Examples. (i) $\mathbf{A}^n$ is the affine scheme with $\mathbf{A}^n(R) = R^n$. It is represented by the polynomial ring $K[X_1, \dots, X_n]$, since

$$\text{Hom}_{K\text{-comm}}(K[X_1, \dots, X_n], R) = R^n.$$

(ii) Any subset S of $K[X_1, \dots, X_n]$ defines a functor $\mathbf{V}(S)$ by

$$\mathbf{V}(S)(R) = \{(x_1, \dots, x_n) \in R^n : f(x_1, \dots, x_n) = 0 \text{ for all } f \in S\}.$$

It is an affine scheme, represented by the algebra $K[X_1, \dots, X_n]/(S)$.

Definition. The affine scheme represented by A is

- *algebraic* if A is f.g. as a K -algebra (and K is a noetherian ring).
- *reduced* if A is reduced.

We immediately get:

Proposition. *If K is an algebraically closed field, then there is an equivalence*

Cat. of affine varieties $\rightarrow$ Cat. of reduced affine algebraic schemes

$$X \mapsto \text{Hom}_{K\text{-comm}}(K[X], -).$$

We usually identify an affine variety with the reduced affine algebraic scheme. Note that if X is the functor, then the underlying set for the variety is $X(K)$.

Lemma. *Given an affine (algebraic) scheme F , there is a reduced affine (algebraic) scheme F_{red} and a morphism $F_{\text{red}} \rightarrow F$ such that for all R the map*

$$F_{\text{red}}(R) \rightarrow F(R)$$

is injective, and a bijection for R reduced. This defines a functor $F \mapsto F_{\text{red}}$ which is right adjoint to the inclusion of reduced affine (algebraic) schemes into affine (algebraic) schemes.

Proof. If $F(-) = \text{Hom}(A, -)$ we set $F_{\text{red}}(-) = \text{Hom}(A_{\text{red}}, -)$. The natural map $A \rightarrow A_{\text{red}}$ gives a morphism $F_{\text{red}} \rightarrow F$. $\square$

For example $\mathbf{V}(S)$ is algebraic. It is reduced if and only if $K[X_1, \dots, X_n]/(S)$ is reduced. The scheme $\mathbf{V}(S)_{\text{red}}$ is represented by $K[X_1, \dots, X_n]/\sqrt{(S)}$

Remark. If K is any commutative ring, then an *affine group scheme* over K is a representable functor $F : K\text{-comm} \rightarrow \text{Groups}$. If A is the commutative K -algebra representing F , then A becomes a Hopf algebra, see Waterhouse §1.4. For example $\mathbf{GL}_n$ is the affine group scheme with $\mathbf{GL}_n(R) = \text{GL}_n(R)$ for all R . It is represented by the algebra $K[X_{ij}, 1/\det]$, so reduced.

Let A be a f.g. K -algebra (possibly non-commutative). A d -dimensional A -module V can be considered as a homomorphism $A \rightarrow \text{End}_K(V)$, or choosing a basis of V , as a homomorphism $A \rightarrow M_d(K)$.

Definition. Let A be a f.g. K -algebra and n a natural number. We define the scheme $\mathbf{Mod}(A, d)$ of d -dimensional A -modules to be the functor

$$K\text{-comm} \rightarrow \text{Sets}, \quad R \mapsto \text{Hom}_{K\text{-algebra}}(A, M_d(R)).$$

Lemma. $\mathbf{Mod}(A, d)$ is an affine algebraic K -scheme.

Proof. Write A as a quotient of a free algebra, say $A = K\langle X_1, \dots, X_k \rangle / I$. Then

$$\mathbf{Mod}(A, d)(R) = \{(A_1, \dots, A_k) \in M_d(R)^k : p(A_1, \dots, A_k) = 0 \text{ for all } p \in I\}.$$

Consider the polynomial ring $S = K[X_{rij} : 1 \leq r \leq k, 1 \leq i, j \leq d]$ and let $U_r \in M_d(S)$ be the matrix with (i, j) entry X_{rij} . If $p \in K\langle X_1, \dots, X_k \rangle$, then considering it as a noncommutative polynomial, we obtain $p(U_1, \dots, U_k) \in M_d(S)$. Then $\mathbf{Mod}(A, d)(R)$ is in bijection with

$$\mathrm{Hom}_{K\text{-algebra}}(S/J, M_d(R)).$$

where J is the ideal generated by all entries of $p(U_1, \dots, U_k)$ with $p \in I$. $\square$

Definition. If K is algebraically closed, the variety corresponding to the reduced scheme is denoted $\mathrm{Mod}(A, d)$. Thus

$$\mathrm{Mod}(A, d) = \mathrm{Hom}_{K\text{-algebra}}(A, M_d(K)).$$

There is an action of $\mathrm{GL}_d(K)$ on $\mathrm{Mod}(A, d)$ by conjugation, so given by $(g \cdot \theta)(a) = g\theta(a)g^{-1}$. The orbits correspond to isomorphism classes of d -dimensional modules.

Writing $A = K\langle X_1, \dots, X_k \rangle / I$, we have

$$\mathrm{Mod}(A, d) = \{(A_1, \dots, A_k) \in M_d(K)^k : p(A_1, \dots, A_k) = 0 \text{ for all } p \in I\}.$$

Examples. (1) $\mathbf{Mod}(A, 1)$ is the affine algebraic scheme given by the targets commutative quotient of A , which is $A/([A, A])$, where $[A, A] = \{ab - ba : a, b \in A\}$. Then the variety has coordinate ring

$$K[\mathrm{Mod}(A, 1)] = (A/([A, A]))/\sqrt{0}.$$

(2) The *nilpotent variety* consists of the $d \times d$ nilpotent matrices over K . In fact that d th power of such a matrix must be zero, so the nilpotent variety is

$$N_d = \{A \in M_d(K) : A^d = 0\} = \mathrm{Mod}(K[x]/(x^d), d)$$

(3) The *commuting variety* consists of the pairs of commuting matrices

$$C_d = \{(A, B) \in M_d(K)^2 : AB = BA\} = \mathrm{Mod}(K[x, y], d).$$

Remark. We can do the same thing with quivers and dimension vectors. If Q is a finite quiver and $\alpha \in \mathbb{N}^{Q_0}$ is a dimension vector, one gets an affine space

$$\text{Rep}(Q, \alpha) = \prod_{a \in Q_1} \text{Hom}_K(K^{\alpha_{t(a)}}, K^{\alpha_{h(a)}})$$

and the group

$$\text{GL}(\alpha) = \prod_{i \in Q_0} \text{GL}_{\alpha_i}(K)$$

acts by conjugation,

$$(g \cdot x)_a = g_{h(a)} x_a (g_{t(a)})^{-1}$$

and the orbits correspond to the isomorphism classes of representations of Q of dimension vector α .

There is a corresponding scheme $\mathbf{Rep}(Q, \alpha)$ with

$$\mathbf{Rep}(Q, \alpha)(R) = \prod_{a: i \rightarrow j} \text{Hom}_R(R^{\alpha_i}, R^{\alpha_j})$$

which is reduced.

For an algebra $A = KQ/I$, one gets an affine algebraic scheme $\mathbf{Rep}(A, \alpha)$ which is not necessarily reduced, and the affine variety $\text{Rep}(A, \alpha)$ which is a closed subset of $\text{Rep}(Q, \alpha)$. Again, the orbits under $\text{GL}(\alpha)$ correspond to the isomorphism classes of representations.

Another way to get this. Let $S = K \times \cdots \times K$ with the factors indexed by Q_0 . There is a homomorphism $S \rightarrow A$ sending the 1 in the i th factor to the idempotent e_i , and restriction gives a morphism

$$\text{res} : \text{Mod}(A, d) \rightarrow \text{Mod}(S, d).$$

Given a dimension vector α with $\sum_{i \in Q_0} \alpha_i = d$, let $z \in \text{Mod}(S, d)$ be the S -module structure given by the decomposition

$$K^d = \bigoplus_{i \in Q_0} K^{\alpha_i}.$$

Then $\text{Rep}(A, \alpha) = \text{res}^{-1}(z)$.

Example. Let Q be the quiver $1 \xrightarrow{a} 2 \xrightarrow{b} 3$ and I the ideal generated by ba .

$$\text{Rep}(KQ/I, (2, 2, 1)) = \{(a, b) \in M_{2 \times 2}(K) \times M_{1 \times 2}(K) : ba = 0\}.$$